

Essential Haematology

FIFTH EDITION

Essential Haematology

EDITED BY

A.V. Hoffbrand

MA DM FRCP FRCPATH FRCP(Edin) DSc FMED Sci
Emeritus Professor of Haematology, Royal Free
and University College Medical School, London

P.A.H. Moss

PhD FRCP FRCPATH
Professor of Haematology,
University of Birmingham

J.E. Pettit

MD FRCPA FRCPATH
Haematologist, Medlab South,
Christchurch, New Zealand



Blackwell
Publishing

© 1980, 1984, 1993, 2001, 2006 AV Hoffbrand, PAH Moss and JE Pettit
Published by Blackwell Publishing Ltd
Blackwell Publishing, Inc., 350 Main Street, Malden, Massachusetts 02148-5020, USA
Blackwell Publishing Ltd, 9600 Garsington Road, Oxford OX4 2DQ, UK
Blackwell Publishing Asia Pty Ltd, 550 Swanston Street, Carlton, Victoria 3053, Australia

The right of the Author to be identified as the Author of this Work has been asserted in accordance with the Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

First published 1980
Reprinted 1981, 1982, 1983 (twice)
Second edition 1984
Reprinted 1985
Reprinted with corrections 1985, 1988 (twice), 1989
German edition 1986 (reprinted 1996)
Japanese edition 1986
Spanish edition 1987 (reprinted twice)
Indonesian edition 1987
Third edition 1993
Hungarian edition 1997
Chinese edition 1998
Reprinted with corrections 1993, 1994, 1995, 1996, 1997, 1998, 1999, 2000
Fourth edition 2001
Fifth edition 2006

12006

Library of Congress Cataloging-in-Publication Data

Hoffbrand, A.V.
Essential haematology / A.V. Hoffbrand, P.A.H. Moss, and J.E. Pettit. — 5th ed.
p. ; cm.
Includes bibliographical references.
ISBN-13: 978-1-4051-3649-5 (alk. paper)
ISBN-10: 1-4051-3649-9 (alk. paper)
1. Blood—Diseases. 2. Hematology. I. Moss, P.A.H. II. Pettit, J.E. III. Title.
[DNLM: 1. Hematologic Diseases. WH 120 H698e 2006]
RC633.H627 2006
616.1'5—dc22

2006016791

ISBN-13: 978-1-4051-3649-5
ISBN-10: 1-4051-3649-9

A catalogue record for this title is available from the British Library

Set in 9/12pt Palatino by Graphicraft Limited, Hong Kong
Printed and bound by Rotolito Lombarda SpA, Italy

Commissioning Editor: Martin Sugden
Development Editor: Rebecca Huxley
Production Controller: Kate Charman

For further information on Blackwell Publishing, visit our website:
<http://www.blackwellpublishing.com>

The publisher's policy is to use permanent paper from mills that operate a sustainable forestry policy, and which has been manufactured from pulp processed using acid-free and elementary chlorine-free practices. Furthermore, the publisher ensures that the text paper and cover board used have met acceptable environmental accreditation standards.

Blackwell Publishing makes no representation, express or implied, that the drug dosages in this book are correct. Readers must therefore always check that any product mentioned in this publication is used in accordance with the prescribing information prepared by the manufacturers. The author and the publishers do not accept responsibility or legal liability for any errors in the text or for the misuse or misapplication of material in this book.

Contents

- Preface to fifth edition, vii
- Preface to first edition, viii
- Bibliography, ix
- 1 Haemopoiesis, 1
- 2 Erythropoiesis and general aspects of anaemia, 12
- 3 Hypochromic anaemias and iron overload, 28
- 4 Megaloblastic anaemias and other macrocytic anaemias, 44
- 5 Haemolytic anaemias, 58
- 6 Genetic disorders of haemoglobin, 72
- 7 The white cells 1: granulocytes, monocytes and their benign disorders, 94
- 8 The white cells 2: lymphocytes and their benign disorders, 108
- 9 The spleen, 123
- 10 The aetiology and genetics of haematological malignancies, 129
- 11 Management of haematological malignancy, 147
- 12 Acute leukaemias, 157
- 13 Chronic myeloid leukaemia, 174
- 14 Myelodysplasia, 182
- 15 The chronic lymphoid leukaemias, 188
- 16 Hodgkin's lymphoma, 197
- 17 Non-Hodgkin's lymphoma, 203
- 18 Multiple myeloma and related disorders, 216
- 19 Myeloproliferative disorders, 230

- 20 Aplastic anaemia and bone marrow failure, 241
- 21 Stem cell transplantation, 249
- 22 Platelets, blood coagulation and haemostasis, 264
- 23 Bleeding disorders caused by vascular and platelet abnormalities, 278
- 24 Coagulation disorders, 290
- 25 Thrombosis and antithrombotic therapy, 303
- 26 Haematological changes in systemic disease, 320
- 27 Blood transfusion, 337
- 28 Pregnancy and neonatal haematology, 352

Appendices

- 1 Principal features of known cluster differentiation (CD) molecules, 360
- 2 Normal values, 365
- 3 World Health Organization (WHO) classification of myeloid and lymphoid neoplasms, 366

Index, 370

Preface to the fifth edition

Major advances in the understanding at a molecular level of the genetic changes underlying many inherited and acquired haematological disorders have necessitated revisions in many chapters for this new edition. The introduction of new diagnostic laboratory tests and imaging techniques for diagnosis and monitoring of malignant blood disorders and the major changes in their therapy has required expansion of these sections with many new figures. Chapters on principles of treatment and support care and on the spleen have been added while Hodgkin's disease and the non-Hodgkin's lymphomas are now given separate chapters. The greatly increased knowledge of iron absorption, metabolism and iron loading diseases is recognized by allocating increased space to these topics. In order to keep the book as concise as possible, topics that are now no longer in routine practice have been omitted (e.g. radioactive chromium red cell survival studies and immunofluorescent microscopy).

Essential Haematology is intended for medical students but we realize there is more information than most medical students can be expected to know, given the new knowledge in all areas of medicine. We have therefore indicated by means of vertical blue lines in the margins, material that we

consider more suitable for honours candidates and for postgraduates, leaving unmarked core material appropriate for medical students taking final examinations. We hope, therefore, that this new edition provides 'two books', one more basic, the other more advanced. We hope the book will be used as previously, by science graduates, medical laboratory technicians and general physicians, indeed all those wishing to learn about the exciting topic of the blood and its diseases.

As previously, we are grateful to Rebecca Huxley for her expert help during the publishing process and to Jane Fallows for the beautiful clear scientific diagrams which illustrate the text throughout. We are also grateful to Dr Clare Taylor for advice in Chapters 27 and 28, and Dr Pratima Chowdary for advice on Chapters 22–25. We also wish to thank Blackwell Publishing for their continuing support for this book since it was first published in 1980.

Sir John Dacie, FRS was mentor to AVH and JEP, and Per Saugman, Chairman of Blackwell Scientific Publications commissioned the First Edition of *Essential Haematology*. Both sadly died during the last year. We wish to dedicate this fifth edition of *Essential Haematology* to their memory.

A.V. Hoffbrand, P.A.H. Moss and J.E. Pettit
September 2006

Preface to the first edition

The major changes that have occurred in all fields of medicine over the last decade have been accompanied by an increased understanding of the biochemical, physiological and immunological processes involved in normal blood cell formation and function and the disturbances that may occur in different diseases. At the same time, the range of treatment available for patients with diseases of the blood and blood-forming organs has widened and improved substantially as understanding of the disease processes has increased and new drugs and means of support care have been introduced.

We hope the present book will enable the medical student of the 1980s to grasp the essential features of modern clinical and laboratory haematology and to achieve an understanding of how many of the manifestations of blood diseases can be explained with this new knowledge of the disease processes.

We would like to thank many colleagues and assistants who have helped with the preparation of the book. In particular, Dr H.G. Prentice cared for the patients whose haematological responses are illustrated in Figs 5.3 and 7.8 and Dr J. McLaughlin supplied Fig. 8.6. Dr S. Knowles reviewed critically the final manuscript and made many helpful suggestions. Any remaining errors are, however, our own. We also thank Mr J.B. Irwin and R.W. McPhee who drew many excellent diagrams, Mr Cedric Gilson for expert photomicrography, Mrs T. Charalambos, Mrs B. Elliot, Mrs M. Evans and Miss J. Allaway for typing the manuscript, and Mr Tony Russell of Blackwell Scientific Publications for his invaluable help and patience.

AVH, JEP

Bibliography

- Arceci R.J., Hann I.M., Smith O.P. (eds) (2006) *Pediatric Hematology*, 3rd edn. Blackwell Publishing Ltd, Oxford.
- Bain B.J. (2005) *A Beginner's Guide to Blood Cells*, 2nd edn. Blackwell Publishing, Oxford.
- Bain B.J., Clark D.M., Lampert I.A., Wilkins B.S. (2001) *Bone Marrow Pathology*, 2nd edn. Blackwell Science, Oxford.
- Beutler E., Kipps T.J., Lichtman M.A. *et al.* (eds) (2000) *William's Hematology*, 6th edn. McGraw-Hill, New York.
- Degos L., Linch D.C., Lowenberg B. (eds) (2005) *Textbook of Malignant Hematology*, 2nd edn. Taylor & Francis, London & New York.
- Hoffbrand A.V. and Pettit J.E. (2000) *Color Atlas of Clinical Hematology*, 3rd edn. Mosby, London.
- Hoffbrand A.V., Tuddenham E.G. and Catovsky D. (eds) (2005) *Postgraduate Hematology*, 5th edn. Blackwell Publishing, Oxford.
- Hoffman R., Benz E.J., Shatill S.J. *et al.* (2005) *Hematology: Basic Principles and Practice*, 4th edn. Elsevier, New York.
- Klein H.G. and Anstee D.J. (2005) *Mollison's Blood Transfusion in Clinical Medicine*, 11th edn. Blackwell Publishing, Oxford.
- Provan D. and Gribben J. (eds) (2005) *Molecular Haematology*, 2nd edn. Blackwell Publishing, Oxford.
- Stamatoyannopoulos G., Perlmutter R.M., Majerus P.W., Varmus H. (eds) (2000) *The Molecular Basis of Blood Diseases*, 3rd edn. W.B. Saunders, Philadelphia.

CHAPTER 1

1

Haemopoiesis

Site of haemopoiesis, 1
Haemopoietic stem and progenitor cells, 1
Bone marrow stroma, 3
Stem cell plasticity, 4
The regulation of haemopoiesis, 5
Haemopoietic growth factors, 5

Growth factor receptors and signal transduction, 6
The cell cycle, 7
Apoptosis, 9
Transcription factors, 10
Adhesion molecules, 11
Bibliography, 11

This first chapter is concerned with the general aspects of blood cell formation (haemopoiesis). The processes that regulate haemopoiesis and the early stages of formation of red cells (erythropoiesis), granulocytes and monocytes (myelopoiesis) and platelets (thrombopoiesis) are also discussed.

Site of haemopoiesis

In the first few weeks of gestation the yolk sac is the main site of haemopoiesis. However, definitive haemopoiesis derives from a population of stem cells first observed on the dorsal aorta termed the AGM (aorta-gonads-mesonephros) region. These common precursors of endothelial and haemopoietic cells (haemangioblasts) are believed to seed the liver, spleen and bone marrow and from 6 weeks until 6–7 months of fetal life the liver and spleen are

the major haemopoietic organs and continue to produce blood cells until about 2 weeks after birth (Table 1.1) (see Fig. 6.1b). The bone marrow is the most important site from 6 to 7 months of fetal life. During normal childhood and adult life the marrow is the only source of new blood cells. The developing cells are situated outside the bone marrow sinuses and mature cells are released into the sinus spaces, the marrow microcirculation and so into the general circulation.

In infancy all the bone marrow is haemopoietic but during childhood there is progressive fatty replacement of marrow throughout the long bones so that in adult life haemopoietic marrow is confined to the central skeleton and proximal ends of the femurs and humeri (Table 1.1). Even in these haemopoietic areas, approximately 50% of the marrow consists of fat (Fig. 1.1). The remaining fatty marrow is capable of reversion to haemopoiesis and in many diseases there is also expansion of haemopoiesis down the long bones. Moreover, the liver and spleen can resume their fetal haemopoietic role ('extramedullary haemopoiesis').

Table 1.1 Sites of haemopoiesis.

Fetus	0–2 months (yolk sac) 2–7 months (liver, spleen) 5–9 months (bone marrow)
Infants	Bone marrow (practically all bones)
Adults	Vertebrae, ribs, sternum, skull, sacrum and pelvis, proximal ends of femur

Haemopoietic stem and progenitor cells

Haemopoiesis starts with a pluripotential stem cell that can give rise to the separate cell lineages.




Fig. 1.1 A normal bone marrow trephine biopsy (posterior iliac crest). Haematoxylin and eosin stain; approximately 50% of the intertrabecular tissue is haemopoietic tissue and 50% is fat.

This *haemopoietic stem cell* is rare, perhaps 1 in every 20 million nucleated cells in bone marrow. Although its exact phenotype is unknown, on immunological testing it is $CD34^+ CD38^-$ and has the appearance of a small or medium-sized lymphocyte (Fig. 21.3). Cell differentiation occurs from the stem cell via the committed *haemopoietic progenitors* which are restricted in their developmental potential (Fig. 1.2). The existence of the separate progenitor cells can be demonstrated by *in vitro* culture techniques. Very early progenitors are assayed by culture on bone marrow stroma as long-term culture initiating cells whereas late progenitors are generally assayed in semi-solid media. An example is the earliest detectable mixed myeloid precursor which gives rise to granulocytes, erythrocytes, monocytes and megakaryocytes and is termed




Fig. 1.2 Diagrammatic representation of the bone marrow pluripotent stem cell and the cell lines that arise from it. Various progenitor cells can be identified by culture in semi-solid medium by the type of colony they form. Baso, basophil; BFU, burst-forming unit; CFU, colony-forming unit; E, erythroid; Eo, eosinophil; GEMM, granulocyte, erythroid, monocyte and megakaryocyte; GM, granulocyte, monocyte; Meg, megakaryocyte; NK, natural killer.




Fig. 1.3 (a) Bone marrow cells are increasingly differentiated and lose the capacity for self-renewal as they mature. (b) A single stem cell gives rise, after multiple cell divisions (shown by vertical lines), to $>10^6$ mature cells.

CFU (colony-forming unit)-GEMM (Fig. 1.2). The bone marrow is also the primary site of origin of lymphocytes (Chapter 8) which differentiate from a common lymphoid precursor.

The stem cell has the capability for *self-renewal* (Fig. 1.3) so that marrow cellularity remains constant in a normal healthy steady state. There is considerable amplification in the system: one stem cell is capable of producing about 10^6 mature blood cells after 20 cell divisions (Fig. 1.3). The precursor cells are, however, capable of responding to haemopoietic growth factors with increased production of one or

other cell line when the need arises. The development of the *mature cells* (red cells, granulocytes, monocytes, megakaryocytes and lymphocytes) is considered further in other sections of this book.

Bone marrow stroma

The bone marrow forms a suitable environment for stem cell survival, growth and development. It is composed of stromal cells and a microvascular network (Fig. 1.4). The stromal cells include adipocytes, fibroblasts, endothelial cells and macrophages and




Fig. 1.4 Haemopoiesis occurs in a suitable microenvironment provided by a stromal matrix on which stem cells grow and divide. There are probably specific recognition and adhesion sites (p. 11); extracellular glycoproteins and other compounds are involved in the binding.

they secrete extracellular molecules such as collagen, glycoproteins (fibronectin and thrombospondin) and glycosaminoglycans (hyaluronic acid and chondroitin derivatives) to form an extracellular matrix. In addition, stromal cells secrete several growth factors necessary for stem cell survival. *Mesenchymal stem cells* are thought to be critical in stromal cell formation.

Stem cells are able to traffic around the body and are found in peripheral blood in low numbers. In order to exit the bone marrow, cells must cross the blood vessel endothelium and this process of *mobilization* is enhanced by administration of cytokines such as granulocyte colony-stimulating factor (G-CSF) or granulocyte-macrophage colony-stimulating factor (GM-CSF) (p. 97). The reverse process of stem cell *homing* appears to depend on a chemokine gradient in which the stromal-derived factor (SDF-1) is critical. Several critical

interactions maintain stem cell viability and production in the stroma including stem cell factor (SCF) and Jagged proteins expressed on stroma and their respective receptors c-Kit and Notch expressed on stem cell.

Stem cell plasticity

There is some evidence that adult stem cells in different organs are *pluripotent* and can generate various types of tissue (Fig. 1.5). Studies in patients and animals who have received haemopoietic stem cell transplants (Chapter 21) have suggested that donor cells may contribute to tissues such as neurons, liver and muscle. The contribution of adult donor bone marrow cells to non-haemopoietic tissues is at most small. The persistence of pluripotent stem cells in postnatal life, organ-specific stem cells and fusion of transplanted cells with host cells




Fig. 1.5 (a) Cells in the early embryo are able to generate all the tissues of the body and are known as totipotent. (b) Specialized adult stem cells of the bone marrow, nervous tissue, epithelial and other tissues give rise to differentiated cells of the same tissue and possibly to other tissues (see text).

have all been proposed, however, to explain many of the findings suggesting stem cell plasticity.

The regulation of haemopoiesis

Haemopoiesis starts with stem cell division in which one cell replaces the stem cell (*self-renewal*) and the other is committed to differentiation. These early committed progenitors express low levels of transcription factors that may commit them to discrete cell lineages. Which cell lineage is selected for differentiation may depend both on chance and on the external signals received by progenitor cells. Several transcription factors have been isolated that regulate differentiation along the major cell lineages. For instance, PU.1 commits cells to the myeloid

lineage whereas GATA-1 has an essential role in erythropoietic and megakaryocytic differentiation.

Haemopoietic growth factors

The haemopoietic growth factors are glycoprotein hormones that regulate the proliferation and differentiation of haemopoietic progenitor cells and the function of mature blood cells. They may act locally at the site where they are produced by cell-cell contact or circulate in plasma. They also bind to the extracellular matrix to form niches to which stem and progenitor cells adhere. The growth factors may cause cell proliferation but can also stimulate differentiation, maturation, prevent apoptosis and affect the function of mature cells (Fig. 1.6).




Fig. 1.6 Growth factors may stimulate proliferation of early bone marrow cells, direct differentiation to one or other cell type, stimulate cell maturation, suppress apoptosis or affect the function of mature non-dividing cells, as illustrated here for granulocyte colony-stimulating factor (G-CSF) for an early myeloid progenitor and a neutrophil.

Table 1.2 General characteristics of myeloid and lymphoid growth factors.

Glycoproteins that act at very low concentrations
Act hierarchically
Usually produced by many cell types
Usually affect more than one lineage
Usually active on stem/progenitor cells and on functional end cells
Usually show synergistic or additive interactions with other growth factors
Often act on the neoplastic equivalent of a normal cell
Multiple actions: proliferation, differentiation, maturation, functional activation, prevention of apoptosis of progenitor cells

Table 1.3 Haemopoietic growth factors.

Act on stromal cells
IL-1
TNF
Act on pluripotential stem cells
SCF
Flt-L
Act on multipotential progenitor cells
IL-3
GM-CSF
IL-6
G-CSF
Thrombopoietin
Act on committed progenitor cells
G-CSF*
M-CSF
IL-5 (eosinophil-CSF)
Erythropoietin
Thrombopoietin*

Flt-L, Flt ligand; G- and GM-CSF, granulocyte and granulocyte-macrophage colony-stimulating factor; IL, interleukin; M-CSF, macrophage colony-stimulating factor; SCF, stem cell factor; TNF, tumour necrosis factor.

* These also act synergistically with early acting factors on pluripotential progenitors.

They share a number of common properties (Table 1.2) and act at different stages of haemopoiesis (Table 1.3; Fig. 1.7). Stromal cells are the major source of growth factors except for erythropoietin, 90% of which is synthesized in the kidney,

and thrombopoietin, made largely in the liver. An important feature of growth factor action is that two or more factors may synergize in stimulating a particular cell to proliferate or differentiate. Moreover, the action of one growth factor on a cell may stimulate production of another growth factor or growth factor receptor. SCF and Flt ligand (Flt-L) act locally on the pluripotential stem cells and on early myeloid and lymphoid progenitors (Fig. 1.7). Interleukin 3 (IL-3) and GM-CSF are multipotential growth factors with overlapping activities. G-CSF and thrombopoietin enhance the effects of SCF, Flt-L, IL-3 and GM-CSF on survival and differentiation of the early haemopoietic cells.

These factors maintain a pool of haemopoietic stem and progenitor cells on which later acting factors erythropoietin, G-CSF, M-CSF, IL-5 and thrombopoietin act to increase production of one or other cell lineage in response to the body's need. Granulocyte and monocyte formation, for example, can be stimulated by infection or inflammation through release of IL-1 and tumour necrosis factor (TNF) which then stimulate stromal cells to produce growth factors in an interacting network (Fig. 7.4). In contrast, cytokines such as transforming growth factor- β (TGF- β) and γ -interferon (IFN- γ) can exert a negative effect on haemopoiesis and may have a role in the development of aplastic anaemia (p. 244).

Growth factor receptors and signal transduction

The biological effects of growth factors are mediated through specific receptors on target cells. Many receptors (e.g. erythropoietin (epo) receptor (R), GM-CSF-R) are from the *haematopoietin receptor superfamily* which dimerize after binding their ligand.

Dimerization of the receptor leads to activation of a complex series of intracellular signal transduction pathways of which the three major ones are the JAK/STAT, the mitogen activated protein (MAP) kinase and the phosphatidylinositol 3 (PI3) kinase pathways (Figs 1.8, 19.2). The Janus associated kinase (JAK) proteins are a family of four tyrosine-specific protein kinases that associate with the intracellular domains of the growth factor receptors (Fig. 1.8). A growth factor molecule binds simultaneously to the extracellular domains of two or three receptor




Fig. 1.7 A diagram of the role of growth factors in normal haemopoiesis. Multiple growth factors act on the earlier marrow stem and progenitor cells. EPO, erythropoietin; PSC, pluripotent stem cell; SCF, stem cell factor; TPO, thrombopoietin. For other abbreviations see Fig. 1.2.

molecules, resulting in their aggregation. Receptor aggregation induces activation of the JAKs which now phosphorylate members of the signal transducer and activator of transcription (STAT) family of transcription factors. This results in their dimerization and translocation from the cell cytoplasm across the nuclear membrane to the cell nucleus. Within the nucleus STAT dimers activate transcription of specific genes. A model for control of gene expression by a transcription factor is shown in Fig. 1.9. The clinical importance of this pathway is revealed by the finding of an activating mutation of the *JAK2* gene as the cause of polycythaemia rubra vera (p. 230).

JAK can also activate the MAPK pathway which is regulated by Ras and controls proliferation. PI3 kinases phosphorylate inositol lipids which have a wide range of downstream effects including activation of AKT (protein kinase B) leading to block of apoptosis and other actions (Fig. 1.8, 19.2). Different domains of the intracellular receptor protein may

signal for the different processes (e.g. proliferation or suppression of apoptosis) mediated by growth factors.

A second smaller group of growth factors, including SCF, Flt-3L and macrophage colony-stimulating factor (M-CSF) (Table 1.3), bind to receptors that have an extracellular immunoglobulin-like domain linked via a transmembrane bridge to a cytoplasmic tyrosine kinase domain. Growth factor binding results in dimerization of these receptors and consequent activation of the tyrosine kinase domain. Phosphorylation of tyrosine residues in the receptor itself generates binding sites for signalling proteins which initiate complex cascades of biochemical events resulting in changes in gene expression, cell proliferation and prevention of apoptosis.

The cell cycle

The cell division cycle, generally known simply as the *cell cycle*, is a complex process that lies at




Fig. 1.8 Control of haemopoiesis by growth factors. The factors act on cells expressing the corresponding receptors. Binding of a growth factor to its receptor activates the JAK/STAT, MAPK and phosphatidylinositol3-kinase (PI3K) pathways (Fig. 19.2) which leads to transcriptional activation of specific genes. E2F is a transcription factor needed for cell transition from G1 to S phase. E2F is inhibited by the tumour suppressor gene *Rb* (retinoblastoma) which can be indirectly activated by p53. The synthesis and degradation of different cyclins (Fig. 1.10) stimulates the cell to pass through the different phases of the cell cycle. The growth factors may also suppress apoptosis by activating AKT (protein kinase B).




Fig. 1.9 Model for control of gene expression by a transcription factor. The DNA-binding domain of a transcription factor binds a specific enhancer sequence adjacent to a structural gene. The transactivation domain then binds a molecule of RNA polymerase, thus augmenting its binding to the TATA box. The RNA polymerase now initiates transcription of the structural gene to form mRNA. Translation of the mRNA by the ribosomes generates the protein encoded by the gene.

the heart of haemopoiesis. Dysregulation of cell proliferation is also the key to the development of malignant disease. The duration of the cell cycle is variable between different tissues but the basic principles remain constant. The cycle is divided into the mitotic phase (*M phase*), during which the cell physically divides, and *interphase* during which the chromosomes are duplicated and cell growth occurs prior to division (Fig. 1.10). The M phase is further partitioned into classical *mitosis* in which nuclear

division is accomplished, and *cytokinesis* in which cell fission occurs.

Interphase is divided into three main stages: a *G₁ phase* in which the cell begins to commit to replication, an *S phase* during which DNA content doubles (Fig. 1.10b) and the chromosomes replicate and the *G₂ phase* in which the cell organelles are copied and cytoplasmic volume is increased. If cells rest prior to division they enter a *G₀* state where they can remain for long periods of time. The number of cells at each stage of the cell cycle can be assessed by exposing cells to a chemical or radiolabel that gets incorporated into newly generated DNA or by flow cytometry.

The cell cycle is controlled by two *checkpoints* which act as brakes to coordinate the division process at the end of the *G₁* and *G₂* phases. Two major classes of molecules control these checkpoints, *cyclin dependent protein kinases* (Cdk) which phosphorylate downstream protein targets and *cyclins* which bind to Cdks and regulate their activity. An example of the importance of these systems is demonstrated by mantle cell lymphoma which results from the constitutive activation of cyclin D1 as a result of a chromosomal translocation (p. 212).





Fig. 1.10 (a) The stages of the cell cycle. Progression through cell cycle is regulated by specific combinations of cyclin dependent protein kinases (Cdk) and cyclin proteins. The synthesis and degradation of different cyclins stimulates the cell to pass through the different phases of the cell cycle although the exact role of each heterodimer is currently uncertain. (b) Relationship between the DNA content of a cell expressed in arbitrary units as 2c increasing to 4c and its position in the cell cycle. (Adapted from Wickramasinghe S.N. (1975) *Human Bone Marrow*, Blackwell Scientific, Oxford, p. 13.)

Apoptosis

Apoptosis is a regulated process of physiological cell death in which cells are triggered to activate intracellular proteins that lead to the death of the cell. Morphologically it is characterized by cell shrinkage, condensation of the nuclear chromatin, fragmentation of the nucleus and cleavage of DNA at internucleosomal sites. It is an important process for maintaining tissue homeostasis in haemopoiesis and lymphocyte development.

Apoptosis results from the action of intracellular cysteine proteases called *caspases* which are activated following cleavage and lead to endonuclease digestion of DNA and disintegration of the cell skeleton (Fig. 1.11). There are two major pathways by which caspases can be activated. The first is by signalling through membrane proteins such as Fas or TNF receptor via their intracellular death domain. An example of this mechanism is shown by activated cytotoxic T cells expressing Fas ligand which induce apoptosis in target cells. The second




Fig. 1.11 Representation of apoptosis. Apoptosis is initiated via two main stimuli: (i) signalling through cell membrane receptors such as FAS or tumour necrosis factor (TNF) receptor; or (ii) release of cytochrome c from mitochondria. Membrane receptors signal apoptosis through an intracellular death domain leading to activation of caspases which digest DNA. Cytochrome c binds to the cytoplasmic protein Apaf-1 leading to activation of caspases. The intracellular ratio of pro- (e.g. BAX) or anti-apoptotic (e.g. BCL-2) members of the BCL-2 family may influence mitochondrial cytochrome c release. Growth factors raise the level of BCL-2 inhibiting cytochrome c release whereas DNA damage, by activating p53, raises the level of BAX which enhances cytochrome c release.

pathway is via the release of cytochrome c from mitochondria. Cytochrome c binds to Apaf-1 which then activates caspases. DNA damage induced by irradiation or chemotherapy may act through this pathway. The protein p53 has an important role in sensing DNA damage. It activates apoptosis by raising the cell level of BAX which then increases cytochrome c release (Fig. 1.11). P53 also shuts down the cell cycle to stop the damaged cell from dividing (Fig. 1.8). The cellular level of p53 is rigidly controlled by a second protein MDM2. Following death, apoptotic cells display molecules that lead to their ingestion by macrophages.

As well as molecules that mediate apoptosis there are several intracellular proteins that protect cells from apoptosis. The best characterized example is BCL-2. BCL-2 is the prototype of a family of related proteins, some of which are anti-apoptotic and some, like BAX, pro-apoptotic. The intracellular ratio of BAX and BCL-2 determines the relative susceptibility of cells to apoptosis and may act through regulation of cytochrome c release from mitochondria.

Many of the genetic changes associated with malignant disease lead to a reduced rate of apoptosis and hence prolonged cell survival. The clearest

example is the translocation of the *BCL-2* gene to the immunoglobulin heavy chain locus in the t(14; 18) translocation in follicle centre lymphoma. Overexpression of the BCL-2 protein makes the malignant B cells less susceptible to apoptosis. Apoptosis is the normal fate for most B cells undergoing selection in the lymphoid germinal centres.

Several translocations leading to the generation of fusion proteins such as t(9; 22), t(1; 14) and t(15; 17) also result in inhibition of apoptosis (Chapter 10). In addition, genes encoding proteins that are involved in mediating apoptosis following DNA damage, such as p53 and ATM, are also frequently mutated and therefore inactivated in haemopoietic malignancies.

Transcription factors

Transcription factors regulate gene expression by controlling the transcription of specific genes or gene families. Typically, they contain at least two domains: a *DNA-binding domain* such as a leucine zipper or helix-loop-helix motif which binds to a specific DNA sequence, and an *activation domain* which contributes to assembly of the transcription complex at a gene promoter.

Adhesion molecules

A large family of glycoprotein molecules termed adhesion molecules mediate the attachment of marrow precursors, leucocytes and platelets to various components of the extracellular matrix, to endothelium, to other surfaces and to each other. The adhesion molecules on the surface of leucocytes are termed receptors and these interact with molecules (termed ligands) on the surface of potential target cells. Three main families exist:

1 Immunoglobulin superfamily This includes receptors that react with antigens (the T-cell receptors and the immunoglobulins) and antigen-independent surface adhesion molecules.

2 Selectins These are mainly involved in leucocyte and platelet adhesion to endothelium during inflammation and coagulation.

3 Integrins These are involved in cell adhesion to extracellular matrix (e.g. to collagen in wound healing and in leucocyte and platelet adhesion).

The adhesion molecules are thus important in the development and maintenance of inflammatory and immune responses, and in platelet–vessel wall and leucocyte–vessel wall interactions. Expression of adhesion molecules can be modified by extracellular and intracellular factors and this alteration of expression may be quantitative or functional. IL-1, TNF, IFN- γ , T-cell activation, adhesion to extracellular proteins and viral infection may all up-regulate expression of these molecules.

The pattern of expression of adhesion molecules on tumour cells may determine their mode of spread and tissue localization (e.g. the pattern of metastasis of carcinoma cells or non-Hodgkin's lymphoma cells into a follicular or diffuse pattern). The adhesion molecules may also determine

whether or not cells circulate in the bloodstream or remain fixed in tissues. They may also partly determine whether or not tumour cells are susceptible to the body's immune defences.

Bibliography

- Adolfsson J., Mansson R., Buza-Vidas N. *et al.* (2005) Identification of Flt3+ lympho-myeloid stem cells lacking erythro-megakaryocytic potential: a revised road map for adult blood lineage commitment. *Cell* **121**,295–306.
- Cantor A.B. and Orkin S.H. (2002) Transcriptional regulation of erythropoiesis: an affair involving multiple partners. *Oncogene* **21**,3368–76.
- Hochedlinger K. and Jaenisch R. (2003) Nuclear transplantation, embryonic stem cells and potential for cell therapy. *N Engl J Med* **349**,275–86.
- Kaushansky K. (2006) Lineage-specific hematopoietic growth factors. *N Engl J Med* **354**,2034–45.
- Lanza R., Gearhart J., Hogan B. *et al.* (eds) (2006) *Essentials of Stem Cell Biology*. Elsevier Academic Press, San Diego, California.
- Lapidot T., Dar A., Kollet O. (2005) How do stem cells find their way home? *Blood* **106**,1901–10.
- Lensch M.W. and Daley G.Q. (2006) Scientific and clinical opportunities for modeling blood disorders with embryonic stem cells. *Blood* **107**,2605–12.
- Lemoli R.M., Bertolini F., Cancedda R. *et al.* (2005) Stem cell plasticity: time for a reappraisal. *Haematologica* **90**,360–81.
- Martin-Rendon E. and Watt S.M. (2003) Skin cell plasticity. *Br J Haematol* **122**,877–91.
- Orkin S.H. and Zon L.I. (2002) Hematopoiesis and stem cells: plasticity versus developmental heterogeneity. *Nat Immunol* **4**,323–8.
- Taichman R.S. (2005) Blood and bone: two tissues whose fates are intertwined to create the hematopoietic stem-cell niche. *Blood* **105**,2631–9.
- Vats A., Bielby R.C., Tolley N.S., Nerem R., Polak J.M. (2005) Stem cells. *Lancet* **366**,592–602.

CHAPTER 2

2

Erythropoiesis and general aspects of anaemia

Erythropoietin, 12
Haemoglobin, 15
The red cell, 18

Anaemia, 20
Red cell lifespan, 27
Bibliography, 27

We each make approximately 10^{12} new erythrocytes (red cells) each day by the complex and finely regulated process of erythropoiesis. Erythropoiesis passes from the stem cell through the progenitor cells colony-forming unit granulocyte, erythroid, monocyte and megakaryocyte (CFU_{GEMM}), burst-forming unit erythroid (BFU_E) and erythroid CFU (CFU_E) (see Figs 1.2 and 2.2) to the first recognizable erythrocyte precursor in the bone marrow, the pronormoblast. This is a large cell with dark blue cytoplasm, a central nucleus with nucleoli and slightly clumped chromatin (Fig. 2.1). The pronormoblast gives rise to a series of progressively smaller normoblasts by a number of cell divisions. They also contain progressively more haemoglobin (which stains pink) in the cytoplasm; the cytoplasm stains paler blue as it loses its RNA and protein synthetic apparatus while nuclear chromatin becomes more condensed (Figs 2.1 and 2.2). The nucleus is finally extruded from the late normoblast within the marrow and a reticulocyte stage results which still contains some ribosomal RNA and is still able to synthesize haemoglobin (Fig. 2.3). This cell is slightly larger than a mature red cell, spends 1–2 days in the marrow and also circulates in the peripheral blood for 1–2 days before maturing, mainly in the spleen, when RNA is completely lost. A completely pink-staining mature erythrocyte results which is a non-nucleated biconcave disc. A single pronormoblast usually gives rise to 16 mature red cells (Fig. 2.2). Nucleated red cells (normoblasts)

are not present in normal human peripheral blood. They appear in the blood if erythropoiesis is occurring outside the marrow (extramedullary erythropoiesis) and also with some marrow diseases. Normoblasts.

Erythropoietin

Erythropoiesis is regulated by the hormone erythropoietin. The erythropoietin gene contains a hypoxia-response element at its 3' end. Erythropoietin is a heavily glycosylated polypeptide of 165 amino acids with a molecular weight of 34 kDa. Normally, 90% of the hormone is produced in the peritubular interstitial cells of the kidney and 10% in the liver and elsewhere. There are no preformed stores and the stimulus to erythropoietin production is the oxygen (O₂) tension in the tissues of the kidney (Fig. 2.4). Erythropoietin production therefore increases in anaemia, when haemoglobin for some metabolic or structural reason is unable to give up O₂ normally, when atmospheric O₂ is low or when defective cardiac or pulmonary function or damage to the renal circulation affects O₂ delivery to the kidney.

Erythropoietin stimulates erythropoiesis by increasing the number of progenitor cells committed to erythropoiesis. The transcription factors GATA-1 and FOG-1 are activated by erythropoietin receptor stimulation and are important in enhancing expression of erythroid-specific genes (e.g. haem




Fig. 2.1 Erythroblasts (normoblasts) at varying stages of development. The earlier cells are larger, with more basophilic cytoplasm and a more open nuclear chromatin pattern. The cytoplasm of the later cells is more eosinophilic as a result of haemoglobin formation.

biosynthetic and red cell membrane proteins) and also enhancing expression of anti-apoptotic genes and of the transferrin receptor (CD71). Late BFU_E and CFU_E which have erythropoietin receptors are stimulated to proliferate, differentiate and produce haemoglobin. The proportion of erythroid cells in the marrow increases and, in the chronic state, there is anatomical expansion of erythropoiesis into fatty marrow and sometimes into extramedullary sites. In infants, the marrow cavity may expand into cortical bone resulting in bone deformities with frontal bossing and protrusion of the maxilla (p. 78).

Conversely, increased O₂ supply to the tissues (because of an increased red cell mass or because haemoglobin is able to release its O₂ more readily than normal) reduces the erythropoietin drive. Tissue hypoxia also stimulates new blood vessel formation by vascular endothelial growth factor (VEGF).

Plasma erythropoietin levels can be valuable in clinical diagnosis. They are high if a tumour-secreting erythropoietin is causing polycythaemia but low in severe renal disease or polycythaemia rubra vera (Fig. 2.5).




Fig. 2.2 The amplification and maturation sequence in the development of mature red cells from the pronormoblast.

	Normoblast	Reticulocyte	Mature RBC
Nuclear DNA	Yes	No	No
RNA in cytoplasm	Yes	Yes	No
In marrow	Yes	Yes	Yes
In blood	No	Yes	Yes

Fig. 2.3 Comparison of the DNA and RNA content, and marrow and peripheral blood distribution, of the erythroblast (normoblast), reticulocyte and mature red blood cell (RBC).

Indications for erythropoietin therapy

Recombinant erythropoietin is of great value in treating anaemia resulting from renal disease or from various other causes. It is given intravenously

or subcutaneously either 3–7 times weekly or once every 1–2 weeks depending on the indication and on the preparation used (erythropoietin alpha or beta or darbepoetin alpha, a heavily glycosylated longer acting form). The main indication is end-stage renal disease (with or without dialysis). Other uses include pre-autologous blood transfusions, the anaemia of chronic disorders (e.g. in rheumatoid arthritis or cancer) and some cases of myelodysplasia or myeloma. In these conditions, often higher doses are used (Table 2.1) and the quality of life is improved (Chapter 11). A low serum erythropoietin

Table 2.1 Clinical uses of erythropoietin.

Anaemia of chronic renal disease
Myelodysplastic syndrome
Anaemia associated with malignancy and chemotherapy
Anaemia of chronic diseases
AIDS
Anaemia of prematurity
Perioperative uses

AIDS, acquired immunodeficiency syndrome.





Fig. 2.4 The production of erythropoietin by the kidney in response to its oxygen (O_2) supply. Erythropoietin stimulates erythropoiesis and so increases O_2 delivery. BFU_E , erythroid burst-forming unit; CFU_E , erythroid colony-forming unit.



level prior to treatment is valuable in predicting an effective response. Oral or parenteral iron is often needed to maximize the response to erythropoietin therapy. Side-effects include a rise in blood pressure and platelet count and local injection site reactions.

The marrow requires many other precursors for effective erythropoiesis. These include metals such as iron or cobalt, vitamins (especially vitamin B_{12} , folate, vitamin C, vitamin E, vitamin B_6 , thiamine and riboflavin) and hormones such as androgens and thyroxine. Deficiency in any of these may be associated with anaemia.

Haemoglobin

Haemoglobin synthesis

The main function of red cells is to carry O_2 to the tissues and to return carbon dioxide (CO_2) from the

Fig. 2.5 The relation between radioimmunoassay estimates of erythropoietin (EPO) in plasma and haemoglobin concentration. Anaemias exclude conditions shown to be associated with impaired production of EPO. (From M. Pippard *et al.* 1992 with permission)

Table 2.2 Normal haemoglobins in adult blood.

	Hb A	Hb F	Hb A ₂
Structure	$\alpha_2\beta_2$	$\alpha_2\gamma_2$	$\alpha_2\delta_2$
Normal (%)	96–98	0.5–0.8	1.5–3.2

tissues to the lungs. In order to achieve this gaseous exchange they contain the specialized protein haemoglobin. Each red cell contains approximately 640 million haemoglobin molecules. Each molecule of normal adult haemoglobin (Hb).A (the dominant haemoglobin in blood after the age of 3–6 months) consists of four polypeptide chains, $\alpha_2\beta_2$, each with its own haem group. The molecular weight of Hb A is 68 000. Normal adult blood also contains small quantities of two other haemoglobins: Hb F and Hb A₂. These also contain α chains, but with γ and δ chains, respectively, instead of β (Table 2.2). The synthesis of the various globin chains in the fetus and adult is discussed in more detail in Chapter 6.

Table 2.3 Normal adult red cell values.

	Male	Female
Haemoglobin (g/dL)	13.5–17.5	11.5–15.5
Haematocrit (PCV) (%)	40–52	36–48
Red cell count ($\times 10^{12}/L$)	4.5–6.5	3.9–5.6
Mean cell haemoglobin (MCH) (pg)	27–34	
Mean cell volume (MCV) (fL)	80–95	
Mean cell haemoglobin concentration (g/dL)	30–35	
Reticulocyte count ($\times 10^9/L$)	25–125	

PCV, packed cell volume.

The major switch from fetal to adult haemoglobin occurs 3–6 months after birth (Table 2.2; see Fig. 6.1b).

Haem synthesis occurs largely in the mitochondria by a series of biochemical reactions commencing with the condensation of glycine and succinyl coenzyme A under the action of the key rate-limiting enzyme δ -aminolaevulinic acid (ALA) synthase (Fig. 2.6). Pyridoxal phosphate (vitamin B₆)




Fig. 2.6 Haemoglobin synthesis in the developing red cell. The mitochondria are the main sites of protoporphyrin synthesis, iron (Fe) is supplied from circulating transferrin; globin chains are synthesized on ribosomes. δ -ALA, δ -aminolaevulinic acid; CoA, coenzyme A.




Fig. 2.7 The structure of haem.




Fig. 2.8 The oxygenated and deoxygenated haemoglobin molecule. α , β , globin chains of normal adult haemoglobin (Hb A). 2,3-DPG, 2,3-diphosphoglycerate.

is a coenzyme for this reaction which is stimulated by erythropoietin. Ultimately, protoporphyrin combines with iron in the ferrous (Fe^{2+}) state to form haem (Fig. 2.7), each molecule of which combines with a globin chain made on the polyribosomes (Fig. 2.6). A tetramer of four globin chains each with its own haem group in a 'pocket' is then formed to make up a haemoglobin molecule (Fig. 2.8).

Haemoglobin function

The red cells in systemic arterial blood carry O_2 from the lungs to the tissues and return in venous blood with CO_2 to the lungs. As the haemoglobin




Fig. 2.9 The haemoglobin oxygen (O_2) dissociation curve. 2,3-DPG, 2,3-diphosphoglycerate.

molecule loads and unloads O_2 the individual globin chains in the haemoglobin molecule move on each other (Fig. 2.8). The $\alpha_1\beta_1$ and $\alpha_2\beta_2$ contacts stabilize the molecule. The β chains slide on the $\alpha_1\beta_2$ and $\alpha_2\beta_1$ contacts during oxygenation and deoxygenation. When O_2 is unloaded the β chains are pulled apart, permitting entry of the metabolite 2,3-diphosphoglycerate (2,3-DPG) resulting in a lower affinity of the molecule for O_2 . This movement is responsible for the sigmoid form of the haemoglobin O_2 dissociation curve (Fig. 2.9). The P_{50} (i.e. the partial pressure of O_2 at which haemoglobin is half saturated with O_2) of normal blood is 26.6 mmHg. With increased affinity for O_2 , the curve shifts to the left (i.e. the P_{50} falls) while with decreased affinity for O_2 the curve shifts to the right (i.e. the P_{50} rises).

Normally *in vivo*, O_2 exchange operates between 95% saturation (arterial blood) with a mean arterial O_2 tension of 95 mmHg and 70% saturation (venous blood) with a mean venous O_2 tension of 40 mmHg.

The normal position of the curve depends on the concentration of 2,3-DPG, H^+ ions and CO_2 in the red cell and on the structure of the haemoglobin molecule. High concentrations of 2,3-DPG, H^+ or CO_2 , and the presence of certain haemoglobins (e.g. sickle haemoglobin, Hb S), shift the curve to

the right (oxygen is given up more easily) whereas fetal haemoglobin (Hb F)—which is unable to bind 2,3-DPG—and certain rare abnormal haemoglobins associated with polycythaemia shift the curve to the left because they give up O_2 less readily than normal.

Methaemoglobinaemia

This is a clinical state in which circulating haemoglobin is present with iron in the oxidized (Fe^{3+}) instead of the usual Fe^{2+} state. It may arise because of a hereditary deficiency of reduced nicotinamide adenine dinucleotide (NADH), diaphorase or inheritance of a structurally abnormal haemoglobin (Hb M). These contain an amino acid substitution affecting the haem pocket of the globin chain. Toxic methaemoglobinaemia (and/or sulphaemoglobinaemia) occurs when a drug or other toxic substance oxidizes haemoglobin. In all these states, the patient is likely to show cyanosis.

The red cell

In order to carry haemoglobin into close contact with the tissues and for successful gaseous exchange, the red cell, 8 μm in diameter, must be able: to pass repeatedly through the microcirculation whose minimum diameter is 3.5 μm , to maintain haemoglobin in a reduced (ferrous) state and to maintain osmotic equilibrium despite the high concentration of protein (haemoglobin) in the cell. Its total journey throughout its 120-day lifespan has been estimated to be 480 km (300 miles). To fulfil these functions, the cell is a flexible biconcave disc with an ability to generate energy as adenosine triphosphate (ATP) by the anaerobic glycolytic (Embden–Meyerhof) pathway (Fig. 2.10) and to generate reducing power as NADH by this pathway and as reduced nicotinamide adenine dinucleotide phosphate (NADPH) by the hexose monophosphate shunt (Fig. 2.11).

Red cell metabolism

Embden–Meyerhof pathway

In this series of biochemical reactions, glucose that enters the red cell from plasma by facilitated transfer is metabolized to lactate (Fig. 2.10). For each

molecule of glucose used, two molecules of ATP and thus two high-energy phosphate bonds are generated. This ATP provides energy for maintenance of red cell volume, shape and flexibility. The red cell has an osmotic pressure five times that of plasma and an inherent weakness of the membrane results in continual Na^+ and K^+ movement. A membrane ATPase sodium pump is needed, and this uses one molecule of ATP to move three sodium ions out and two potassium ions into the cell.

The Embden–Meyerhof pathway also generates NADH which is needed by the enzyme methaemoglobin reductase to reduce functionally dead methaemoglobin (oxidized haemoglobin) containing ferric iron (produced by oxidation of approximately 3% of haemoglobin each day) to functionally active, reduced haemoglobin. The Luebering–Rapoport shunt, or side arm, of this pathway (Fig. 2.10b) generates 2,3-DPG which forms a 1 : 1 complex with haemoglobin and, as mentioned above, is important in the regulation of haemoglobin's oxygen affinity.

Hexose monophosphate (pentose phosphate) pathway

Approximately 10% of glycolysis occurs by this oxidative pathway in which glucose-6-phosphate is converted to 6-phosphogluconate and so to ribulose-5-phosphate (Fig. 2.11). NADPH is generated and is linked with glutathione which maintains sulphhydryl (SH) groups intact in the cell including those in haemoglobin and the red cell membrane. NADPH is also used by another methaemoglobin reductase to maintain haemoglobin iron in the functionally active Fe^{2+} state. In one of the most common inherited abnormalities of red cells, glucose-6-phosphate dehydrogenase (G6PD) deficiency, the red cells are extremely susceptible to oxidant stress (p. 63).

Red cell membrane

The red cell membrane comprises a lipid bilayer, integral membrane proteins and a membrane skeleton (Fig. 2.12). Approximately 50% of the membrane is protein, 20% phospholipids, 20% cholesterol molecules and up to 10% is carbohydrate. Carbohydrates occur only on the external surface while





Fig. 2.10 (a) The Embden-Meyerhof glycolytic pathway. (b) The Luebering-Rapoport shunt which regulates the concentration of 2,3-diphosphoglycerate (2,3-DPG) in the red cell. ADP, adenosine diphosphate; ATP, adenosine triphosphate; Hb, haemoglobin; NAD, NADH, nicotinamide adenine dinucleotide; PG, phosphoglycerate.



proteins are either peripheral or integral, penetrating the lipid bilayer. Several red cell proteins have been numbered according to their mobility on polyacrylamide gel electrophoresis (PAGE), e.g. Band 3, proteins 4.1, 4.2 (Fig. 2.12).

The membrane skeleton is formed by structural proteins that include α and β spectrin, ankyrin, protein 4.1 and actin. These proteins form an horizontal lattice on the internal side of the red cell membrane and are important in maintaining the biconcave




Fig. 2.11 The hexose monophosphate shunt pathway. GSH, GSSG, glutathione; NADP, NADPH, nicotinamide adenine dinucleotide phosphate; P, phosphate; PG, phosphoglycerate.

shape. Spectrin is the most abundant and consists of two chains, α and β , wound around each other to form heterodimers which then self-associate head-to-head to form tetramers. These tetramers are linked at the tail end to actin and are attached to protein band 4.1. At the head end, the β spectrin chains attach to ankyrin which connects to band 3, the transmembrane protein that acts as an anion channel ('vertical connections') (Fig. 2.12). Protein 4.2 enhances this interaction.




Fig. 2.12 The structure of the red cell membrane. Some of the penetrating and integral proteins carry carbohydrate antigens; other antigens are attached directly to the lipid layer.

Defects of the proteins may explain some of the abnormalities of shape of the red cell membrane (e.g. hereditary spherocytosis and elliptocytosis) (Chapter 5) while alterations in lipid composition because of congenital or acquired abnormalities in plasma cholesterol or phospholipid may be associated with other membrane abnormalities. For instance, an increase in cholesterol and phospholipid has been suggested as one cause of target cells whereas a large selective increase in cholesterol may cause acanthocyte formation (see Fig. 2.16).

Anaemia

This is defined as a reduction in the haemoglobin concentration of the blood. Although normal values can vary between laboratories, typical values would be less than 13.5 g/dL in adult males and less than 11.5 g/dL in adult females (Fig. 2.13). From the age of 2 years to puberty, less than 11.0 g/dL indicates anaemia. As newborn infants have a high haemoglobin level, 14.0 g/dL is taken as the lower limit at birth (Fig. 2.13). Reduction of haemoglobin is usually accompanied by a fall in red cell count and packed cell volume (PCV) but these may be normal in some patients with subnormal haemoglobin levels (and therefore by definition anaemic). Alterations in total circulating plasma volume as well as of total circulating haemoglobin mass determine the haemoglobin concentration. Reduction in plasma volume (as in dehydration) may mask anaemia or even

Table 2.4 Classification of anaemia.

Microcytic, hypochromic	Normocytic, normochromic	Macrocytic
MCV <80 fL	MCV 80–95 fL	MCV >95 fL
MCH <27 pg	MCH ≥27 pg	Megaloblastic: vitamin B ₁₂ or folate deficiency
Iron deficiency	Many haemolytic anaemias	Non-megaloblastic: alcohol, liver disease, myelodysplasia, aplastic anaemia, etc. (Table 4.11)
Thalassaemia	Anaemia of chronic disease (some cases)	
Anaemia of chronic disease (some cases)	After acute blood loss	
Lead poisoning	Renal disease	
Sideroblastic anaemia (some cases)	Mixed deficiencies	
	Bone marrow failure (e.g. post-chemotherapy, infiltration by carcinoma, etc.)	

MCH, mean corpuscular haemoglobin; MCV, mean corpuscular volume.




Fig. 2.13 The lower limit of normal blood haemoglobin concentration in men, women and children of various ages.

cause (pseudo) polycythaemia (p. 236); conversely, an increase in plasma volume (as with splenomegaly or pregnancy) may cause anaemia even with a normal total circulating red cell and haemoglobin mass.

After acute major blood loss, anaemia is not immediately apparent because the total blood volume is reduced. It takes up to a day for the plasma volume to be replaced and so for the degree of anaemia to become apparent (p. 350). Regeneration of the haemoglobin mass takes substantially longer. The initial clinical features of major blood loss are therefore a result of reduction in blood volume rather than anaemia.

Clinical features of anaemia

The major adaptations to anaemia are in the cardio-

vascular system (with increased stroke volume and tachycardia) and in the haemoglobin O₂ dissociation curve. In some patients with quite severe anaemia there may be no symptoms or signs, whereas others with mild anaemia may be severely incapacitated. The presence or absence of clinical features can be considered under four major headings.

1 Speed of onset Rapidly progressive anaemia causes more symptoms than anaemia of slow onset because there is less time for adaptation in the cardiovascular system and in the O₂ dissociation curve of haemoglobin.

2 Severity Mild anaemia often produces no symptoms or signs but these are usually present when the haemoglobin is less than 9–10 g/dL. Even severe anaemia (haemoglobin concentration as low as 6.0 g/dL) may produce remarkably few symptoms,



(a)



(b)

Fig. 2.14 Pallor of the conjunctival mucosa (a) and of the nail bed (b) in two patients with severe anaemia (haemoglobin 6.0 g/dL).

however, when there is very gradual onset in a young subject who is otherwise healthy.

3 Age The elderly tolerate anaemia less well than the young because of the effect of lack of oxygen on organs when normal cardiovascular compensation (increased cardiac output caused by increased stroke volume and tachycardia) is impaired.

4 Haemoglobin O_2 dissociation curve Anaemia, in general, is associated with a rise in 2,3-DPG in the red cells and a shift in the O_2 dissociation curve to the right so that oxygen is given up more readily to tissues. This adaptation is particularly marked in some anaemias which either affect red cell metabolism directly (e.g. the anaemia of pyruvate kinase deficiency which causes a rise in 2,3-DPG concentration in the red cells) or which are associated with a low affinity haemoglobin (e.g. Hb S) (Fig. 2.9).

Symptoms

If the patient does have symptoms these are usually shortness of breath particularly on exercise, weakness, lethargy, palpitation and headaches. In older subjects, symptoms of cardiac failure, angina pectoris or intermittent claudication or confusion may be present. Visual disturbances because of retinal haemorrhages may complicate very severe anaemia, particularly of rapid onset.

Signs

These may be divided into *general* and *specific*. General signs include pallor of mucous membranes

which occurs if the haemoglobin level is less than 9–10 g/dL (Fig. 2.14). Conversely, skin colour is not a reliable sign. A hyperdynamic circulation may be present with tachycardia, a bounding pulse, cardiomegaly and a systolic flow murmur especially at the apex. Particularly in the elderly, features of congestive heart failure may be present. Retinal haemorrhages are unusual (Fig. 2.15).

Specific signs are associated with particular types of anaemia (e.g. koilonychia 'spoon nails' with iron deficiency, jaundice with haemolytic or megaloblastic anaemias, leg ulcers with sickle cell and other haemolytic anaemias, bone deformities with




Fig. 2.15 Retinal haemorrhages in a patient with severe anaemia (haemoglobin 2.5 g/dL) caused by severe chronic haemorrhage.

thalassaemia major and other severe congenital haemolytic anaemias).

The association of features of anaemia with excess infections or spontaneous bruising suggest that neutropenia or thrombocytopenia may be present, possibly as a result of bone marrow failure.

Classification and laboratory findings in anaemia

Red cell indices

The most useful classification is that based on red cell indices (Table 2.4) and divides the anaemia into microcytic, normocytic and macrocytic. As well as suggesting the nature of the primary defect, this approach may also indicate an underlying abnormality before overt anaemia has developed.

In two common physiological situations the mean corpuscular volume (MCV) may be outside the normal adult range. In the newborn for a few weeks the MCV is high but in infancy it is low (e.g. 70 fL at 1 year of age) and rises slowly throughout childhood to the normal adult range. In normal pregnancy there is a slight rise in MCV, even in the absence of other causes of macrocytosis (e.g. folate deficiency).

Other laboratory findings

Although the red cell indices will indicate the type of anaemia, further useful information can be obtained from the initial blood sample.

Leucocyte and platelet counts

Measurement of these helps to distinguish 'pure' anaemia from 'pancytopenia' (a drop in red cells, granulocytes and platelets) which suggests a more general marrow defect (e.g. caused by marrow hypoplasia or infiltration) or general destruction of cells (e.g. hypersplenism). In anaemias caused by

haemolysis or haemorrhage, the neutrophil and platelet counts are often raised; in infections and leukaemias the leucocyte count is also often raised and there may be abnormal leucocytes or neutrophil precursors present.

Reticulocyte count

The normal percentage is 0.5–2.5%, and the absolute count $25\text{--}125 \times 10^9/\text{L}$. This should rise in anaemia because of erythropoietin increase and be higher the more severe the anaemia. This is particularly so when there has been time for erythroid hyperplasia to develop in the marrow as in chronic haemolysis. After an acute major haemorrhage there is an erythropoietin response in 6 h, the reticulocyte count rises within 2–3 days, reaches a maximum in 6–10 days and remains raised until the haemoglobin returns to the normal level. If the reticulocyte count is not raised in an anaemic patient this suggests impaired marrow function or lack of erythropoietin stimulus (Table 2.5).

Blood film

It is essential to examine the blood film in all cases of anaemia. Abnormal red cell morphology (Fig. 2.16) or red cell inclusions (Fig. 2.17) may suggest a particular diagnosis. When causes of both microcytosis and macrocytosis are present (e.g. mixed iron and folate or B_{12} deficiency) the indices may be normal but the blood film reveals a 'dimorphic' appearance (a dual population of large well-haemoglobinized cells and small hypochromic cells). During the blood film examination the white cell differential count is performed, platelet number and morphology are assessed and the presence or absence of abnormal cells (e.g. normoblasts, granulocyte precursors or blast cells) is noted.

Table 2.5 Factors impairing the normal reticulocyte response to anaemia.

Marrow diseases (e.g. hypoplasia, infiltration by carcinoma, lymphoma, myeloma, acute leukaemia, tuberculosis)
Deficiency of iron, vitamin B_{12} or folate
Lack of erythropoietin (e.g. renal disease)
Reduced tissue O_2 consumption (e.g. myxoedema, protein deficiency)
Ineffective erythropoiesis (e.g. thalassaemia major, megaloblastic anaemia, myelodysplasia, myelofibrosis, congenital dyserythropoietic anaemia)
Chronic inflammatory or malignant disease















Red cell abnormality	Causes	Red cell abnormality	Causes
 Normal		 Microspherocyte	Hereditary spherocytosis, autoimmune haemolytic anaemia, septicaemia
 Macrocyte	Liver disease, alcoholism. Oval in megaloblastic anaemia	 Fragments	DIC, microangiopathy, HUS, TTP, burns, cardiac valves
 Target cell	Iron deficiency, liver disease, haemoglobinopathies, post-splenectomy	 Elliptocyte	Hereditary elliptocytosis
 Stomatocyte	Liver disease, alcoholism	 Tear drop poikilocyte	Myelofibrosis, extramedullary haemopoiesis
 Pencil cell	Iron deficiency	 Basket cell	Oxidant damage—e.g. G6PD deficiency, unstable haemoglobin
 Echinocyte	Liver disease, post-splenectomy	 Sickle cell	Sickle cell anaemia
 Acanthocyte	Liver disease, abetalipoproteinaemia, renal failure	 Microcyte	Iron deficiency, haemoglobinopathy

Fig. 2.16 Some of the more frequent variations in size (anisocytosis) and shape (poikilocytosis) that may be found in different anaemias. DIC, disseminated intravascular coagulopathy; G6PD, glucose-6-phosphate dehydrogenase; HUS, haemolytic uraemic syndrome; TTP, thrombotic thrombocytopenic purpura.

Bone marrow examination

This may be performed by aspiration or trephine biopsy (Fig. 2.18). During bone marrow aspiration a needle is inserted into the marrow and a liquid sample of marrow is sucked into a syringe. This is then spread on a slide for microscopy and stained by the usual Romanowsky technique. A great deal of morphological information can be obtained by examining aspirate slides. The detail of the developing cells can be examined (e.g. normoblastic or megaloblastic), the proportion of the different cell lines assessed (myeloid : erythroid ratio) and the presence of cells foreign to the marrow (e.g. secondary carcinoma) observed. The cellularity of the marrow can also be viewed provided fragments are obtained.

An iron stain is performed routinely so that the amount of iron in reticuloendothelial stores (macrophages) and as fine granules ('siderotic' granules) in the developing erythroblasts can be assessed.

An aspirate sample may also be used for a number of other specialized investigations (Table 2.6).

A trephine biopsy provides a solid core of bone including marrow and is examined as a histological specimen after fixation in formalin, decalcification and sectioning. It is less valuable than aspiration when individual cell detail is to be examined but provides a panoramic view of the marrow from which overall marrow architecture, cellularity and presence of fibrosis or abnormal infiltrates can be reliably determined.

Fig. 2.17 Red blood cell (RBC) inclusions which may be seen in the peripheral blood film in various conditions. The reticulocyte RNA and Heinz bodies are only demonstrated by supravital staining (e.g. with new methylene blue). Heinz bodies are oxidized denatured haemoglobin. Siderotic granules (Pappenheimer bodies) contain iron. They are purple on conventional staining but blue with Perls' stain. The Howell-Jolly body is a DNA remnant. Basophilic stippling is denatured RNA.





Fig. 2.18 (a) The bone marrow aspiration needle and a smear made from a bone marrow aspirate. (b) The bone marrow trephine (biopsy) needle and normal trephine section.

Table 2.6 Comparison of bone marrow aspiration and trephine biopsy.

	Aspiration	Trephine
Site	Posterior iliac crest or sternum (tibia in infants)	Posterior iliac crest
Stains	Romanowsky; Perls' reaction (for iron)	Haematoxylin and eosin; reticulin (silver stain)
Result available	1–2 h	1–7 days (according to decalcification method)
Main indications	Investigation of anaemia, pancytopenia, suspected leukaemia or myeloma, neutropenia, thrombocytopenia, etc.	Indications for additional trephine: suspicion of polycythaemia vera, myelofibrosis and other myeloproliferative disorders, aplastic anaemia, malignant lymphoma, secondary carcinoma, cases of splenomegaly or pyrexia of undetermined cause. Any case where aspiration gives a 'dry' tap
Special tests	Cytogenetics, microbiological culture, biochemical analysis, immunological and cytochemical markers, DNA or RNA analysis for gene abnormalities, progenitor cell culture	Immunological staining

Ineffective erythropoiesis

Erythropoiesis is not entirely efficient because approximately 10–15% of developing erythroblasts die within the marrow without producing mature cells. This is termed ineffective erythropoiesis and it is substantially increased in a number of chronic anaemias. The serum unconjugated bilirubin

(derived from breaking down haemoglobin) and lactate dehydrogenase (LDH, derived from breaking down cells) are usually raised when ineffective erythropoiesis is marked. The reticulocyte count is low in relation to the degree of anaemia and to the proportion of erythroblasts in the marrow.




Fig. 2.19 The relative proportions of marrow erythroblastic activity, circulating red cell mass and red cell lifespan in normal subjects and in three types of anaemia.

Assessment of erythropoiesis

Total erythropoiesis and the amount of erythropoiesis that is effective in producing circulating red cells can be assessed by examining the bone marrow, haemoglobin level and reticulocyte count.

Total erythropoiesis is assessed from the marrow cellularity and the myeloid : erythroid ratio (i.e. the proportion of granulocyte precursors to red cell precursors in the bone marrow, normally 2.5 : 1 to 12 : 1). This ratio falls and may be reversed when total erythropoiesis is selectively increased.

Effective erythropoiesis is assessed by the reticulocyte count. This is raised in proportion to the degree of anaemia when erythropoiesis is effective, but is low when there is ineffective erythropoiesis or an abnormality preventing normal marrow response (Table 2.5).

Red cell lifespan

This used to be measured by ^{51}Cr -labelled red cell survival. A sample of the subject's blood was

incubated with ^{51}Cr which binds firmly to haemoglobin and the labelled cells reinjected into the circulation. The disappearance of ^{51}Cr from the blood was measured sequentially over the next 3 weeks. The sites of red cell destruction were determined by surface counting over the spleen, liver and heart (as an index of blood activity). This test has now fallen out of routine use: Figure 2.19 shows typical changes in marrow erythropoiesis, circulating red cell mass and red cell life span in some of the different types of anaemia.

Bibliography

- Bennett C.L., Luminari S., Nissenson A.R. *et al.* (2004) Pure red-cell aplasia and epoetin therapy. *N Engl J Med* 351,1403–8.
- Cantor A.B. and Orkin S.H. (2002) Transcriptional regulation of erythropoiesis: an affair involving multiple partners. *Oncogene* 21,3368–76.
- Eschbach J.W. (2005) Iron requirements in erythropoietin therapy. *Best Pract Res Clin Haematol* 18,347–61.
- Samol J. and Littlewood T.J. (2003) The efficacy of rHuEPO in cancer-related anaemia. *Br J Haematol* 121,3–11.

CHAPTER 3

3

Hypochromic anaemias and iron overload

Nutritional and metabolic aspects of iron, 28

Iron deficiency, 33

Anaemia of chronic disorders, 39

Sideroblastic anaemia, 40

Iron overload, 41

Bibliography, 43

Iron deficiency is the most common cause of anaemia in every country of the world. It is the most important cause of a microcytic hypochromic anaemia, in which the two red cell indices MCV (mean corpuscular volume) and MCH (mean corpuscular haemoglobin) are reduced and the blood film shows small (microcytic) and pale (hypochromic) red cells. This appearance is caused by a defect in haemoglobin synthesis (Fig. 3.1). The major differential diagnosis in microcytic hypochromic anaemia is thalassaemia which is considered in Chapter 6 and anaemia of chronic disease which is dealt with in this chapter.

Nutritional and metabolic aspects of iron

Iron is one of the most common elements in the Earth's crust, yet iron deficiency is the most common cause of anaemia, affecting about 500 million people worldwide. This is because the body has a limited ability to absorb iron and excess loss of iron as a result of haemorrhage is frequent.

Body iron distribution and transport

The transport and storage of iron is largely mediated by three proteins: transferrin, the transferrin receptor 1 (TfR1) and ferritin.




Fig. 3.1 The causes of a hypochromic, microcytic anaemia. These include lack of iron (iron deficiency) or of iron release from macrophages to serum (anaemia of chronic inflammation or malignancy), failure of protoporphyrin synthesis (sideroblastic anaemia) or of globin synthesis (α- or β-thalassaemia). Lead also inhibits haem and globin synthesis.




Fig. 3.2 Daily iron cycle. Most of the iron in the body is contained in circulating haemoglobin (Table 3.1) and is reutilized for haemoglobin synthesis after the red cells die. Iron is transferred from macrophages to plasma transferrin and so to bone marrow erythroblasts. Iron absorption is normally just sufficient to make up for iron loss. The dashed line indicates ineffective erythropoiesis.

Transferrin can contain up to two atoms of iron. It delivers iron to tissues that have transferrin receptors, especially erythroblasts in the bone marrow which incorporate the iron into haemoglobin (Fig. 3.2). The transferrin is then reutilized. At the end of their life, red cells are broken down in the macrophages of the reticuloendothelial system (RES) and the iron is released from haemoglobin, enters the plasma and provides most of the iron on transferrin. Only a small proportion of plasma transferrin iron comes from dietary iron, absorbed through the duodenum and jejunum.

Some iron is stored in the macrophages as ferritin and haemosiderin, the amount varying widely according to overall body iron status. Ferritin is a water-soluble protein-iron complex of molecular weight 465 000. It is made up of an outer protein shell, apoferritin, consisting of 22 subunits and an iron-phosphate-hydroxide core. It contains up to

20% of its weight as iron and is not visible by light microscopy. Each molecule of apoferritin may bind up to 4000–5000 atoms of iron.

Haemosiderin is an insoluble protein-iron complex of varying composition containing approximately 37% iron by weight. It is derived from partial lysosomal digestion of aggregates of ferritin molecules and is visible in macrophages and other cells by light microscopy after staining by Perl's (Prussian blue) reaction. Iron in ferritin and haemosiderin is in the ferric form. It is mobilized after reduction to the ferrous form, vitamin C being involved. A copper-containing enzyme, caeruloplasmin, catalyses oxidation of the iron to the ferric form for binding to plasma transferrin.

Iron is also present in muscle as myoglobin and in most cells of the body in iron-containing enzymes (e.g. cytochromes, succinic dehydrogenase, catalase) (Table 3.1). This tissue iron is less likely to

Table 3.1 The distribution of body iron.

Amount of iron in average adult	Male (g)	Female (g)	Percentage of total
Haemoglobin	2.4	1.7	65
Ferritin and haemosiderin	1.0 (0.3–1.5)	0.3 (0–1.0)	30
Myoglobin	0.15	0.12	3.5
Haem enzymes (e.g. cytochromes, catalase, peroxidases, flavoproteins)	0.02	0.015	0.5
Transferrin-bound iron	0.004	0.003	0.1

become depleted than haemosiderin, ferritin and haemoglobin in states of iron deficiency, but some reduction of haem-containing enzymes may occur.

Regulation of ferritin and transferrin receptor 1 synthesis

The levels of ferritin and TfR1 are linked to iron status so that iron overload causes a rise in tissue ferritin and a fall in TfR1, whereas in iron deficiency ferritin is low and TfR1 increased. This linkage arises through the binding of an iron regulatory protein (IRP) to iron response elements (IREs) on the

ferritin and TfR1 mRNA molecules. Iron deficiency increases the ability of IRP to bind to the IREs whereas iron overload reduces the binding. The site of IRP binding to IREs, whether upstream (5') or downstream (3') from the coding gene, determines whether the amount of mRNA and so protein produced is increased or decreased (Fig. 3.3). Upstream binding reduces translation whereas downstream binding stabilizes the mRNA, increasing protein translation.




Fig. 3.3 Regulation of transferrin receptor 1 (TfR1), DMT-1 (divalent metal transporter) and ferritin expression by iron regulatory protein (IRP) sensing of intracellular iron levels. IRPs (●) are able to bind to stem-loop structures called iron response elements (IREs) (⌋) in transferrin receptor or ferritin mRNAs. IRP binding to the IRE within the 3' untranslated region of TfR or DMT-1 mRNA leads to stabilization of the mRNA and increased protein

synthesis whereas IRP binding to the IRE within the 5' untranslated region of ferritin mRNA reduces translation. IRPs can exist in two states: at times of high iron levels the IRP binds iron and exhibits a reduced affinity for the IREs whereas when iron levels are low the binding of IRPs to IREs is increased. In this way synthesis of TfR, DMT-1 and ferritin is coordinated to physiological requirements.




Fig. 3.4 The role of hepcidin in the regulation of iron-absorption and iron-release from macrophages. BMP, bone morphogenetic protein; HJV, hemojuvelin; Smad4, transcription factor Smad4 which stimulates hepcidin synthesis; Tfr2, transferrin receptor 2.

When plasma iron is raised and transferrin is saturated the amount of iron transferred to parenchymal cells (e.g. those of the liver, endocrine organs, pancreas and heart) is increased and this is the basis of the pathological changes associated with iron loading conditions.

Hepcidin

Hepcidin is a 25-amino acid polypeptide produced by liver cells. It is both an acute phase protein and the major hormonal regulator of iron homeostasis (Fig. 3.4). It inhibits iron release from macrophages, intestinal epithelial cells and from placental syncytiotrophoblasts by its interaction with the transmembrane iron exporter ferroportin, accelerating degradation of ferroportin mRNA. Increased production of hepcidin is induced by inflammation via interleukin 6 (IL-6). Hepcidin synthesis and secretion are controlled by three proteins: HFE, hemojuvelin and transferrin receptor 2. Decreased production of hepcidin occurs in response to iron deficiency, hypoxia and ineffective erythropoiesis.

Transferrin receptor 2

This senses the degree of saturation of transferrin and is a key regulator of hepcidin synthesis (Fig. 3.4). High saturation levels of transferrin stimulate hepcidin synthesis by this pathway whereas low saturation levels as in iron deficiency reduce hepcidin synthesis. HFE and hemojuvelin are also involved in hepcidin synthesis. Transferrin receptor 2 is restricted to erythroid, duodenal crypt and liver cells.

Dietary iron

Iron is present in food as ferric hydroxides, ferric-protein and haem-protein complexes. Both the iron content and the proportion of iron absorbed differ from food to food; in general, meat—in particular liver—is a better source than vegetables, eggs or dairy foods. The average Western diet contains 10–15 mg iron daily from which only 5–10% is normally absorbed. The proportion can be increased to 20–30% in iron deficiency or pregnancy (Table 3.2)

Table 3.2 Iron absorption.

Factors favouring absorption	Factors reducing absorption
Haem iron	Inorganic iron
Ferrous form (Fe^{2+})	Ferric form (Fe^{3+})
Acids (HCl, vitamin C)	Alkalis—antacids, pancreatic secretions
Solubilizing agents (e.g. sugars, amino acids)	Precipitating agents—phytates, phosphates
Iron deficiency	Iron excess
Ineffective erythropoiesis	Decreased erythropoiesis
Pregnancy	Infection
Hereditary haemochromatosis	Tea
Increased expression of DMT-1 and ferroportin in duodenal enterocytes	Decreased expression of DMT-1 and ferroportin in duodenal enterocytes
	Increased hepcidin

but even in these situations most dietary iron remains unabsorbed.

Iron absorption

Organic dietary iron is partly absorbed as haem and partly broken down in the gut to inorganic iron. Absorption occurs through the duodenum. Haem is absorbed through a specific receptor, HCP-1, exposed on the apical membrane of the duodenal

enterocyte. Haem is then digested to release iron. Inorganic iron absorption is favoured by factors such as acid and reducing agents that keep iron in the gut lumen in the Fe^{2+} rather than the Fe^{3+} state (Table 3.2). The protein DMT-1 (divalent metal transporter) is involved in transfer of iron from the lumen of the gut across the enterocyte microvilli (Fig. 3.5). Ferroportin at the basolateral surface controls exit of iron from the cell into portal plasma. The amount of iron absorbed is partly regulated




Fig. 3.5 The regulation of iron-absorption. Dietary ferric (Fe^{3+}) iron is reduced to Fe^{2+} and its entry to the enterocyte is through the divalent cation binder DMT-1. Its export into portal plasma is controlled by ferroportin. It is oxidized before binding to transferrin in plasma. Haem is absorbed after binding to its receptor protein HCP-1.

Table 3.3 Estimated daily iron requirements. Units are mg/day.

	Urine, sweat, faeces	Menses	Pregnancy	Growth	Total
Adult male	0.5–1				0.5–1
Postmenopausal female	0.5–1				0.5–1
Menstruating female*	0.5–1	0.5–1			1–2
Pregnant female*	0.5–1		1–2		1.5–3
Children (average)	0.5			0.6	1.1
Female (age 12–15)*	0.5–1	0.5–1		0.6	1.6–2.6

* These groups are more likely to develop iron deficiency.

according to the body's needs by changing the levels of DMT-1 according to the iron status of the duodenal villous crypt enterocyte. In iron deficiency less iron is delivered to the crypt cell from transferrin which is largely unsaturated with iron. The consequent iron deficiency in the crypt cell results in increased expression of DMT-1. This occurs by the same mechanism (IRP/IRE binding) by which transferrin receptor is increased in iron deficiency (Fig. 3.3). The increased expression of DMT-1 results, when the enterocyte reaches the apical absorptive surface of the duodenal villous 24–48 h later, in increased transfer of iron from the gut lumen into the enterocyte. Hepcidin is also a major regulator by affecting ferroportin concentration. Low hepcidin levels in iron deficiency increase ferroportin levels and allow more iron to enter portal plasma. Thus, less iron is lost when the enterocyte is shed into the gut lumen from the apex of the villous.

Ferrireductase present at the apical surface converts iron from the Fe^{3+} to Fe^{2+} state and another

enzyme, hephaestin (which contains copper), converts Fe^{2+} to Fe^{3+} at the basal surface prior to binding to transferrin.

Iron requirements


The amount of iron required each day to compensate for losses from the body and for growth varies with age and sex; it is highest in pregnancy, adolescent and menstruating females (Table 3.3). Therefore these groups are particularly likely to develop iron deficiency if there is additional iron loss or prolonged reduced intake.

Iron deficiency

Clinical features

When iron deficiency is developing the reticuloendothelial stores (haemosiderin and ferritin) become completely depleted before anaemia occurs (Fig. 3.6).

Fig. 3.6 The development of iron deficiency anaemia. Reticuloendothelial (macrophage) stores are lost completely before anaemia develops. MCH, mean corpuscular haemoglobin; MCV, mean corpuscular volume.






(a)



(b)



(c)

Fig. 3.7 Iron deficiency anaemia. (a) Koilonychia: typical 'spoon' nails. (b) Angular cheilosis: fissuring and ulceration of the corner of the mouth. (c) Paterson–Kelly (Plummer–Vinson) syndrome: barium swallow X-ray showing a filling defect (arrow) caused by a postcricoid web.

As the condition develops the patient may develop the general symptoms and signs of anaemia (p. 21) and also show a painless glossitis, angular stomatitis, brittle, ridged or spoon nails (koilonychia), dysphagia as a result of pharyngeal webs (Paterson–Kelly or Plummer–Vinson syndrome) (Fig. 3.7) and unusual dietary cravings (pica). The cause of the epithelial cell changes is not clear but may be related to reduction of iron in iron-containing enzymes. In children, iron deficiency is particularly significant as it can cause irritability, poor cognitive function and a decline in psychomotor development.

Causes of iron deficiency

Chronic blood loss, especially uterine or from the gastrointestinal tract, is the dominant cause (Table 3.4). In contrast, in developed countries dietary deficiency is rarely a cause on its own. Half a

litre of whole blood contains approximately 250 mg of iron and, despite the increased absorption of food iron at an early stage of iron deficiency, negative iron balance is usual in chronic blood loss.

Increased demands during infancy, adolescence, pregnancy, lactation and in menstruating women account for the high risk of anaemia in these particular clinical groups. Newborn infants have a store of iron derived from delayed clamping of the cord and the breakdown of excess red cells. From 3 to 6 months there is a tendency for negative iron balance because of growth. From 6 months supplemented formula milk and mixed feeding, particularly with iron-fortified foods, prevents iron deficiency.

In pregnancy increased iron is needed for an increased maternal red cell mass of approximately 35%, transfer of 300 mg of iron to the fetus and because of blood loss at delivery. Although iron absorption is also increased, iron therapy is often

Table 3.4 Causes of iron deficiency.*Chronic blood loss*

Uterine

Gastrointestinal, e.g. peptic ulcer, oesophageal varices, aspirin (or other non-steroidal anti-inflammatory drugs) ingestion, partial gastrectomy, carcinoma of the stomach, caecum, colon or rectum, hookworm, angiodysplasia, colitis, piles, diverticulosis

Rarely, haematuria, haemoglobinuria, pulmonary haemosiderosis, self-inflicted blood loss

Increased demands (see also Table 3.3)

Prematurity

Growth

Pregnancy

Erythropoietin therapy

Malabsorption

Gluten-induced enteropathy, gastrectomy

Poor diet

A contributory factor in many developing countries but rarely the sole cause except in infants and children

needed if the haemoglobin (Hb) falls below 10 g/dL or the mean cell volume (MCV) is below 82 fL in the third trimester.

Menorrhagia (a loss of 80 mL or more of blood at each cycle) is difficult to assess clinically, although the loss of clots, the use of large numbers of pads

or tampons or prolonged periods all suggest excessive loss.

It has been estimated to take 8 years for a normal adult male to develop iron deficiency anaemia solely as a result of a poor diet or malabsorption resulting in no iron intake at all. In clinical practice inadequate intake or malabsorption are only rarely the sole cause of iron deficiency anaemia although in developing countries iron deficiency may occur as a result of a life-long poor diet, consisting mainly of cereals and vegetables. Gluten-induced enteropathy, partial or total gastrectomy and atrophic gastritis (often autoimmune and with *Helicobacter pylori* infection) may, however, predispose to iron deficiency.

Laboratory findings

These are summarized and contrasted with those in other hypochromic anaemias in Table 3.7.

Red cell indices and blood film

Even before anaemia occurs, the red cell indices fall and they fall progressively as the anaemia becomes more severe. The blood film shows hypochromic microcytic cells with occasional target cells and pencil-shaped poikilocytes (Fig. 3.8). The reticulocyte count is low in relation to the degree of anaemia. When iron deficiency is associated with severe folate or vitamin B₁₂ deficiency a 'dimorphic' film

Fig. 3.8 The peripheral blood film in severe iron deficiency anaemia. The cells are microcytic and hypochromic with occasional target cells.



occurs with a dual population of red cells of which one is macrocytic and the other microcytic and hypochromic; the indices may be normal. A dimorphic blood film is also seen in patients with iron deficiency anaemia who have received recent iron therapy and produced a population of new haemoglobinized normal-sized red cells (Fig. 3.9) and when the patient has been transfused. The platelet count is often moderately raised in iron deficiency, particularly when haemorrhage is continuing.

Bone marrow iron

Bone marrow examination is not essential to assess iron stores except in complicated cases. In iron deficiency anaemia there is a complete absence of iron from stores (macrophages) and from developing erythroblasts (Fig. 3.10). The erythroblasts are small and have a ragged cytoplasm.

Serum iron and total iron-binding capacity

The serum iron falls and total iron-binding capacity (TIBC) rises so that the TIBC is less than 10% saturated (Fig. 3.11). This contrasts both with the anaemia of chronic disorders (see below) when the



Fig. 3.9 Dimorphic blood film in iron deficiency anaemia responding to iron therapy. Two populations of red cells are present: one microcytic and hypochromic, the other normocytic and well haemoglobinized.

serum iron and the TIBC are both reduced and with other hypochromic anaemias where the serum iron is normal or even raised.

Serum transferrin receptor

Transferrin receptor is shed from cells into plasma.




Fig. 3.10 Bone marrow iron assessed by Perls' stain. (a) Normal iron stores indicated by blue staining in the macrophages. Inset: normal siderotic granule in erythroblast. (b) Absence of blue staining (absence of hemosiderin) in iron deficiency. Inset: absence of siderotic granules in erythroblasts.




Fig. 3.11 The serum iron, unsaturated serum iron-binding capacity (UIBC) and serum ferritin in normal subjects and in those with iron deficiency, anaemia of chronic disorders and iron overload. The total iron-binding capacity (TIBC) is made up of the serum iron and the UIBC. In some laboratories, the transferrin content of serum is

measured directly by immunodiffusion, rather than by its ability to bind iron, and is expressed in g/L. Normal serum contains 2–4 g/L transferrin (1 g/L transferrin = 20 $\mu\text{mol/L}$ binding capacity). Normal ranges for serum iron are 10–30 $\mu\text{mol/L}$; for TIBC, 40–75 $\mu\text{mol/L}$; for serum ferritin, male, 40–340 $\mu\text{g/L}$; female, 14–150 $\mu\text{g/L}$.

The level of serum transferrin receptor (sTfR) is increased in iron deficiency anaemia but not in the anaemia of chronic disease or thalassaemia trait. The level is also raised if the overall level of erythropoiesis is increased.

Serum ferritin

A small fraction of body ferritin circulates in the serum, the concentration being related to tissue, particularly reticuloendothelial, iron stores. The normal range in men is higher than in women (Fig. 3.11). In iron deficiency anaemia the serum ferritin is very low while a raised serum ferritin indicates iron overload or excess release of ferritin from damaged tissues or an acute phase response (e.g. in inflammation). The serum ferritin is normal or raised in the anaemia of chronic disorders.

Investigation of the cause of iron deficiency (Table 3.4)

In premenopausal women, menorrhagia and/or repeated pregnancies are the usual causes of the deficiency. If these are not present other causes must be sought. In some patients with menorrhagia

a clotting or platelet abnormality (e.g. von Willebrand disease) is present. In men and postmenopausal women, gastrointestinal blood loss is the main cause of iron deficiency and the exact site is sought from the clinical history, physical and rectal examination, by occult blood tests, and by appropriate use of upper and lower gastrointestinal endoscopy and/or radiology (e.g. computed tomography (CT) of the pneumocolon) or virtual colonoscopy using the 3D colon system (Figs 3.12 and 3.13). In difficult cases a camera in a capsule can be swallowed which relays pictures of the gastrointestinal tract electronically. Tests for endomysial and transglutaminase antibodies and duodenal biopsy to look for gluten-induced enteropathy can be valuable. Hookworm ova are sought in stools of subjects from areas where this infestation occurs. Rarely, a coeliac axis angiogram is needed to demonstrate angiodysplasia.

If gastrointestinal blood loss is excluded, loss of iron in the urine as haematuria or haemosiderinuria (resulting from chronic intravascular haemolysis) is considered. A normal chest X-ray excludes the rare condition of pulmonary haemosiderosis. Rarely, patients bleed themselves producing iron deficiency.




Fig. 3.12 Investigation and management of iron deficiency anaemia. GI, gastrointestinal; TIBC, total iron-binding capacity.



Fig. 3.13 Virtual colonoscopy to show carcinoma of colon causing colonic obstruction and iron deficiency.

Treatment

The underlying cause is treated as far as possible. In addition, iron is given to correct the anaemia and replenish iron stores.

Oral iron

The best preparation is ferrous sulphate which is cheap, contains 67 mg of iron in each 200 mg tablet

and is best given on an empty stomach in doses spaced by at least 6 h. If side-effects occur (e.g. nausea, abdominal pain, constipation or diarrhoea), these can be reduced by giving iron with food or by using a preparation of lower iron content (e.g. ferrous gluconate which contains less iron (37 mg) per 300 mg tablet). An elixir is available for children. Slow-release preparations should not be used.

Oral iron therapy should be given for long enough both to correct the anaemia and to replenish body iron stores, which usually means for at least 6 months. The haemoglobin should rise at the rate of approximately 2 g/dL every 3 weeks. Failure of response to oral iron has several possible causes (Table 3.5) which should all be considered before parenteral iron is used.

Iron fortification of the diet in infants in Africa reduces the incidence of iron deficiency anaemia but increases susceptibility to malaria.

Parenteral iron

Ferric hydroxide-sucrose (Venofer®) is the safest form. It is administered by slow intravenous injection or infusion, usually 200 mg iron in each infusion. Iron dextran (CosmoFer®) can be given as slow injection or infusion either in small single doses or as a total dose infusion given in one day. Iron sorbitol (Jectofer®) is given by deep intramuscular injection but not intravenously. There may be hypersensitivity or anaphylactoid reactions so

Table 3.5 Failure of response to oral iron.

Continuing haemorrhage
Failure to take tablets
Wrong diagnosis—especially thalassaemia trait, sideroblastic anaemia
Mixed deficiency—associated folate or vitamin B ₁₂ deficiency
Another cause for anaemia (e.g. malignancy, inflammation)
Malabsorption—coeliac disease, atrophic gastritis
Use of slow-release preparation

parenteral iron is only given when there are high iron requirements as in gastrointestinal bleeding, severe menorrhagia, chronic haemodialysis, with erythropoietin therapy, and when oral iron is ineffective (e.g. iron malabsorption resulting from gluten-induced enteropathy or atrophic gastritis) or impractical (e.g. active Crohn's disease). The haematological response to parenteral iron is no faster than to adequate dosage of oral iron but the stores are replenished faster.

Anaemia of chronic disorders

One of the most common anaemias occurs in

Table 3.6 Causes of the anaemia of chronic disorders.

<i>Chronic inflammatory diseases</i>
Infections (e.g. pulmonary abscess, tuberculosis, osteomyelitis, pneumonia, bacterial endocarditis)
Non-infectious (e.g. rheumatoid arthritis, systemic lupus erythematosus and other connective tissue diseases, sarcoidosis, Crohn's disease)
<i>Malignant diseases</i>
Carcinoma, lymphoma, sarcoma

patients with a variety of chronic inflammatory and malignant diseases (Table 3.6). The characteristic features are:

- 1 Normochromic, normocytic or mildly hypochromic (MCV rarely <75 fL) indices and red cell morphology.
- 2 Mild and non-progressive anaemia (haemoglobin rarely <9.0 g/dL)—the severity being related to the severity of the disease.
- 3 Both the serum iron and TIBC are reduced; sTfR levels are normal.
- 4 The serum ferritin is normal or raised.
- 5 Bone marrow storage (reticuloendothelial) iron is normal but erythroblast iron is reduced (Table 3.7).

Table 3.7 Laboratory diagnosis of a hypochromic anaemia.

	Iron deficiency	Chronic inflammation or malignancy	Thalassaemia trait (α or β)	Sideroblastic anaemia
MCV	Reduced in relation to severity of anaemia	Normal or mild reduction	Reduced; very low for degree of anaemia	Usually low in congenital type but MCV often raised in acquired type
MCH				
Serum iron	Reduced	Reduced	Normal	Raised
TIBC	Raised	Reduced	Normal	Normal
Serum transferrin receptor	Raised	Normal/low	Variable	Normal
Serum ferritin	Reduced	Normal or raised	Normal	Raised
Bone marrow iron stores	Absent	Present	Present	Present
Erythroblast iron	Absent	Absent	Present	Ring forms
Haemoglobin electrophoresis	Normal	Normal	Hb A ₂ raised in β form	Normal

MCH, mean corpuscular haemoglobin; MCV, mean corpuscular volume; TIBC, total iron-binding capacity.

The pathogenesis of this anaemia appears to be related to decreased release of iron from macrophages to plasma, reduced red cell lifespan and an inadequate erythropoietin response to anaemia caused by the effects of cytokines such as IL-1 and tumour necrosis factor (TNF) on erythropoiesis.

Hepcidin, released by the liver in response to inflammation, inhibits macrophage release of iron as well as iron absorption. The anaemia is corrected by successful treatment of the underlying disease and does not respond to iron therapy. Recombinant erythropoietin improves the anaemia in some cases. In many conditions this anaemia is complicated by anaemia resulting from other causes (e.g. iron, vitamin B₁₂ or folate deficiency, renal failure, bone marrow failure, hypersplenism, endocrine abnormality, leucoerythroblastic anaemia) and these are discussed in Chapter 26.

Sideroblastic anaemia

This is a refractory anaemia with hypochromic cells in the peripheral blood and increased marrow iron; it is defined by the presence of many pathological ring sideroblasts in the bone marrow (Fig. 3.14). These are abnormal erythroblasts containing numerous




Fig. 3.14 Ring sideroblasts with a perinuclear ring of iron granules in sideroblastic anaemia.

Table 3.8 Classification of sideroblastic anaemia.

Hereditary

Usually occurs in males, transmitted by females; also occurs rarely in females (see text)

Acquired

Primary

Myelodysplasia (refractory anaemia with ring sideroblasts) (p. 185)

Secondary

Ring sideroblast formation may also occur in the bone marrow in:

Other malignant diseases of the marrow (e.g. other types of myelodysplasia, myelofibrosis, myeloid leukaemia, myeloma)

Drugs, e.g. antituberculous (isoniazid, cycloserine), alcohol, lead

Other benign conditions (e.g. haemolytic anaemia, megaloblastic anaemia, malabsorption, rheumatoid arthritis)

iron granules arranged in a ring or collar around the nucleus instead of the few randomly distributed iron granules seen when normal erythroblasts are stained for iron. Sideroblastic anaemia is diagnosed when 15% or more of marrow erythroblasts are ring sideroblasts but they can be found at lower numbers in a variety of haematological conditions.

Sideroblastic anaemia is classified into different types (Table 3.8) and the common link is a defect in haem synthesis. In the hereditary forms the anaemia is usually characterized by a markedly hypochromic and microcytic blood picture. The most common mutations are in the δ -aminolaevulinic acid synthase (ALA-S) gene which is on the X chromosome. Pyridoxal-6-phosphate is a coenzyme for ALA-S. Other rare types include mitochondrial defects, thiamine-responsive and other autosomal defects. The much more common primary acquired form is one subtype of myelodysplasia. It is also termed 'refractory anaemia with ring sideroblasts'. This condition is discussed together with the other types of myelodysplasia in Chapter 14.

In some patients, particularly with the hereditary type, there is a response to pyridoxine therapy. Folate deficiency may occur and folic acid therapy may also be tried. Other treatments that have been

tried in myelodysplasia (e.g. erythropoietin) may be tried in the primary acquired form (Chapter 14). In many severe cases, however, repeated blood transfusions are the only method of maintaining a satisfactory haemoglobin concentration and transfusional iron overload becomes a major problem.

Lead poisoning

Lead inhibits both haem and globin synthesis at a number of points. In addition it interferes with the breakdown of RNA by inhibiting the enzyme pyrimidine 5' nucleotidase, causing accumulation of denatured RNA in red cells, the RNA giving an appearance called basophilic stippling on the ordinary (Romanowsky) stain (see Fig. 2.17). The anaemia may be hypochromic or predominantly haemolytic, and the bone marrow may show ring sideroblasts. Free erythrocyte protoporphyrin is raised.

Differential diagnosis of hypochromic anaemia

Table 3.7 lists the laboratory investigations that may be necessary. The clinical history is particularly important as the source of the haemorrhage leading to iron deficiency or the presence of a chronic disease may be revealed. The country of origin and the family history may suggest a possible diagnosis of thalassaemia or other haemoglobinopathy. Physical examination may also be helpful in determining a site of haemorrhage, features of a chronic inflammatory or malignant disease, koilonychia or, in some haemoglobinopathies, an enlarged spleen or bony deformities.

In thalassaemia trait the red cells tend to be small, often with an MCV of 70 fL or less, even when anaemia is mild or absent; the red cell count is usually over $5.5 \times 10^{12}/L$. Conversely, in iron deficiency anaemia the indices fall progressively with the degree of anaemia and when anaemia is mild the indices are often only just reduced below normal (e.g. MCV 75–80 fL). In the anaemia of chronic disorders the indices are also not markedly low, an MCV in the range 75–82 fL being usual.

It is usual to perform a serum iron and TIBC measurement, or alternatively serum ferritin estimation, to confirm a diagnosis of iron deficiency. sTfR assay is also useful in distinguishing iron defi-

ciency anaemia from anaemia of chronic disease but not widely available. Haemoglobin electrophoresis with an estimation of Hb A₂ and Hb F is carried out in all patients suspected of thalassaemia or other haemoglobinopathy because of the family history, country of origin, red cell indices and blood film. Iron deficiency or the anaemia of chronic disorders may also occur in these subjects. β -Thalassaemia trait is characterized by a raised Hb A₂ above 3.5%, but in α -thalassaemia trait there is no abnormality on simple haemoglobin studies so the diagnosis is usually made by exclusion of all other causes of hypochromic red cells and by the presence of a red cell count $>5.5 \times 10^{12}/L$. DNA studies can be used to confirm the diagnosis. In some α -thalassaemia patients, however, occasional red cells show deposits of Hb H (β_4) in reticulocyte preparations (Chapter 6).

Bone marrow examination is essential if a diagnosis of sideroblastic anaemia is suspected but is not usually needed in diagnosis of the other hypochromic anaemias.

Iron overload

There is no physiological mechanism for eliminating excess iron from the body and so iron absorption is normally carefully regulated to avoid accumulation. Iron overload can occur in disorders associated with excessive absorption or chronic blood transfusion. Excessive iron deposition in tissues can cause serious damage to organs, particularly the heart, liver and endocrine organs. The causes of iron overload are listed in Tables 3.9 and 3.10. Transfusional iron overload and iron chelation therapy are discussed in Chapter 6.

Hereditary (genetic, primary) haemochromatosis

This is a group of diseases in which there is excessive absorption of iron from the gastrointestinal tract leading to iron overload of the parenchymal cells of the liver (Fig. 3.15), of the endocrine organs and, in severe cases, of the heart.

The most common gene involved is *HFE* and most patients are homozygous for a missense mutation (845 G to A) which leads to insertion of a tyrosine residue rather than cysteine in the mature

Table 3.9 The causes of iron overload.

Increased iron absorption	Hereditary (primary) haemochromatosis Ineffective erythropoiesis, e.g. thalassaemia intermedia, sideroblastic anaemia Chronic liver disease
Increased iron intake	African siderosis (dietary and genetic)
Repeated red cell transfusions	Transfusion siderosis

Table 3.10 Genetic causes of haemochromatosis, iron overload and hyperferritinaemia.

Type	Inheritance	Clinical condition	Defect in
I	AR	Classical hereditary haemochromatosis	HFE
II	AR	Juvenile haemochromatosis	Hemojuvelin Hepcidin
III	AR	Hereditary haemochromatosis	Transferrin receptor 2
IV	AD	Marked increase in RE iron, less hepatic iron	Ferroportin 1
V	AD	Hereditary hyperferritinaemia—cataract syndrome (no iron deposition)	Low molecular weight ferritin

AD, autosomal dominant; AR, autosomal recessive; RE, reticuloendothelial.




Fig. 3.15 Liver biopsy. Iron loading of hepatic parenchymal cells (Perl's stain). (Courtesy of Professor A.P. Dhillon)

protein (C282Y). This allele has a prevalence of approximately 1 in 300 within the white North European population. The *HFE* gene is situated close to the major histocompatibility complex (MHC) locus on chromosome 6. The abnormal allele is associated with HLA-A3 and -B8. Only a small proportion of those homozygous for the mutation

present with clinical features of the disease. A second mutation resulting in a histidine to aspartic acid substitution H63D is found with the C282Y mutation in approximately 5% of patients but homozygotes for the H63D mutation do not have the disease.

Serum hepcidin levels are low in patients with mutated *HFE* implying *HFE* is involved in hepcidin synthesis or secretion (Fig. 3.4). Low serum hepcidin levels lead to high levels of ferroportin on the basolateral surface of the duodenal enterocyte and so increased iron absorption. Low levels of hepcidin also lead by the same mechanism to increased release of iron from macrophages.

The consequent iron overload damages parenchymal cells and patients present usually in adult life with hepatic disease, endocrine disturbances such as diabetes mellitus or impotence, melanin skin pigmentation and arthropathy (resulting from pyrophosphate deposition). In some severe cases there is cardiac failure or arrhythmia (Chapter 6). Diagnosis is suspected by increased serum iron, increased serum transferrin saturation and ferritin. It is confirmed by testing for the *HFE* mutation. Liver biopsy may quantify the degree of iron overload and assess liver damage. Magnetic resonance

imaging (MRI) can also be used to measure liver and cardiac iron. Treatment is with regular venesection, initially at 1–2 week intervals, each unit of blood lost removing 200–250 mg iron. It is monitored by serum iron, TIBC, serum ferritin and by tests of organ function.

Rarer forms of genetic haemochromatosis are caused by mutations in the genes for hemojuvelin, transferrin receptor 2 and hepcidin (Table 3.10). All three are associated with low levels of hepcidin in serum. They often present as severe iron overload with cardiomyopathy in children, adolescents or young adults. On the other hand, ferroportin mutations cause reticuloendothelial but not parenchymal cell iron overload. Mutations of the ferritin light chain gene cause a raised monoclonal serum ferritin with cataracts resulting from ferritin deposition in the eye but no tissue iron overload.

Bibliography

- Adams P.C., Reboussin D.M., Barton J.C. *et al.* (2005) Hemochromatosis and iron-overload screening in a racially diverse population. *N Engl J Med* 352,1769–78.
- Andrews N.C. (2005) Understanding heme transport. *N Engl J Med* 353,2508–9.
- Bomford A. (2002) Genetics of haemochromatosis. *Lancet* 360,1673–4.
- Camaschella C. (2005) Understanding iron homeostasis through genetic analysis of hemochromatosis and related disorders. *Blood* 106,3710–7.
- De Domenico I., Ward D.M., Musci G. and Kaplan J. (2006) Iron overload due to mutations in ferroportin. *Haematologica* 91,92–5.
- Fleming R.E. and Bacon B.R. (2005) Orchestration of iron homeostasis. *N Engl J Med* 352,1741–4.
- Ganz T. (2005) Hepcidin: a regulator of intestinal iron absorption and iron recycling by macrophages. *Clin Haematol* 18,171–82.
- Hentze M.W., Muckenthaler M.U., Andrews N.C. (2004) Balancing acts: molecular control of mammalian iron metabolism. *Cell* 117,285–97.
- Hershko C. (ed.) (2005) Iron diseases. *Best Pract Res Clin Haematol* 18,157–380.
- Hershko C., Hoffbrand A.V., Keret D. *et al.* (2005) Role of autoimmune gastritis, *Helicobacter pylori* and celiac disease in refractory or unexplained iron deficiency anemia. *Haematologica* 90,585–95.
- Lee P.L., Beutler E., Rao S.V. *et al.* (2004) Genetic abnormalities and juvenile hemochromatosis: mutations of the *HJV* gene encoding hemojuvelin. *Blood* 103,4669–71.
- Nemeth E. and Ganz T. (2006) Hepcidin and iron-loading anemias. *Haematologica* 91,727–32.
- Nemeth E., Rivera S., Gabayan V. *et al.* (2004) IL-6 mediates hypoferremia of inflammation by inducing the synthesis of the iron regulatory hormone hepcidin. *J Clin Invest* 113,1271–6.
- Papanikolaou G., Tzilianos M., Christakis J.I. *et al.* (2005) Hepcidin in iron overload disorders. *Blood* 105,4103–5.
- Pietrangelo A. (2004) Hereditary hemochromatosis: a new look at an old disease. *N Engl J Med* 350,2383–97.
- Shayeghi M., Latunde-Dada G.O., Oakhill J.S. *et al.* (2005) Identification of an intestinal heme transporter. *Cell* 122,789–801.
- Wang R.H., Li C., Xu X. *et al.* (2005) A role of SMAD4 in iron metabolism through the positive regulation of hepcidin expression. *Cell Metab* 2,399–409.
- Weiss G. and Goodnough L. (2005) Anemia of chronic disorders. *N Engl J Med* 352,1011–22.

CHAPTER 4

4

Megaloblastic anaemias and other macrocytic anaemias

Introduction to macrocytic anaemia, 44

Megaloblastic anaemias, 44

Vitamin B₁₂ (B₁₂, cobalamin), 44

Folate, 47

Vitamin B₁₂ deficiency, 48

Folate deficiency, 49

Clinical features of megaloblastic anaemia, 49

Diagnosis of vitamin B₁₂ or folate deficiency, 52

Other megaloblastic anaemias, 56

Systemic diseases associated with folate or vitamin B₁₂ deficiency, 56

Other macrocytic anaemias, 56

Bibliography, 57

Introduction to macrocytic anaemia

In macrocytic anaemia the red cells are abnormally large (mean corpuscular volume, MCV >95 fL). There are several causes (see Table 2.4) but they can be broadly subdivided into megaloblastic and non-megaloblastic, based on the appearance of developing erythroblasts in the bone marrow.

Megaloblastic anaemias

This is a group of anaemias in which the erythroblasts in the bone marrow show a characteristic abnormality—maturation of the nucleus being

delayed relative to that of the cytoplasm. The underlying defect accounting for the asynchronous maturation of the nucleus is defective DNA synthesis and in clinical practice this is usually caused by deficiency of vitamin B₁₂ or folate. Less commonly, abnormalities of metabolism of these vitamins or other lesions in DNA synthesis may cause an identical haematological appearance (Table 4.1). Dietary and metabolic aspects of the two vitamins are reviewed before considering the anaemia.

Vitamin B₁₂ (B₁₂, cobalamin)

This vitamin is synthesized in nature by micro-organisms; animals acquire it by eating other animal foods, by internal production from intestinal bacteria (not in humans) or by eating bacterially contaminated foods. The vitamin consists of a small group of compounds, the cobalamins, which have the same basic structure, with a cobalt atom at the centre of a corrin ring which is attached to a nucleotide portion (Fig. 4.1). The vitamin is found in foods of animal origin such as liver, meat, fish and dairy produce but does not occur in fruit, cereals or vegetables. Table 4.2 compares nutritional aspects of B₁₂ and folate.

Table 4.1 Causes of megaloblastic anaemia.

Vitamin B₁₂ deficiency

Folate deficiency

Abnormalities of vitamin B₁₂ or folate metabolism

(e.g. transcobalamin deficiency, nitrous oxide, antifolate drugs)

Other defects of DNA synthesis

congenital enzyme deficiencies (e.g. orotic aciduria)

acquired enzyme deficiencies (e.g. alcohol, therapy with hydroxyurea, cytosine arabinoside)




Fig. 4.1 The structure of methylcobalamin (methyl B₁₂), the main form of vitamin B₁₂ in human plasma. Other forms include deoxyadenosylcobalamin (ado B₁₂), the main form in human tissues; hydroxocobalamin (hydroxo B₁₂), the main form in treatment; and cyanocobalamin (cyano B₁₂), the radioactively labelled (⁵⁷Co or ⁵⁸Co) form used to study vitamin B₁₂ absorption or metabolism.

Absorption

A normal diet contains a large excess of B₁₂ compared with daily needs (Table 4.2). B₁₂ is combined with the glycoprotein intrinsic factor (IF) (molecular weight, MW 45 000) which is synthesized by the gastric parietal cells. The IF-B₁₂ complex can then bind to a specific surface receptor for IF, cubilin, which then binds to a second protein, amnionless which directs endocytosis of the cubilin IF-B₁₂ complex in the distal ileum where B₁₂ is absorbed and IF destroyed (Fig. 4.2).

Transport: the transcobalamins

Vitamin B₁₂ is absorbed into portal blood where it becomes attached to the plasma-binding protein transcobalamin (TC, previously called transcobalamin II) which delivers B₁₂ to bone marrow and other tissues. Although TC is the essential plasma protein for transferring B₁₂ into the cells of the body, the amount of B₁₂ on TC is normally very low (<50 ng/L). TC deficiency causes megaloblastic anaemia because of failure of B₁₂ to enter marrow (and other cells) from plasma but the serum B₁₂ level

Table 4.2 Vitamin B₁₂ and folate: nutritional aspects.

	Vitamin B ₁₂	Folate
Normal daily dietary intake	7–30 µg	200–250 µg
Main foods	Animal produce only	Most, especially liver, greens and yeast
Cooking	Little effect	Easily destroyed
Minimal adult daily requirement	1–2 µg	100–150 µg
Body stores	2–3 mg (sufficient for 2–4 years)	10–12 mg (sufficient for 4 months)
Absorption		
Site	Ileum	Duodenum and jejunum
Mechanism	Intrinsic factor	Conversion to methyltetrahydrofolate
Limit	2–3 µg/day	50–80% of dietary content
Enterohepatic circulation	5–10 µg/day	90 µg/day
Transport in plasma	Most bound to haptocorrin; TC essential for cell uptake	Weakly bound to albumin
Major intracellular physiological forms	Methyl- and deoxyadenosylcobalamin	Reduced polyglutamate derivatives
Usual therapeutic form	Hydroxocobalamin	Folic (pteroylglutamic) acid

TC, transcobalamin.




Fig. 4.2 The absorption of dietary vitamin B₁₂ after combination with intrinsic factor (IF), through the ileum. Folate absorption occurs through the duodenum and jejunum after conversion of all dietary forms to methyltetrahydrofolate (methyl THF). TC, transcobalamin.

in TC deficiency is normal. This is because most B₁₂ in plasma is bound to another transport protein, haptocorrin (previously called transcobalamin I). This is a glycoprotein largely synthesized by granulocytes and macrophages. In myeloproliferative diseases where granulocyte production is greatly increased, the haptocorrin and B₁₂ levels in serum both rise considerably. B₁₂ bound to haptocorrin does not transfer to marrow; it appears to be functionally 'dead'. Closely related glycoproteins to plasma haptocorrin are present in gastric juice, milk and other body fluids.

Biochemical function

Vitamin B₁₂ is a coenzyme for two biochemical reactions in the body: first, as methyl B₁₂ it is a cofactor for methionine synthase, the enzyme responsible for methylation of homocysteine to methionine using methyl tetrahydrofolate (methyl THF) as methyl donor (Fig. 4.3a); and, secondly, as deoxyadenosyl B₁₂ (ado B₁₂) it assists in conversion of methylmalonyl coenzyme A (CoA) to succinyl CoA (Fig. 4.3b). Assay of homocysteine and methylmalonic acid in plasma may be used as tests for B₁₂ deficiency (p. 52).





Fig. 4.3 The biochemical reactions of vitamin B₁₂ in humans. Ado B₁₂, deoxyadenosylcobalamin; CoA, coenzyme A; THF, tetrahydrofolate.

Fig. 4.4 The structure of folic (pteroylglutamic) acid. Dietary folates may contain: (a) additional hydrogen atoms at positions 7 and 8 (dihydrofolate) or 5, 6, 7 and 8 (tetrahydrofolate); (b) a formyl group at N₅ or N₁₀, a methyl group at N₅ or other 1-carbon groups; and (c) additional glutamate moiety attached to the γ -carboxyl group of the glutamate moiety.



Folate

Folic (pteroylglutamic) acid is the parent compound of a large group of compounds, the folates, that are derived from it (Fig. 4.4). Humans are unable to synthesize the folate structure and thus require preformed folate as a vitamin.

Absorption, transport and function

Dietary folates are converted to methyl THF (which, like folic acid, contains only one glutamate moiety) during absorption through the upper small intestine. Once inside the cell they are converted to folate polyglutamates (Fig. 4.5). Folate binding proteins are present on cell surfaces including the enterocyte and facilitate entry of reduced folates into cells. There is no specific plasma protein that enhances cellular folate uptake.

Folates are needed in a variety of biochemical reactions in the body involving single carbon unit transfer, in amino acid interconversions (e.g. homocysteine conversion to methionine) (Fig. 4.5) and serine to glycine or in synthesis of purine precursors of DNA.

Biochemical basis for megaloblastic anaemia (Fig. 4.5)

DNA is formed by polymerization of the four deoxyribonucleoside triphosphates. Folate deficiency is thought to cause megaloblastic anaemia by inhibiting thymidylate synthesis, a rate-limiting step in DNA synthesis in which thymidine monophosphate (dTMP) is synthesized, as this reaction needs 5,10-methylene THF polyglutamate as coenzyme.

All body cells, including those of the bone marrow, receive folate from plasma as methyl




Fig. 4.5 The biochemical basis of megaloblastic anaemia caused by vitamin B₁₂ or folate deficiency. Folate is required in one of its coenzyme forms, 5,10-methylene tetrahydrofolate (THF) polyglutamate, in the synthesis of thymidine monophosphate from its precursor deoxyuridine monophosphate. Vitamin B₁₂ is needed to convert methyl THF, which enters the cells from plasma, to THF, from which polyglutamate forms of folate are synthesized. Dietary folates are all converted to methyl THF (a monoglutamate) by the small intestine. A, adenine; C, cytosine; d, deoxyribose; DHF, dihydrofolate; DP, diphosphate; G, guanine; MP, monophosphate; T, thymine; TP, triphosphate; U, uracil.

THF. B_{12} is needed in the conversion of this methyl THF to THF, a reaction in which homocysteine is methylated to methionine. THF (but not methyl THF) is a substrate for folate polyglutamate synthesis inside cells. The folate polyglutamates act as intracellular folate coenzymes, including 5,10-methylene THF polyglutamate, the coenzyme form of folate involved in the thymidylate synthetase reaction (Fig. 4.5). Lack of B_{12} prevents the demethylation of methyl THF, thus depriving cells of THF and so of folate polyglutamate coenzymes.

Other congenital or acquired causes of megaloblastic anaemia (e.g. antimetabolite drug therapy) inhibit purine or pyrimidine synthesis at one or other step. The result is a reduced supply of one or other of the four precursors needed for DNA synthesis.

Folate reduction

During thymidylate synthesis, the folate polyglutamate coenzyme becomes oxidized from the THF state to dihydrofolate (DHF) (Fig. 4.5). Regeneration of active THF requires the enzyme DHF reductase. Inhibitors of this enzyme (e.g. methotrexate) therefore inhibit all folate coenzyme reactions, and so DNA synthesis (Fig. 4.5). Methotrexate is a useful drug, mainly in the treatment of malignant or inflammatory disease (e.g. of the skin) with excessive cell turnover. The weaker antagonist, pyrimethamine, is used primarily against malaria. Trimethoprim, active against bacterial DHF reductase but only very weakly against the human enzyme, is used alone or in combination with a sulfonamide, as co-trimoxazole. Toxicity caused by methotrexate or pyrimethamine is reversed by giving the reduced folate, folinic acid (5-formyl THF).

Vitamin B_{12} deficiency

In Western countries, severe deficiency is usually caused by (Addisonian) pernicious anaemia (Table 4.3). Less commonly, it may be caused by veganism in which the diet lacks B_{12} (usually in Hindu Indians), gastrectomy or small intestinal lesions. There is no syndrome of B_{12} deficiency as a result of increased utilization or loss of the vitamin. The deficiency takes at least 2 years to develop (i.e. the time needed for body stores to deplete at the rate of 1–2 $\mu\text{g/day}$) when there is severe malabsorption

Table 4.3 Causes of severe vitamin B_{12} deficiency.

Nutritional

Especially vegans

Malabsorption

Gastric causes

Pernicious anaemia

Congenital lack or abnormality of intrinsic factor

Total or partial gastrectomy

Intestinal causes

Intestinal stagnant loop syndrome—jejunal diverticulosis, blind-loop, stricture, etc.

Chronic tropical sprue

Ileal resection and Crohn's disease

Congenital selective malabsorption with proteinuria (autosomal recessive megaloblastic anaemia)

Fish tapeworm

Causes of mild vitamin B_{12} deficiency; other causes of malabsorption of vitamin B_{12} (e.g. malabsorption, atrophic gastritis, severe pancreatitis, gluten-induced enteropathy, HIV infection or therapy with metformin) do not usually lead to food vitamin B_{12} deficiency sufficient to cause anaemia or neuropathy.

of B_{12} from the diet. Nitrous oxide, however, may rapidly inactivate body B_{12} (p. 56).

Pernicious anaemia

This is caused by autoimmune attack on the gastric mucosa leading to atrophy of the stomach. The wall of the stomach becomes thin, with a plasma cell and lymphoid infiltrate of the lamina propria. Intestinal metaplasia may occur. There is achlorhydria and secretion of IF is absent or almost absent. Serum gastrin levels are raised. *Helicobacter pylori* infection may initiate an autoimmune gastritis which presents in younger subjects as iron deficiency and in the elderly as pernicious anaemia.

More females than males are affected (1.6 : 1), with a peak occurrence at 60 years, and there may be associated autoimmune disease including the autoimmune polyendocrine syndrome (Table 4.4). The disease is found in all races but is most common in northern Europeans and tends to occur in families. There is also an increased incidence of carcinoma of the stomach (approximately 2–3% of all cases of pernicious anaemia).

Table 4.4 Pernicious anaemia: associations.

Female	Vitiligo
Blue eyes	Myxoedema
Early greying	Hashimoto's disease
Northern European	Thyrotoxicosis
Familial	Addison's disease
Blood group A	Hypoparathyroidism
	Hypogammaglobulinaemia
	Carcinoma of the stomach

Antibodies

Ninety per cent of patients show parietal cell antibody in the serum directed against gastric H^+/K^+ -ATPase, and 50% type I or blocking antibody to IF which inhibits IF binding to B_{12} . Thirty-five per cent show a second (type II or precipitating) antibody to IF which inhibits its ileal binding site. IF antibodies are virtually specific for pernicious anaemia but occur in the serum of only half of patients, whereas the more common parietal cell antibody is less specific and occurs quite commonly in older subjects (e.g. 16% of normal women over 60 years).

Other causes of vitamin B_{12} deficiency

Congenital lack or abnormality of IF usually presents at approximately 2 years of age when stores of B_{12} that were derived from the mother *in utero* have been used up. There is also a form of autoimmune pernicious anaemia that presents in childhood. Specific malabsorption of B_{12} is brought about by genetic mutation of the IF- B_{12} receptor, cubilin or of amnionless which is involved in processing the IF- B_{12} complex. It usually presents in infancy or childhood and is associated with proteinuria in 90% of cases.

Lesser degrees of B_{12} deficiency occur resulting from inadequate intake of B_{12} , malabsorption of food B_{12} , atrophic gastritis (possibly triggered by *Helicobacter pylori*) without IF antibodies and other conditions listed in Table 4.3. Serum homocysteine and methylmalonic acid levels may be mildly raised and serum B_{12} levels subnormal but megaloblastic anaemia or neuropathy rarely occur.

Table 4.5 Causes of folate deficiency.

Nutritional

Especially old age, institutions, poverty, famine, special diets, goat's milk anaemia, etc.

Malabsorption

Tropical sprue, gluten-induced enteropathy (adult or child). Possible contributory factor to folate deficiency in some patients with partial gastrectomy, extensive jejunal resection or Crohn's disease

Excess utilization

Physiological

Pregnancy and lactation, prematurity

Pathological

Haematological diseases: haemolytic anaemias, myelofibrosis

Malignant disease: carcinoma, lymphoma, myeloma

Inflammatory diseases: Crohn's disease, tuberculosis, rheumatoid arthritis, psoriasis, exfoliative dermatitis, malaria

Excess urinary folate loss

Active liver disease, congestive heart failure

Drugs

Anticonvulsants, sulfasalazine

Mixed

Liver disease, alcoholism, intensive care

Folate deficiency

This is most often a result of a poor dietary intake of folate alone or in combination with a condition of increased folate utilization or malabsorption (Table 4.5). Excess cell turnover of any sort, including pregnancy, is the main cause of an increased need for folate, because the folate molecule becomes degraded when DNA synthesis is increased. The mechanism by which anticonvulsants and barbiturates cause the deficiency is still controversial. Alcohol, sulfasalazine and other drugs may have multiple effects on folate metabolism.

Clinical features of megaloblastic anaemia

The onset is usually insidious with gradually progressive symptoms and signs of anaemia (Chapter 2).

The patient may be mildly jaundiced (lemon yellow tint) (Fig. 4.6) because of the excess breakdown of haemoglobin resulting from increased ineffective erythropoiesis in the bone marrow. Glossitis (a beefy-red sore tongue) (Fig. 4.7), angular stomatitis (Fig. 4.8) and mild symptoms of malabsorption




Fig. 4.6 Megaloblastic anaemia: pallor and mild icterus in a patient with a haemoglobin count of 7.0 g/dL and a mean corpuscular volume of 132 fL.



Fig. 4.7 Megaloblastic anaemia: glossitis—the tongue is beefy-red and painful.



Fig. 4.8 Megaloblastic anaemia: angular cheilosis (stomatitis).

Table 4.6 Effects of vitamin B₁₂ or folate deficiency.

Megaloblastic anaemia
Macrocytosis of epithelial cell surfaces
Neuropathy (for vitamin B ₁₂ only)
Sterility
Rarely, reversible melanin skin pigmentation
Decreased osteoblast activity
Neural tube defects in the fetus are related to folate or B ₁₂ deficiency
Cardiovascular disease (see text and Chapter 25)

with loss of weight may be present because of the epithelial abnormality. Purpura as a result of thrombocytopenia and widespread melanin pigmentation (the cause of which is unclear) are less frequent presenting features (Table 4.6). Many asymptomatic patients are diagnosed when a blood count that has been performed for another reason reveals macrocytosis.



Fig. 4.9 Cross-section of the spinal cord in a patient who died with subacute combined degeneration of the cord (Weigert–Pal stain). There is demyelination of the dorsal and dorsolateral columns.

Vitamin B₁₂ neuropathy (subacute combined degeneration of the cord)

Severe B₁₂ deficiency can cause a progressive neuropathy affecting the peripheral sensory nerves and posterior and lateral columns (Fig. 4.9). The neuropathy is symmetrical and affects the lower limbs more than the upper limbs. The patient notices tingling in the feet, difficulty in walking and may fall over in the dark. Rarely, optic atrophy or severe psychiatric symptoms are present. Anaemia may be severe, mild or even absent, but the blood film and bone marrow appearances are always abnormal. The cause of the neuropathy is likely to be related to the accumulation of *S*-adenosyl homocysteine and reduced levels of *S*-adenosyl methionine in nervous tissue resulting in defective methylation of myelin and other substrates. The evidence that folate deficiency in the adult can cause a neuropathy is conflicting although there are some data suggesting it causes psychiatric changes.

Neural tube defect

Folate or B₁₂ deficiency in the mother predisposes to neural tube defect (NTD) (anencephaly, spina bifida or encephalocele) in the fetus (Fig. 4.10). The lower the maternal serum or red cell folate or serum B₁₂ levels (even when these are in the normal range), the higher the incidence of NTDs. Moreover,



Fig. 4.10 A baby with neural tube defect (spina bifida). (Courtesy of Professor C.J. Schorah)

supplementation of the diet with folic acid at the time of conception and in early pregnancy reduces the incidence of NTD by 75%. The exact mechanism is uncertain but is thought to be related to build-up of homocysteine and *S*-adenosyl homocysteine in the fetus which may impair methylation of various proteins and lipids. A common polymorphism in the enzyme 5,10-methylene tetrahydrofolate reductase (5,10-MTHFR) (677C → T) (p. 308) results in higher serum homocysteine and lower serum and red cell folate levels compared with controls. The incidence of the mutation is higher in the parents and fetus with NTD than in controls. Other studies suggest that an autoantibody against folate cell membrane receptors may occur more frequently than in controls in the sera of women who have a baby with NTD.

Other tissue abnormalities

Sterility is frequent in either sex with severe B₁₂ or folate deficiency. Macrocytosis, excess apoptosis and other morphological abnormalities of cervical, buccal, bladder and other epithelia occur. Widespread reversible melanin pigmentation may also occur. B₁₂ deficiency is associated with reduced osteoblastic activity. The associations of folate deficiency with cardiovascular and malignant diseases are discussed on page 56.

Laboratory findings

The anaemia is macrocytic (MCV >95 fL and often as high as 120–140 fL in severe cases) and the macrocytes are typically oval in shape (Fig. 4.11). The reticulocyte count is low and the total white cell and platelet counts may be moderately reduced, especially in severely anaemic patients. A proportion of the neutrophils show hypersegmented nuclei (with



Fig. 4.11 Megaloblastic anaemia: peripheral blood film showing oval macrocytes.

six or more lobes). The bone marrow is usually hypercellular and the erythroblasts are large and show failure of nuclear maturation maintaining an open, fine, lacy primitive chromatin pattern but normal haemoglobinization (Fig. 4.12). Giant and abnormally shaped metamyelocytes are characteristic.

The serum unconjugated bilirubin and lactate dehydrogenase (LDH) are raised as a result of marrow cell breakdown.

Diagnosis of vitamin B₁₂ or folate deficiency

It is usual to assay serum B₁₂, and serum and red cell folate (Table 4.7). The serum B₁₂ is low in megaloblastic anaemia or neuropathy caused by B₁₂ deficiency. The serum and red cell folate are both low in megaloblastic anaemia caused by folate deficiency. In B₁₂ deficiency, the serum folate tends to rise but the red cell folate falls. In the absence of B₁₂ deficiency, however, the red cell folate is a more accurate guide than the serum folate of tissue folate status. Measurement of serum methylmalonic acid is a test for B₁₂ deficiency and of homocysteine for folate or B₁₂ deficiency. These are not specific and it is difficult to establish normal levels in different age groups. These tests are not widely available.

Tests for cause of vitamin B₁₂ or folate deficiency

For B₁₂ deficiency, absorption tests (Table 4.8) using an oral dose of radioactive cobalt (⁵⁷Co)-labelled cyanocobalamin are valuable in distinguishing malabsorption from an inadequate diet. When the test is repeated with an active IF preparation, gastric lesions

Table 4.7 Laboratory tests for vitamin B₁₂ and folate deficiency.

Test	Normal values*		Result in	
			Vitamin B ₁₂ deficiency	Folate deficiency
Serum vitamin B ₁₂	160–925 ng/L	120–680 pmol/L	Low	Normal or borderline
Serum folate	3.0–15.0 µg/L	4–30 nmol/L	Normal or raised	Low
Red cell folate	160–640 µg/L	360–1460 nmol/L	Normal or low	Low

* Normal values differ slightly with different commercial kits.




Fig. 4.12 Megaloblastic changes in the bone marrow in a patient with severe megaloblastic anaemia. (a–c) Erythroblasts showing fine, open stippled (primitive) appearance of the nuclear chromatin even in late cells (pale cytoplasm with some haemoglobin formation). (d) Abnormal giant metamyelocytes and band forms.

such as those associated with pernicious anaemia can be distinguished from intestinal lesions (Table 4.9). Absorption is most frequently measured indirectly by the urinary excretion (Schilling) technique in which absorbed radiolabelled B_{12} is 'flushed' into a 24-h urine sample by a large (1 mg) dose of non-radioactive B_{12} given simultaneously with the labelled oral dose. Problems with using human IF,

non-virally inactivated and availability of radioactive B_{12} have led to less frequent use of the test.

Other useful tests are listed in Table 4.8. These are mainly concerned with assessing gastric function and testing for antibodies to gastric antigens. In all cases of pernicious anaemia endoscopy studies should be performed to confirm the presence of gastric atrophy and exclude carcinoma of the stomach.

Table 4.8 Tests for cause of vitamin B₁₂ or folate deficiency.

Vitamin B ₁₂	Folate
Diet history	Diet history
B ₁₂ absorption ± IF	Tests for intestinal malabsorption
IF, parietal cell antibodies	Anti-transglutaminase and endomysial antibodies
Endoscopy or barium meal and follow through	Duodenal biopsy
	Underlying disease

IF, intrinsic factor.

Table 4.9 Results of absorption tests of radioactive vitamin B₁₂.

	Dose of labelled B ₁₂ given alone	Dose of labelled B ₁₂ given with IF
Vegan	Normal	Normal
Pernicious anaemia or gastrectomy	Low	Normal
Ileal lesion	Low	Low
Intestinal blind-loop syndrome	Low*	Low*

IF, intrinsic factor.

* Corrected by antibiotic therapy.

For folate deficiency, the dietary history is most important, although it is difficult to estimate folate intake accurately. Unsuspected gluten-induced enteropathy or other underlying conditions should also be considered (Table 4.5).

Treatment

Most cases only need therapy with the appropriate

vitamin (Table 4.10). If large doses of folic acid (e.g. 5 mg/day) are given in B₁₂ deficiency they cause a haematological response but may aggravate the neuropathy. They should therefore not be given alone unless B₁₂ deficiency has been excluded. In severely anaemic patients who need treatment urgently it may be safer to initiate treatment with both vitamins after blood has been taken for B₁₂ and folate examinations and a bone marrow test has

Table 4.10 Treatment of megaloblastic anaemia.

	Vitamin B ₁₂ deficiency	Folate deficiency
Compound	Hydroxocobalamin	Folic acid
Route	Intramuscular*	Oral
Dose	1000 µg	5 mg
Initial dose	6 × 1000 µg over 2–3 weeks	Daily for 4 months
Maintenance	1000 µg every 3 months	Depends on underlying disease; life-long therapy may be needed in chronic inherited haemolytic anaemias, myelofibrosis, renal dialysis
Prophylactic	Total gastrectomy Ileal resection	Pregnancy, severe haemolytic anaemias, dialysis, prematurity

* Some authors have recommended daily oral or sublingual therapy of vitamin B₁₂ deficiency.




Fig. 4.13 Typical haematological response to vitamin B₁₂ (hydroxocobalamin) therapy in pernicious anaemia. Hb, haemoglobin; Retics, reticulocytes; WBC, white blood cells.

been performed. In the elderly, the presence of heart failure should be corrected with diuretics and oral potassium supplements given for 10 days (because hypokalaemia has been found to occur during the response in some cases). Blood transfusion should be avoided if possible as it may cause circulatory overload.

Response to therapy

The patient feels better after 24–48 h of correct vitamin therapy with increased appetite and well-being. The haemoglobin should rise by 2–3 g/dL each fortnight. The white cell and platelet counts become normal in 7–10 days (Fig. 4.13) and the marrow is normoblastic in about 48 h, although giant metamyelocytes persist for up to 12 days.

The peripheral neuropathy may partly improve but spinal cord damage is irreversible. The shorter the history of neurological symptoms, the greater the chance of recovery.

Prophylactic therapy

Vitamin B₁₂ is given to patients who have total gastrectomy or ileal resection. Folic acid is given in pregnancy at a recommended dosage of 400 µg/day and all women of child-bearing age are recommended to have an intake of at least 400 µg/day (by increased intake of folate-rich or folate-supplemented foods or as folic acid) to prevent a first occurrence of an NTD in the fetus. Folic acid is also given to patients undergoing chronic dialysis and with severe haemolytic anaemias and chronic myelofibrosis, and to premature babies. Food forti-

fication with folic acid (e.g. in flour) is currently recommended in the UK to reduce the incidence of NTDs and is already practised in over 40 countries including North America.

Other megaloblastic anaemias

See Table 4.1.

Abnormalities of vitamin B₁₂ or folate metabolism

These include congenital deficiencies of enzymes concerned in B₁₂ or folate metabolism or of the serum transport protein for B₁₂, TC. Nitrous oxide (N₂O) anaesthesia causes rapid inactivation of body B₁₂ by oxidizing the reduced cobalt atom of methyl B₁₂. Megaloblastic marrow changes occur with several days of N₂O administration and can cause pancytopenia. Chronic exposure (as in dentists and anaesthetists) has been associated with neurological damage resembling B₁₂ deficiency neuropathy. Antifolate drugs, particularly those which inhibit DHF reductase (e.g. methotrexate and pyrimethamine) may also cause megaloblastic change. Trimethoprim, which inhibits bacterial DHF reductase, has only a slight action against the human enzyme and causes megaloblastic change only in patients already B₁₂ or folate deficient.

Systemic diseases associated with folate or vitamin B₁₂ deficiency

Cardiovascular diseases

Raised serum homocysteine levels are associated with an increased incidence of myocardial infarct, peripheral and cerebral vascular disease and venous thrombosis (Chapter 25). Raised serum homocysteine levels are associated with low serum and red cell folate and low serum B₁₂ or vitamin B₆ levels. In addition, homocysteine levels tend to be higher in men than in premenopausal women, in old age, in heavy smokers and those with excess alcohol consumption, with impaired renal function and with some drugs. Although folate deficiency (and in some studies, the presence of the polymorphism

in the 5,10-MTHFR gene) has been associated with an increased incidence of cardiovascular disease, recent large randomised studies have not shown a reduction in the rate of myocardial infarction or stroke by the use of prophylactic folic acid.

Malignant diseases

Various associations have been found between folate status or polymorphisms in folate metabolizing enzymes and malignant diseases such as colon or breast cancer and acute lymphoblastic leukaemia in childhood. In most but not all, reduced folate status has been associated with an increased risk of malignancy.

Defects of DNA synthesis not related to vitamin B₁₂ or folate

Congenital deficiency of one or other enzyme concerned in purine or pyrimidine synthesis can cause megaloblastic anaemia identical in appearance to that caused by a deficiency of B₁₂ or folate. The best known is orotic aciduria. Therapy with drugs that inhibit purine or pyrimidine synthesis (such as hydroxyurea, cytosine arabinoside, 6-mercaptopurine and zidovudine (AZT)) and some forms of acute myeloid leukaemia or myelodysplasia also cause megaloblastic anaemia.

Other macrocytic anaemias

There are many non-megaloblastic causes of macrocytic anaemia (Table 4.11). The exact mechanisms creating large red cells in each of these conditions is not clear although increased lipid deposition on the red cell membrane or alterations of erythroblast maturation time in the marrow may be implicated. Alcohol is the most frequent cause of a raised MCV in the absence of anaemia. Reticulocytes are bigger than mature red cells and so haemolytic anaemia is an important cause of macrocytic anaemia. The other underlying conditions listed in Table 4.11 are usually easily diagnosed provided that they are considered and the appropriate investigations to exclude B₁₂ or folate deficiency are carried out.

Table 4.11 Causes of macrocytosis other than megaloblastic anaemia.

Alcohol
Liver disease
Myxoedema
Myelodysplastic syndromes
Cytotoxic drugs
Aplastic anaemia
Pregnancy
Smoking
Reticulocytosis
Myeloma and paraproteinaemia
Neonatal

Differential diagnosis of macrocytic anaemias

The clinical history and physical examination may suggest B₁₂ or folate deficiency as the cause. Diet, drugs, alcohol intake, family history, history suggestive of malabsorption, presence of autoimmune diseases or other associations with pernicious anaemia (Table 4.4), previous gastrointestinal disease or operations are all important. The presence of jaundice, glossitis or a neuropathy are also valuable indications of megaloblastic anaemia.

The laboratory features of particular importance are the shape of macrocytes (oval in megaloblastic anaemia), the presence of hypersegmented neutrophils and of leucopenia and thrombocytopenia in megaloblastic anaemia and the bone marrow appearance. Assay of B₁₂ and folate is straightforward. Exclusion of alcoholism (particularly if the patient is not anaemic), liver and thyroid function tests, and bone marrow examination for myelodysplasia, aplasia or myeloma are important in the investigation of macrocytosis not caused by B₁₂ or folate deficiency.

Bibliography

- Chan J.C.W., Liu H.S.Y., Kho B.C.S. *et al.* (2006) Pernicious anemia in Chinese: A study of 181 patients in a Hong Kong hospital. *Medicine* 85,129–38.
- Clarke R., Grimley Evans J., Schneede J. *et al.* (2004) Vitamin B₁₂ and folate deficiency in later life. *Age Ageing* 33,34–41.
- Czeizel A.E., Dobo M. and Vargha P. (2004) Hungarian cohort controlled trial of periconceptional multi-vitamin supplementation shows a reduction in certain congenital abnormalities. *Birth Defects Research* 70,853–61.
- Eicholzer M., Tonz O. and Zimmerman R. (2006) Folic acid: a public health challenge. *Lancet* 367,1352–61.
- Fyfe J.C., Madsen M., Hojrup P. *et al.* (2004) The functional cobalamin (vitamin B₁₂)-intrinsic factor receptor is a novel complex of cubilin and amnionless. *Blood* 103,1573–9.
- He Q., Masden M., Kilkenney A. *et al.* (2005) Amnionless function is required for cubilin brush-border expression and intrinsic factor-cobalamin (vitamin B₁₂) absorption *in vivo*. *Blood* 106,1447–53.
- Hershko C., Ronson A., Souroujon M. *et al.* (2006) Variable hematologic presentation of autoimmune gastritis: age-related progression from iron deficiency to cobalamin depletion. *Blood* 107,1673–9.
- The Hope Investigators (2006) Homocysteine lowering with folic acid and B vitamins in vascular disease. *N Engl J Med* 354,1567–77.
- Hvas A.-M., Buhl T., Laursen N.B. *et al.* (2006) The effect of recombinant human intrinsic factor on the uptake of vitamin B₁₂ in patients with evident vitamin B₁₂ deficiency. *Haematologica* 91,805–8.
- Marks P.W. and Zukerberg L.R. (2004) Case records of Massachusetts General Hospital. Weekly clinicopathological exercises. Case 30-2004. A 37-year-old woman with paresthesiae of the arms and legs. *N Engl J Med* 351,1333–41.
- Rothenberg S.P., da Costa M.P., Sequeira J.M. *et al.* (2004) Autoantibodies against folate receptors in women with a pregnancy complicated by a neural-tube defect. *N Engl J Med* 350,134–42.
- Solomon L.R. (2005) Cobalamin-responsive disorders in the ambulatory care setting: unreliability of cobalamin, methylmalonic acid, and homocysteine testing. *Blood* 105,978–85.
- Whitehead V.M. (2006) Acquired and inherited disorders of cobalamin and folate in children. *Br J Haematol* 134,125–36.
- Wickramasinghe S.N. (ed.) (1995) Megaloblastic anaemia. *Clin Haematol* 8,441–703.

CHAPTER 5

5

Haemolytic anaemias

Normal red cell destruction, 58
Introduction to haemolytic anaemias, 58
Intravascular and extravascular haemolysis, 60

Hereditary haemolytic anaemias, 61
Acquired haemolytic anaemias, 66
Bibliography, 70

Normal red cell destruction

Red cell destruction usually occurs after a mean lifespan of 120 days when the cells are removed extravascularly by the macrophages of the reticuloendothelial (RE) system, especially in the marrow but also in the liver and spleen. As the cells have no nucleus, red cell metabolism gradually deteriorates as enzymes are degraded and not replaced and the cells become non-viable. The breakdown of haem from red cells liberates iron for recirculation via plasma transferrin to marrow erythroblasts, and protoporphyrin which is broken down to bilirubin. Bilirubin circulates to the liver where it is conjugated to glucuronides which are excreted into the gut via bile and converted to stercobilinogen and stercobilin (excreted in faeces) (Fig. 5.1). Stercobilinogen and stercobilin are partly reabsorbed and excreted in urine as urobilinogen and urobilin. Globin chains are broken down to amino acids which are reutilized for general protein synthesis in the body. Haptoglobins are proteins present in normal plasma capable of binding haemoglobin. The haemoglobin-haptoglobin complex is removed from plasma by the RE system. Intravascular haemolysis (breakdown of red cells within blood vessels) plays little or no part in normal red cell destruction.

Introduction to haemolytic anaemias

Haemolytic anaemias are defined as those anaemias

that result from an increase in the rate of red cell destruction. Because of erythropoietic hyperplasia and anatomical extension of bone marrow, red cell destruction may be increased several-fold before the patient becomes anaemic—compensated haemolytic disease. The normal adult marrow, after full expansion, is able to produce red cells at 6–8 times the normal rate provided this is ‘effective’. It leads to a marked reticulocytosis, particularly in the more anaemic cases. Therefore, haemolytic anaemia may not be seen until the red cell lifespan is less than 30 days.

Classification

Table 5.1 is a simplified classification of the haemolytic anaemias. Hereditary haemolytic anaemias are the result of ‘intrinsic’ red cell defects whereas acquired haemolytic anaemias are usually the result of an ‘extracorporeal’ or ‘environmental’ change. Paroxysmal nocturnal haemoglobinuria (PNH) is the exception because although it is an acquired disorder the PNH red cells have an intrinsic defect.

Clinical features

The patient may show pallor of the mucous membranes, mild fluctuating jaundice and splenomegaly. There is no bilirubin in urine but this may turn dark on standing because of excess urobilinogen. Pigment (bilirubin) gallstones may complicate




Fig. 5.1 (a) Normal red blood cell (RBC) breakdown. This takes place extravascularly in the macrophages of the reticuloendothelial system. (b) Intravascular haemolysis occurs in some pathological disorders.

the condition (Fig. 5.2) and some patients (particularly with sickle cell disease) develop ulcers around the ankle. Aplastic crises may occur, usually precipitated by infection with parvovirus which 'switches off' erythropoiesis, and are characterized by a sudden increase in anaemia and drop in reticulocyte count (see Fig. 20.4).

Rarely, folate deficiency may cause an aplastic crisis in which the bone marrow is megaloblastic.

Laboratory findings

The laboratory findings are conveniently divided into three groups.

1 Features of increased red cell breakdown:

- (a) serum bilirubin raised, unconjugated and bound to albumin;

- (b) urine urobilinogen increased;
- (c) faecal stercobilinogen increased;
- (d) serum haptoglobins absent because the haptoglobins become saturated with haemoglobin and the complex is removed by RE cells.

2 Features of increased red cell production:

- (a) reticulocytosis;
- (b) bone marrow erythroid hyperplasia; the normal marrow myeloid : erythroid ratio of 2 : 1 to 12 : 1 is reduced to 1 : 1 or reversed.

3 Damaged red cells:

- (a) morphology (e.g. microspherocytes, elliptocytes, fragments);
- (b) osmotic fragility, autohaemolysis, etc.;
- (c) red cell survival shortened; this was shown by ^{51}Cr labelling with study of the sites of destruction. This test is now not widely available.

Table 5.1 Classification of haemolytic anaemias.

Hereditary	Acquired
Membrane	Immune
Hereditary spherocytosis, hereditary elliptocytosis	<i>Autoimmune</i>
	Warm antibody type (see Table 5.5)
Metabolism	Cold antibody type
G6PD deficiency, pyruvate kinase deficiency	<i>Alloimmune</i>
Haemoglobin	Haemolytic transfusion reactions
Genetic abnormalities (Hb S, Hb C, unstable); see Chapter 6	Haemolytic disease of the newborn
	Allografts, especially marrow transplantation
	<i>Drug associated</i>
	Red cell fragmentation syndromes
	See Table 5.6
	March haemoglobinuria
	Infections
	Malaria, clostridia
	Chemical and physical agents
	Especially drugs, industrial/domestic substances, burns
	Secondary
	Liver and renal disease
	Paroxysmal nocturnal haemoglobinuria

G6PD, glucose-6-phosphate dehydrogenase; Hb, haemoglobin.



Fig. 5.2 Ultrasound of pigment gallstones (arrowed) in a 6-year-old male patient with hereditary spherocytosis. (Courtesy of L. Berger)

Intravascular and extravascular haemolysis

There are two main mechanisms whereby red cells are destroyed in haemolytic anaemia. There may be excessive removal of red cells by cells of the RE system (extravascular haemolysis) or they may be broken down directly in the circulation in a process known as intravascular haemolysis (Fig. 5.1; Table 5.2). Whichever mechanism dominates will depend on the pathology involved. In intravascular haemolysis, free haemoglobin is released which rapidly saturates plasma haptoglobins and the excess free haemoglobin is filtered by the glomerulus. If the rate of haemolysis saturates the renal tubular reabsorptive capacity, free haemoglobin enters urine (Fig. 5.3) and, as iron is released, the renal tubules become loaded with haemosiderin. Methaemalbumin and haemopexin are also formed from the process of intravascular haemolysis.

Table 5.2 Causes of intravascular haemolysis.

Mismatched blood transfusion (usually ABO)
G6PD deficiency with oxidant stress
Red cell fragmentation syndromes
Some autoimmune haemolytic anaemias
Some drug- and infection-induced haemolytic anaemias
Paroxysmal nocturnal haemoglobinuria
March haemoglobinuria
Unstable haemoglobin

G6PD, glucose-6-phosphate dehydrogenase

The main laboratory features of intravascular haemolysis are as follows.

- 1 Haemoglobinaemia and haemoglobinuria.
- 2 Haemosiderinuria (iron storage protein in the spun deposit of urine).

3 Methaemalbuminaemia (detected spectrophotometrically by Schumm's test).

Hereditary haemolytic anaemias

Membrane defects

Hereditary spherocytosis

Hereditary spherocytosis (HS) is the most common hereditary haemolytic anaemia in northern Europeans.

Pathogenesis

HS is usually caused by defects in the proteins involved in the vertical interactions between the membrane skeleton and the lipid bilayer of the red cell (Table 5.3; see Fig. 2.12). The loss of membrane may be caused by the release of parts of the lipid




Fig. 5.3 (a) Progressive urine samples in an acute episode of intravascular haemolysis showing haemoglobinuria of decreasing severity. (b) Prussian blue-positive deposits of haemosiderin in a urine spun deposit (Perls' stain).



(a)

(b)

Fig. 5.4 (a) Blood film in hereditary spherocytosis. The spherocytes are deeply staining and of small diameter. Larger polychromatic cells are reticulocytes (confirmed by supravital staining). (b) Blood film in hereditary elliptocytosis.

bilayer that are not supported by the skeleton. The marrow produces red cells of normal biconcave shape but these lose membrane and become increasingly spherical (loss of surface area relative to volume) as they circulate through the spleen and the rest of the RE system. Ultimately, the spherocytes are unable to pass through the splenic microcirculation where they die prematurely.

Clinical features

The inheritance is autosomal dominant with variable expression; rarely it may be autosomal recessive. The anaemia can present at any age from infancy to old age. Jaundice is typically fluctuating and is particularly marked if the haemolytic anaemia is associated with Gilbert's disease (a defect of hepatic conjugation of bilirubin); splenomegaly occurs in most patients. Pigment gallstones are frequent (Fig. 5.2); aplastic crises, usually precipitated by parvovirus infection, may cause a sudden increase in severity of anaemia (see Fig. 20.4).

Haematological findings

Anaemia is usual but not invariable; its severity tends to be similar in members of the same family. Reticulocytes are usually 5–20%. The blood film shows microspherocytes (Fig. 5.4a) which are densely staining with smaller diameters than normal red cells.

Investigation and treatment

The classic finding is that the osmotic fragility is increased (Fig. 5.5). The abnormality may require 24-h incubation at 37°C to become obvious. Autohaemolysis is increased and corrected by glucose. The cells are incubated with their own plasma for 48 h with or without glucose. The direct antiglobulin (Coombs') test is normal, excluding an autoimmune cause of spherocytosis and haemolysis.

The principal form of treatment is splenectomy




Fig. 5.5 The osmotic fragility in hereditary spherocytosis. The curve is shifted to the right of the normal range (in yellow), but a tail of more resistant cells (reticulocytes) is also present.

although this should not be performed unless clinically indicated because of anaemia or gallstones because of the risk of post-splenectomy sepsis, particularly in early childhood (p. 127). Splenectomy should always produce a rise in the haemoglobin level to normal, even though microspherocytes formed in the rest of the RE system will remain. Folic acid is given in severe cases to prevent folate deficiency.

Hereditary elliptocytosis

This has similar clinical and laboratory features to HS except for the appearance of the blood film (Fig. 5.4b), but it is usually a clinically milder disorder. It is usually discovered by chance on a blood film and there may be no evidence of haemolysis. Occasional patients require splenectomy. The basic defect is a failure of spectrin heterodimers to self-associate into heterotetramers. A number of genetic mutations affecting horizontal interactions have been detected (Table 5.3). Patients with homozygous or doubly heterozygous elliptocytosis present with a severe haemolytic anaemia with microspherocytes, poikilocytes and splenomegaly (hereditary pyropoikilocytosis).

South-East Asian ovalocytosis

This is common in Melanesia, Malaysia, Indonesia and the Philippines and is caused by a nine amino acid deletion at the junction of the cytoplasmic and transmembrane domains of the band 3 protein.

Table 5.3 Molecular basis of hereditary spherocytosis and elliptocytosis.

Hereditary spherocytosis

Ankyrin deficiency or abnormalities
Spectrin deficiency or abnormalities
Band 3 abnormalities
Pallidin (protein 4.2) abnormalities

Hereditary elliptocytosis

α - or β -spectrin mutants leading to defective spectrin dimer formation
 α - or β -spectrin mutants leading to defective spectrin-ankyrin associations
Protein 4.1 deficiency or abnormality

South-East Asian ovalocytosis band 3 deletion

The cells are rigid and resist invasion by malarial parasites. Most cases are not anaemic and are asymptomatic.

Defective red cell metabolism

Glucose-6-phosphate dehydrogenase deficiency

Glucose-6-phosphate dehydrogenase (G6PD) functions to reduce nicotinamide adenine dinucleotide phosphate (NADP) while oxidizing glucose-6-phosphate. It is the only source of NADPH in red cells and as NADPH is needed for the production of reduced glutathione a deficiency renders the red cell susceptible to oxidant stress (Fig. 5.6).

Epidemiology

There is a wide variety of normal genetic variants of the enzyme G6PD, the most common being type B (Western) and type A in Africans. In addition, more than 400 variants caused by point mutations or deletions of the enzyme G6PD have been characterized




Fig. 5.6 Haemoglobin and red blood cell (RBC) membranes are usually protected from oxidant stress by reduced glutathione (GSH). In G6PD deficiency, NADPH and GSH synthesis is impaired. F6P, fructose-6-phosphate; G6P, glucose-6-phosphate; G6PD, glucose-6-phosphate dehydrogenase; GSSG, glutathione (oxidized form); NADP, NADPH, nicotinamide adenine dinucleotide phosphate.




Fig. 5.7 Global distribution of *G6PD* gene variants causing *G6PD* deficiency. Shaded areas indicate the prevalence of *G6PD* deficiency. (From Luzzatto & Notaro 2001, *Science* 293: 442, with permission)

which show less activity than normal and world-wide over 400 million people are *G6PD* deficient in enzyme activity (Fig. 5.7).

The inheritance is sex-linked, affecting males, and carried by females who show approximately half the normal red cell *G6PD* values. The female heterozygotes have an advantage of resistance to *Falciparum* malaria. The main races affected are in West Africa, the Mediterranean, the Middle East and South-East Asia. The degree of deficiency varies, often being mild (10–15% of normal activity) in black Africans, more severe in Orientals and most severe in Mediterraneans. Severe deficiency occurs occasionally in white people.

Clinical features

G6PD deficiency is usually asymptomatic. Although *G6PD* is present in all cells, the main syndromes that occur are as follows. (i) Acute haemolytic anaemia in response to oxidant stress, e.g. drugs, fava beans or infections (Table 5.4). The acute haemolytic anaemia is caused by rapidly developing intravascular haemolysis with haemoglobinuria (Fig. 5.3a). The anaemia may be self-limiting as new young red

Table 5.4 Agents that may cause haemolytic anaemia in glucose-6-phosphate dehydrogenase (*G6PD*) deficiency.

Infections and other acute illnesses (e.g. diabetic ketoacidosis)

Drugs

Antimalarials (e.g. primaquine, pamaquine, chloroquine, Fansidar, Maloprim)

Sulphonamides and sulphones (e.g. co-trimoxazole, sulfanilamide, dapsone, Salazopyrin)

Other antibacterial agents (e.g. nitrofurans, chloramphenicol)

Analgesics (e.g. aspirin), moderate doses are safe

Antihelmintics (e.g. β -naphthol, stibophen)

Miscellaneous (e.g. vitamin K analogues, naphthalene (mothballs), probenecid)

Fava beans (possibly other vegetables)

NB. Many common drugs have been reported to precipitate haemolysis in *G6PD* deficiency in some patients (e.g. aspirin, quinine and penicillin) but not at conventional dosage.

Fig. 5.8 Blood film in G6PD deficiency with acute haemolysis after an oxidant stress. Some of the cells show loss of cytoplasm with separation of remaining haemoglobin from the cell membrane ('blister' cells). There are also numerous contracted and deeply staining cells. Supravital staining (as for reticulocytes) showed the presence of Heinz bodies (see Fig. 2.17).



cells are made with near normal enzyme levels. (ii) Neonatal jaundice. (iii) Rarely, a congenital non-spherocytic haemolytic anaemia. These may result from different types of enzyme deficiency.

Diagnosis

Between crises the blood count is normal. The enzyme deficiency is detected by one of a number of screening tests or by direct enzyme assay on red cells. During a crisis the blood film may show contracted and fragmented cells, 'bite' cells and 'blister' cells (Fig. 5.8) which have had Heinz bodies removed by the spleen. Heinz bodies (oxidized, denatured haemoglobin) may be seen in the reticulocyte preparation, particularly if the spleen is absent. There are also features of intravascular haemolysis. Because of the higher enzyme level in young red cells, red cell enzyme assay may give a 'false' normal level in the phase of acute haemolysis with a reticulocyte response. Subsequent assay after the acute phase reveals the low G6PD level when the red cell population is of normal age distribution.

Treatment

The offending drug is stopped, any underlying infection is treated, a high urine output is maintained and blood transfusion undertaken where necessary for severe anaemia. G6PD-deficient babies are prone to neonatal jaundice and in severe cases phototherapy and exchange transfusion may

be needed. The jaundice is usually not caused by excess haemolysis but by deficiency of G6PD affecting neonatal liver function.

Glutathione deficiency and other syndromes

Other defects in the pentose phosphate pathway leading to similar syndromes to G6PD deficiency have been described—particularly glutathione deficiency.

Glycolytic (Embden–Meyerhof) pathway defects

These are all uncommon and lead to a congenital non-spherocytic haemolytic anaemia. In some there are defects of other systems (e.g. a myopathy). The most frequently encountered is pyruvate kinase (PK) deficiency.

Pyruvate kinase deficiency

This is inherited as an autosomal recessive, the affected patients being homozygous or doubly heterozygous. Over 100 different mutations have been described. The red cells become rigid as a result of reduced adenosine triphosphate (ATP) formation. The severity of the anaemia varies widely (haemoglobin 4–10 g/dL) and causes relatively mild symptoms because of a shift to the right in the oxygen (O_2) dissociation curve caused by a rise in intracellular 2,3-diphosphoglycerate (2,3-DPG). Clinically, jaundice is usual and gallstones frequent. Frontal bossing may be present. The blood

Table 5.5 Immune haemolytic anaemias: classification.

Warm type	Cold type
Autoimmune	
<i>Idiopathic</i>	<i>Idiopathic</i>
<i>Secondary</i>	<i>Secondary</i>
SLE, other 'autoimmune' diseases	Infections— <i>Mycoplasma pneumoniae</i> , infectious mononucleosis
CLL, lymphomas	Lymphoma
Drugs (e.g. methyldopa)	Paroxysmal cold haemoglobinuria (rare, sometimes associated with infections, e.g. syphilis)
Alloimmune	
<i>Induced by red cell antigens</i>	
Haemolytic transfusion reactions	
Haemolytic disease of the newborn post stem cell grafts	
<i>Drug induced</i>	
Drug–red cell membrane complex	
Immune complex	

CLL, chronic lymphocytic leukaemia; SLE, systemic lupus erythematosus.

film shows poikilocytosis and distorted 'prickle' cells, particularly post-splenectomy. Laboratory tests show that autohaemolysis is increased but, in contrast to HS, it is not corrected by glucose; direct enzyme assay is needed to make the diagnosis. Splenectomy may alleviate the anaemia but does not cure it and is indicated in those patients who need frequent transfusions.

Hereditary disorders of haemoglobin synthesis

Several of these cause clinical haemolysis. They are discussed in Chapter 6.

Acquired haemolytic anaemias

Immune haemolytic anaemias

Autoimmune haemolytic anaemias

Autoimmune haemolytic anaemias (AIHAs) are caused by antibody production by the body against its own red cells. They are characterized by a positive direct antiglobulin test (DAT) also known as the Coombs' test (see Fig. 27.5) and divided into 'warm' and 'cold' types (Table 5.5) according to whether the antibody reacts more strongly with red cells at 37°C or 4°C.

Warm autoimmune haemolytic anaemias

The red cells are coated with immunoglobulin (Ig), usually immunoglobulin G (IgG) alone or with complement, and are therefore taken up by RE macrophages which have receptors for the Ig Fc fragment. Part of the coated membrane is lost so the cell becomes progressively more spherical to maintain the same volume and is ultimately prematurely destroyed, predominantly in the spleen. When the cells are coated with IgG and complement (C3d, the degraded fragment of C3) or complement alone, red cell destruction occurs more generally in the RE system.

Clinical features

The disease may occur at any age, in either sex, and presents as a haemolytic anaemia of varying severity. The spleen is often enlarged. The disease tends to remit and relapse. It may occur alone or in association with other diseases, or arise in some patients as a result of methyldopa therapy (Table 5.5). When associated with idiopathic thrombocytopenic purpura (ITP), which is a similar condition affecting platelets (p. 281), it is known as Evans' syndrome. When secondary to systemic lupus erythematosus the cells typically are coated with immunoglobulin and complement.




Fig. 5.9 (a) Blood film in warm autoimmune haemolytic anaemia. Numerous microspherocytes are present and larger polychromatic cells (reticulocytes). (b) Blood film in cold autoimmune haemolytic anaemia. Marked red cell agglutination is present in films made at room temperature. The background is caused by the raised plasma protein concentration.

Laboratory findings

The haematological and biochemical findings are typical of an extravascular haemolytic anaemia with spherocytosis prominent in the peripheral blood (Fig. 5.9a). The DAT is positive as a result of IgG, IgG and complement or IgA on the cells and, in some cases, the autoantibody shows specificity within the rhesus system. The antibodies both on the cell surface and free in serum are best detected at 37°C.

Treatment

- 1 Remove the underlying cause (e.g. methyl dopa).
- 2 Corticosteroids. Prednisolone is the usual first-line treatment; 60 mg/day is a typical starting dose in adults and should then be tapered down. Those with predominantly IgG on red cells do best whereas those with complement often respond poorly, both to corticosteroids and splenectomy.
- 3 Splenectomy may be of value in those who fail to respond well or fail to maintain a satisfactory haemoglobin level on an acceptably small steroid dosage.
- 4 Immunosuppression may be tried after other measures have failed. Azathioprine, cyclophos-

phamide, chlorambucil, ciclosporin and mycophenolate mofetil have been tried.

5 Monoclonal antibodies. Anti-CD20 (rituximab) has produced prolonged remissions in a proportion of cases and anti-CD52 (Campath-1H) has been tried successfully in a few cases.

6 Folic acid is given to severe cases.

7 Blood transfusion may be needed if anaemia is severe and causing symptoms. The blood should be the least incompatible and if the specificity of the autoantibody is known, donor blood is chosen that lacks the relevant antigen(s). The patients also readily make alloantibodies against donor red cells.

8 High-dose immunoglobulin has been used but with less success than in ITP (p. 283).

Cold autoimmune haemolytic anaemias

In these syndromes the autoantibody, whether monoclonal (as in the idiopathic cold haemagglutinin syndrome or associated with lymphoproliferative disorders) or polyclonal (as following infection, e.g. infectious mononucleosis or *Mycoplasma pneumoniae*) attaches to red cells mainly in the peripheral circulation where the blood temperature is cooled

(Table 5.5). The antibody is usually IgM and binds to red cells best at 4°C. IgM antibodies are highly efficient at fixing complement and both intravascular and extravascular haemolysis can occur. Complement alone is usually detected on the red cells, the antibody having eluted off the cells in warmer parts of the circulation. Interestingly, in nearly all these cold AIHA syndrome the antibody is directed against the 'I' antigen on the red cell surface. In infectious mononucleosis it is anti-i.

Clinical features

The patient may have a chronic haemolytic anaemia aggravated by the cold and often associated with intravascular haemolysis. Mild jaundice and splenomegaly may be present. The patient may develop acrocyanosis (purplish skin discoloration) at the tip of the nose, ears, fingers and toes caused by the agglutination of red cells in small vessels.

Laboratory findings are similar to those of warm AIHA except that spherocytosis is less marked, red cells agglutinate in the cold (Fig. 5.9b) and the DAT reveals complement (C3d) only on the red cell surface.

Treatment consists of keeping the patient warm and treating the underlying cause, if present. Alkylating agents such as chlorambucil may be helpful in the chronic varieties. Both anti-CD20 (rituximab) and anti-CD52 (Campath-1H) have been used. Rituximab is particularly effective when there is an associated B-lymphoproliferative disease. Splenectomy does not usually help unless massive splenomegaly is present, and steroids are not helpful. Underlying lymphoma should be excluded in 'idiopathic' cases.

Paroxysmal cold haemoglobinuria is a rare syndrome of acute intravascular haemolysis after exposure to

the cold. It is caused by the Donath–Landsteiner antibody, an IgG antibody with specificity for the P blood group antigens, which binds to red cells in the cold but causes lysis with complement in warm conditions. Viral infections and syphilis are predisposing causes and the condition is usually self-limiting.

Alloimmune haemolytic anaemias

In these anaemias, antibody produced by one individual reacts with red cells of another. Two important situations are transfusion of ABO-incompatible blood and rhesus disease of the newborn which are considered in Chapters 27 and 28. The increased use of allogeneic transplantation for renal, hepatic, cardiac and bone marrow diseases has led to the recognition of alloimmune haemolytic anaemia resulting from the production of red cell antibodies in the recipient by donor lymphocytes transferred in the allograft.

Drug-induced immune haemolytic anaemias

Drugs may cause immune haemolytic anaemias via three mechanisms (Fig. 5.10):

- 1 Antibody directed against a drug–red cell membrane complex (e.g. penicillin, ampicillin);
- 2 Deposition of complement via a drug–protein (antigen)–antibody complex onto the red cell surface (e.g. quinidine, rifampicin); or
- 3 A true autoimmune haemolytic anaemia in which the role of the drug is unclear (e.g. methyl dopa).

In each case, the haemolytic anaemia gradually disappears when the drug is discontinued but with methyl dopa the autoantibody may persist for several months. The penicillin-induced immune haemolytic anaemias only occur with massive doses of the antibiotic.




Fig. 5.10 Three different mechanisms of drug-induced immune haemolytic anaemia. In each case the coated (opsonized) cells are destroyed in the reticuloendothelial system.

Red cell fragmentation syndromes

These arise through physical damage to red cells either on abnormal surfaces (e.g. artificial heart valves or arterial grafts), arteriovenous malformations or as a microangiopathic haemolytic anaemia. This is caused by red cells passing through abnormal small vessels. The latter may be caused by deposition of fibrin strands often associated with disseminated intravascular coagulation (DIC) or platelet adherence as in thrombotic thrombocytopenic purpura (TTP) (p. 284) or vasculitis (e.g. polyarteritis nodosa; Table 5.6). The peripheral blood contains many deeply staining red cell fragments (Fig. 5.11). Clotting abnormalities typical of DIC (p. 298) with a low platelet count are also present when DIC underlies the haemolysis. TTP is discussed in detail on page 284.

March haemoglobinuria

This is caused by damage to red cells between the small bones of the feet, usually during prolonged marching or running. The blood film does not show fragments.

Infections

Infections can cause haemolysis in a variety of ways. They may precipitate an acute haemolytic crisis

Table 5.6 Red cell fragmentation syndromes.

Cardiac haemolysis	Prosthetic heart valves Patches, grafts Perivalvular leaks
Arteriovenous malformations	
Microangiopathic	TTP-HUS Disseminated intravascular coagulation Malignant disease Vasculitis (e.g. polyarteritis nodosa) Malignant hypertension Pre-eclampsia/HELLP Renal vascular disorders/HELLP syndrome Ciclosporin Homograft rejection

HELLP, haemolysis with elevated liver function tests and low platelets; HUS, haemolytic uraemic syndrome; TTP, thrombotic thrombocytopenic purpura.

in G6PD deficiency or cause microangiopathic haemolytic anaemia (e.g. with meningococcal or pneumococcal septicaemia). Malaria causes haemolysis by extravascular destruction of parasitized red cells as well as by direct intravascular lysis. Blackwater fever is an acute intravascular haemolysis accompanied by acute renal failure caused by *Falciparum* malaria. *Clostridium perfringens* septicaemia

Fig. 5.11 Blood film in microangiopathic haemolytic anaemia (in this patient Gram-negative septicaemia). Numerous contracted and deeply staining cells and cell fragments are present.



can cause intravascular haemolysis with marked microspherocytosis.

Chemical and physical agents

Certain drugs (e.g. dapsone and Salazopyrin) in high doses cause oxidative intravascular haemolysis with Heinz body formation in normal subjects. In Wilson's disease an acute haemolytic anaemia can occur as a result of high levels of copper in the blood. Chemical poisoning (e.g. with lead, chlorate or arsine) can cause severe haemolysis. Severe burns damage red cells causing acanthocytosis or spherocytosis.

Secondary haemolytic anaemias

In many systemic disorders red cell survival is shortened. This may contribute to anaemia (Chapter 26).

Paroxysmal nocturnal haemoglobinuria

PNH is a rare, acquired, clonal disorder of marrow stem cells in which there is deficient synthesis of the glycosylphosphatidylinositol (GPI) anchor, a structure that attaches several surface proteins to the cell membrane. It results from mutations in the X chromosome gene coding for phosphatidylinositol glycan protein A (PIG-A) which is essential for the formation of the GPI anchor. The net result is that GPI-linked proteins (such as CD55 and CD59) are absent from the cell surface of all the cells derived from the abnormal stem cell (Fig. 5.12). The lack of surface molecules decay-activating factor (DAF, CD55) and membrane inhibitor of reactive lysis (MIRL, CD59) render red cells sensitive to lysis by complement and the result is chronic intravascular haemolysis. Haemosiderinuria is a constant feature and can give rise to iron deficiency which may exacerbate the anaemia. CD55 and CD59 are also present on white cells and platelets. The other main clinical problem seen in PNH is thrombosis and patients may develop recurrent thromboses of large veins including portal and hepatic veins, as well as intermittent abdominal pain brought about by thrombosis of mesenteric veins.

PNH is almost invariably associated with some form of bone marrow hypoplasia, often frank aplastic anaemia. It appears that the PNH clone may




Fig. 5.12 Schematic representation of the phosphatidylinositol glycan which anchors many different proteins to the cell membrane, e.g. CD59 (MIRL, membrane inhibitor of reactive lysis).

expand as a result of a selective pressure, possibly immunologically mediated, against cells that have normal GPI-linked membrane proteins.

PNH may be diagnosed by flow cytometry which shows loss of expression of the GPI-linked proteins, CD55 (DAF) and CD59 (MIRL). This has replaced the demonstration of red cell lysis in serum at low pH—the Ham's test.

Eculizumab, a humanized antibody against complement C5, inhibits the activation of terminal components of complement, reduces haemolysis and reduces transfusion requirements. Iron therapy is used for iron deficiency and long-term anticoagulation with warfarin may be needed. Immunosuppression can be useful and allogeneic stem cell transplantation is a definitive treatment. The disease occasionally remits but the median survival is approximately 10 years.

Bibliography

- Beutler E. (1996) Glucose-6-phosphate-dehydrogenase: population genetics and clinical manifestations. *Blood Rev* 10,45–52.
- Bolton-Maggs P.H.B. (2000) The diagnosis and management of hereditary spherocytosis. *Clin Haematol* 13,327–42.

- Dacie J.V. (1988, 1999) *The Haemolytic Anaemias*. Vol. 2, *The Hereditary Haemolytic Anaemias*; Vol. 3, *The Haemolytic Anaemias of Immune Origin*; Vol. 4, *Secondary and Symptomatic Haemolytic Anaemias*; Vol. 5, *Drug and Chemical-induced Haemolytic Anaemias*; PNH; *Haemolytic Disease of the Newborn*, 3rd edn. Churchill Livingstone, Edinburgh.
- Delaunay J. (2003) Red cell membrane and erythropoiesis genetic defects. *Hematol J* 4,225–32.
- Ekvall H. (2003) Malaria and anemia. *Curr Opin Hematol* 10,108–14.
- Gertz M.A. (2006) Cold agglutinin disease. *Haematologica* 91,439–45.
- Hillmen P., Hall C., Marsh J.C. *et al.* (2004) Effect of eculizumab on hemolysis and transfusion requirements in patients with paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 350,552–9.
- Luzzatto L. and Notaro R. (2001) Malaria: protecting against bad air. *Science* 293,442–3.
- Parker C., Omine M., Richards S. *et al.* (2005) Diagnosis and management of paroxysmal nocturnal hemoglobinuria. *Blood* 106,3699–709.
- Rees D.C., Duley J.A., Marinaki A.M. (2003) Pyrimidine 5' nucleotidase deficiency. *Br J Haematol* 120,375–83.
- Zanella A. (ed.) (2000) Inherited disorders of red cell metabolism. *Clin Haematol* 13,1–150.
- Zanella A., Fermo E., Bianchi P., Valentini G. (2005) Red cell pyruvate kinase deficiency: molecular and clinical aspects. *Br J Haematol* 130,11–25.

CHAPTER 6

6

Genetic disorders of haemoglobin

Haemoglobin synthesis, 72

Haemoglobin abnormalities, 74

Thalassaemias, 74

Sickle cell anaemia, 85

Prenatal diagnosis of genetic haemoglobin disorders, 90

Bibliography, 92

This chapter deals with inherited diseases caused by reduced or abnormal synthesis of globin. Mutations in the globin genes are the most prevalent monogenic disorders worldwide and affect approximately 7% of the world's population. The synthesis of normal haemoglobin in the fetus and adult is described first.

Haemoglobin synthesis

Normal adult blood contains three types of haemoglobin (see Table 2.2). The major component is haemoglobin A with the molecular structure $\alpha_2\beta_2$. The minor haemoglobins contain γ (fetal Hb or Hb F) or δ (Hb A₂) globin chains instead of β chains. In the embryo and fetus, Gower 1, Portland, Gower 2 and fetal Hb dominate at different stages (Fig. 6.1). The genes for the globin chains occur in two clusters: ϵ , γ , δ and β on chromosome 11 and ζ and α on chromosome 16. Two types of γ chain occur, G γ and A γ , which differ by a glycine or alanine amino acid at position 136 in the polypeptide chain. The α -chain gene is duplicated and both α_1 genes (α_1 and α_2) on each chromosome are active (Fig. 6.1).

Molecular aspects

All the globin genes have three exons (coding regions) and two introns (non-coding regions whose DNA is not represented in the finished protein). The initial RNA is transcribed from both introns and exons,

and from this transcript the RNA derived from introns is removed by a process known as splicing (Fig. 6.2). The introns always begin with a G-T dinucleotide and end with an A-G dinucleotide. The splicing machinery recognizes these sequences as well as neighbouring conserved sequences. The RNA in the nucleus is also 'capped' by addition of a structure at the 5' end which contains a seven methyl-guanosine group. The cap structure may be important for attachment of the mRNA to ribosomes. The newly formed mRNA is also polyadenylated at the 3' end (Fig. 6.2). This stabilizes it. Thalassaemia may arise from mutations or deletions of any of these sequences.

A number of other conserved sequences are important in globin synthesis and mutations at these sites may also give rise to thalassaemia. These sequences influence gene transcription, ensure its fidelity, specify sites for the initiation and termination of translation, and ensure the stability of newly synthesized mRNA. Promoters are found 5' of the gene, either close to the initiation site or more distally. They are the sites where RNA polymerases bind and catalyse gene transcription (see Fig. 1.9). Enhancers occur either 5' or 3' to the gene (Fig. 6.2). Enhancers are important in the tissue-specific regulation of globin gene expression and in regulation of the synthesis of the various globin chains during fetal and postnatal life. The locus control region (LCR) is a genetic regulatory element, situated a long way upstream of the β -globin cluster,




Fig. 6.1 (a) The globin gene clusters on chromosomes 16 and 11. In embryonic, fetal and adult life different genes are activated or suppressed. The different globin chains are synthesized independently and then combine with each other to produce the different haemoglobins.

The γ gene may have two sequences, which code for either a glutamic acid or alanine residue at position 136 (G_γ or A_γ , respectively). LCR, locus control region, HS-40, see text below. (b) Synthesis of individual globin chains in prenatal and postnatal life.

that controls the genetic activity of each domain, probably by physically interacting with the promoter region and opening up the chromatin to allow transcription factors to bind. The α -globin gene cluster also contains an LCR-like region termed HS40. GATA-1, FoG and NF-E2 transcription factors, expressed mainly in erythroid precursors, are important in determining the expression of globin genes in erythroid cells.

Globin mRNA enters the cytoplasm and attaches to ribosomes (translation) where the synthesis of globin chains takes place. This occurs by attachment

of transfer RNAs, each with its individual amino acid, by codon–anticodon base pairing to an appropriate position on the mRNA template.

Switch from fetal to adult haemoglobin

The globin genes are arranged on chromosomes 11 and 16 in the order in which they are expressed (Fig. 6.1). Certain embryonic haemoglobins are usually only expressed in yolk sac erythroblasts. The β -globin gene is expressed at a low level in early fetal life, but the main switch to adult haemoglobin




Fig. 6.2 The expression of a human globin gene from transcription, excision of introns, splicing of exons and translation to ribosomes. The primary transcript is 'capped' at the 5' end and a poly A tail is then added.

occurs 3–6 months after birth when synthesis of the γ chain is largely replaced by the β chain. How this switch comes about is largely unknown. However, it is clear that the methylation state of the gene (expressed genes tend to be hypomethylated, non-expressed hypermethylated), the state of the chromosome packaging and various enhancer sequences all play a part in determining whether a particular gene will be transcribed.

Haemoglobin abnormalities

These result from the following:

- 1 Synthesis of an abnormal haemoglobin.
- 2 Reduced rate of synthesis of normal α - or β -globin chains (the α - and β -thalassaemias).

Table 6.1 shows some of the first group of syndromes that arise from synthesis of an α or β chain with an amino acid substitution. In many cases, however, the abnormality is completely silent. The clinically most important abnormality is sickle cell anaemia. Haemoglobin (Hb) C, D and E are also common and, like Hb S, are substitutions in the β chain. Unstable haemoglobins are rare and cause a

Table 6.1 The clinical syndromes produced by haemoglobin abnormalities.

Syndrome	Abnormality
Haemolysis	Crystalline haemoglobins (Hb S, C, D, E, etc.)
Thalassaemia	Unstable haemoglobin α or β resulting from reduced globin chain synthesis
Familial polycythaemia	Altered oxygen affinity
Methaemoglobinaemia	Failure of reduction (Hb Ms)

chronic haemolytic anaemia of varying severity with intravascular haemolysis (see Table 5.2). Abnormal haemoglobins may also cause (familial) polycythaemia (Chapter 19) or congenital methaemoglobinaemia (Chapter 2).

The genetic defects of haemoglobin are the most common genetic disorders worldwide. They occur in tropical and subtropical areas (Fig. 6.3) and most appear to have been selected because the carrier state affords some protection against malaria. β -Thalassaemia is more common in the Mediterranean region while α -thalassaemia is more common in the Far East.

Thalassaemias

These are a heterogeneous group of genetic disorders that result from a reduced rate of synthesis of α or β chains.

α -Thalassaemia syndromes

These are usually caused by gene deletions and are listed in Table 6.2. As there are normally four copies of the α -globin gene, the clinical severity can be classified according to the number of genes that are missing or inactive. Loss of all four genes completely suppresses α -chain synthesis (Fig. 6.4) and because the α chain is essential in fetal as well as in adult haemoglobin this is incompatible with life and leads to death *in utero* (hydrops fetalis; Fig. 6.5). Three α gene deletions leads to a moderately severe (haemoglobin 7–11 g/dL) microcytic, hypochromic anaemia (Fig. 6.6) with splenomegaly. This is




Fig. 6.3 The geographical distribution of the thalassaemias and the more common, inherited, structural haemoglobin abnormalities.

Table 6.2 Classification of thalassaemia.

Clinical	
<i>Hydrops fetalis</i>	<i>Thalassaemia minor</i>
Four gene deletion α -thalassaemia	β^0 -Thalassaemia trait
<i>Thalassaemia major</i>	β^+ -Thalassaemia trait
Transfusion dependent, homozygous	Hereditary persistence of fetal haemoglobin
β^0 -thalassaemia or other combinations of β -thalassaemia trait	$\delta\beta$ -Thalassaemia trait
<i>Thalassaemia intermedia</i>	α^0 -Thalassaemia trait
See Table 6.5	α^+ -Thalassaemia trait

Genetic

Type	Haplotype	Heterozygous thalassaemia trait (minor)*	Homozygous
α -Thalassaemias [†]			
α^0	--/	MCV, MCH low	Hydrops fetalis
α^+	- α /	MCV, MCH minimally reduced	As heterozygous α^0 -thalassaemia
β -Thalassaemias			
β^0		MCV, MCH low (Hb A ₂ >3.5%)	Thalassaemia major (Hb F 98%, Hb A ₂ 2%)
β^+		MCV, MCH low (Hb A ₂ >3.5%)	Thalassaemia major or intermedia (Hb F 70–80%, Hb A 10–20%, Hb A ₂ variable)
$\delta\beta$ -Thalassaemia and hereditary persistence of fetal haemoglobin		MCV, MCH low (Hb F 5–20%, Hb A ₂ normal)	Thalassaemia intermedia (Hb F 100%)
Hb Lepore		MCV, MCH low (Hb A 80–90%, Hb Lepore 10%, Hb A ₂ reduced)	Thalassaemia major or intermedia (Hb F 80%, Hb Lepore 10–20%, Hb A, Hb A ₂ absent)

MCH, mean corpuscular haemoglobin; MCV, mean corpuscular volume.

* Occasionally heterozygous β -thalassaemia is dominant (associated with the clinical picture of thalassaemia intermedia). There are several explanations.

[†] Compound heterozygote $\alpha^0\alpha^+$ (--/- α) is haemoglobin H disease.




Fig. 6.4 The genetics of α -thalassaemia. Each α gene may be deleted or (less frequently) dysfunctional. The orange boxes represent normal genes, and the blue boxes represent gene deletions or dysfunctional genes.






Fig. 6.5 α -Thalassaemia: hydrops fetalis, the result of deletion of all four α -globin genes (homozygous α^0 -thalassaemia). The main haemoglobin present is Hb Barts (γ_4). The condition is incompatible with life beyond the fetal stage. (Courtesy of Professor D. Todd)



(a)


Fig. 6.6 (a) α -Thalassaemia: haemoglobin H disease (three α -globin gene deletion). The blood film shows marked hypochromic, microcytic cells with target cells and poikilocytosis. (b) α -Thalassaemia: haemoglobin H disease. Supravital staining with brilliant cresyl blue



(b)

reveals multiple fine, deeply stained deposits (‘golf ball’ cells) caused by precipitation of aggregates of β -globin chains. Hb H can also be detected as a fast-moving band on haemoglobin electrophoresis (see Fig. 6.14).

Fig. 6.7 Distribution of different mutations of β -thalassaemia major in the Mediterranean area. IVS1, IVS2 intervening sequences; 1, 6, 39, 110, 745 are mutations of corresponding codons. (Courtesy of Professor A. Cao)



known as Hb H disease because haemoglobin H (β_4) can be detected in red cells of these patients by electrophoresis or in reticulocyte preparations (Fig. 6.6). In fetal life, Hb Barts (γ_4) occurs.

The α -thalassaemia traits are caused by loss of one or two genes and are usually not associated with anaemia, although the mean corpuscular volume (MCV) and mean corpuscular haemoglobin (MCH) are low and the red cell count is over $5.5 \times 10^{12}/L$. Haemoglobin electrophoresis is normal and α/β -chain synthesis studies or DNA analyses are needed to be certain of the diagnosis. The normal α/β -synthesis ratio is 1 : 1 and this is reduced in the α -thalassaemias and raised in β -thalassaemias. Uncommon non-deletional forms of α -thalassaemia are caused by point mutations producing dysfunction of the genes or rarely by mutations affecting termination of translation which give rise to an elongated but unstable chain (e.g. Hb Constant Spring). Two rare forms of α -thalassaemia are associated with mental retardation. They are caused by mutation in a gene on chromosome 16 (ATR-16) or on chromosome X (ATR-X) which control the transcription of the α globin and other genes.

β -Thalassaemia syndromes

β -Thalassaemia major

This condition occurs on average in one in four offspring if both parents are carriers of the β -thalassaemia trait. Either no β chain (β^0) or small amounts (β^+) are synthesized. Excess α chains

precipitate in erythroblasts and in mature red cells causing the severe ineffective erythropoiesis and haemolysis that are typical of this disease. The greater the α -chain excess, the more severe the anaemia. Production of γ chains helps to 'mop up' excess α chains and to ameliorate the condition. Over 200 different genetic defects have now been detected (Figs 6.7 and 6.8).

Unlike α -thalassaemia, the majority of genetic lesions are point mutations rather than gene deletions. These mutations may be within the gene complex itself or in promoter or enhancer regions. Certain mutations are particularly frequent in some communities (Fig. 6.7) and this may simplify antenatal diagnosis aimed at detecting the mutations in fetal DNA. Thalassaemia major is often a result of inheritance of two different mutations, each affecting β -globin synthesis (compound heterozygotes). In some cases, deletion of the β gene, δ and β genes or even δ , β and γ genes occurs. In others, unequal crossing-over has produced $\delta\beta$ fusion genes (so called Lepore syndrome named after the first family in which this was diagnosed) (p. 84).

Clinical features

- 1 Severe anaemia becomes apparent at 3–6 months after birth when the switch from γ - to β -chain production should take place.
- 2 Enlargement of the liver and spleen occurs as a result of excessive red cell destruction, extramedullary haemopoiesis and later because of iron overload. The large spleen increases blood requirements by




Fig. 6.8 Examples of mutations that produce β -thalassaemia. These include single base changes, small deletions and insertions of one or two bases affecting introns, exons or the flanking regions of the β -globin gene. FS, 'frameshifts': deletion of nucleotide(s) that places the reading frame out of phase downstream of the lesion; NS, 'non-sense': premature chain termination as a result of a

new translational stop codon (e.g. UAA); SPL, 'splicing': inactivation of splicing or new splice sites generated (aberrant splicing) in exons or introns; promoter, CAP, initiation: reduction of transcription or translation as a result of lesion in promoter, CAP or initiation regions; Poly A, mutations on the poly A addition signal resulting in failure of poly A addition and an unstable mRNA.



Fig. 6.9 The facial appearance of a child with β -thalassaemia major. The skull is bossed with prominent frontal and parietal bones; the maxilla is enlarged.

increasing red cell destruction and pooling, and by causing expansion of the plasma volume.

3 Expansion of bones caused by intense marrow hyperplasia leads to a thalassaemic facies (Fig. 6.9) and to thinning of the cortex of many bones with a tendency to fractures and bossing of the skull with a 'hair-on-end' appearance on X-ray (Fig. 6.10).

4 The patient can be sustained by blood transfusions but iron overload caused by repeated transfusions is inevitable unless chelation therapy is given (Table 6.3). Each 500 mL of transfused blood contains approximately 250 mg iron. To make matters worse, iron absorption from food is *increased* in β -thalassaemia, probably secondary

Table 6.3 Causes of refractory anaemia that may lead to transfusional iron overload.

Congenital	Acquired
β -Thalassaemia major	Myelodysplasia
β -Thalassaemia/Hb E disease	Red cell aplasia
Sickle cell anaemia (some cases)	Aplastic anaemia
Red cell aplasia (Diamond–Blackfan)	Myelofibrosis
Sideroblastic anaemia	
Dyserythropoietic anaemia	





Fig. 6.10 The skull X-ray in β -thalassaemia major. There is a 'hair-on-end' appearance as a result of expansion of the bone marrow into cortical bone.

to ineffective erythropoiesis and inappropriately low serum hepcidin levels. Iron damages the liver (Fig. 6.11) and the endocrine organs with failure of growth, delayed or absent puberty, diabetes mellitus, hypothyroidism and hypoparathyroidism.


Skin pigmentation as a result of excess melanin and haemosiderin gives a slaty grey appearance at an early stage of iron overload.

Most importantly, iron damages the heart. In the absence of intensive iron chelation, death occurs in the second or third decade, usually from congestive heart failure or cardiac arrhythmias. T_2^* magnetic resonance imaging (MRI) is a valuable measure of cardiac (or liver) iron (Fig. 6.12). It can detect increased cardiac iron before sensitive tests detect impaired cardiac function. Serum ferritin and liver iron show poor correlation with cardiac iron estimated by T_2^* MRI (Fig. 6.12).

5 Infections can occur for a variety of reasons. In infancy, without adequate transfusion, the anaemic



(a)



(b)

Fig. 6.11 β -Thalassaemia major: needle biopsy of liver. (a) Grade IV siderosis with iron deposition in the hepatic parenchymal cells, bile duct epithelium, macrophages and fibroblasts (Perls' stain). (b) Reduction of iron excess in liver after intensive chelation therapy.




Fig. 6.12 T_2^* magnetic resonance images (MRIs) showing tissue appearance in iron overload (a) normal volunteer, (b) severe iron overload. Green arrow, normal appearance; red arrow, iron overload. Lack of correlation: liver and cardiac iron in two cases of thalassaemia major (c) and (d).

child is prone to bacterial infections. Pneumococcal, *Haemophilus* and meningococcal infections are likely if splenectomy has been carried out and prophylactic penicillin is not taken. *Yersinia enterocolitica* occurs, particularly in iron-loaded patients being treated with deferoxamine; it may cause severe gastroenteritis. Transfusion of viruses by blood transfusion may occur. Liver disease in thalassaemia is most frequently a result of hepatitis C but hepatitis B is also common where the virus is endemic. Human immunodeficiency virus (HIV) has been transmitted to some patients by blood transfusion.

6 Osteoporosis may occur in well-transfused pati-

ents. It is more common in diabetic patients with endocrine abnormalities and with marrow expansion resulting from ineffective erythropoiesis.

Laboratory diagnosis

1 There is a severe hypochromic, microcytic anaemia, raised reticulocyte percentage with normoblasts, target cells and basophilic stippling in the blood film (Fig. 6.13).

2 Haemoglobin electrophoresis reveals absence or almost complete absence of Hb A, with almost all the circulating haemoglobin being Hb F. The Hb A₂ percentage is normal, low or slightly raised

Fig. 6.13 Blood film in β -thalassaemia major post-splenectomy. There are hypochromic cells, target cells and many nucleated red cells (normoblasts). Howell-Jolly bodies are seen in some red cells.



(Fig. 6.14a). High performance liquid chromatography is now usually used as first-line method to diagnose haemoglobin disorders (Fig. 6.14b). α/β -Globin chain synthesis studies on reticulocytes show an increased $\alpha : \beta$ ratio with reduced or absent β -chain synthesis. DNA analysis is used to identify the defect on each allele.

Assessment of iron status

The tests that can be performed to assess iron overload are listed in Table 6.4. Tests may also be carried out to determine the degree of organ damage caused by iron. The serum ferritin is the most widely used test. It is usual in thalassaemia major to attempt to keep the level between 1000 and 1500 $\mu\text{g/L}$, when

Table 6.4 Assessment of iron overload.

Assessment of iron stores

Serum ferritin

Serum iron and percentage saturation of transferrin (iron-binding capacity)

Serum non-transferrin bound iron

Bone marrow biopsy (Perls' stain) for reticuloendothelial stores

DNA test for mutation resulting in Cys282Tyr in the *HFE* gene

Liver biopsy (parenchymal and reticuloendothelial stores)

Liver CT scan or MRI

Cardiac MRI (T_2^* technique)

Deferoxamine or deferiprone iron excretion test (chelatable iron)

Repeated phlebotomy until iron deficiency occurs

Assessment of tissue damage caused by iron overload

Cardiac

Clinical; chest X-ray; ECG; 24-h monitor; echocardiography; radionuclide (MUGA) scan to check left ventricular ejection fraction at rest and with stress


Liver

Liver function tests; liver biopsy; CT scan or MRI


Endocrine

Clinical examination (growth and sexual development); glucose tolerance test; pituitary gonadotrophin release tests; thyroid, parathyroid, gonadal, adrenal function, growth hormone assays; radiology for bone age; isotopic bone density study

CT, computed tomography; ECG, electrocardiography; MRI, magnetic resonance imaging; MUGA, multiple gated acquisition.



(a)



(b)

Fig. 6.14 (a) Haemoglobin electrophoretic patterns in normal adult human blood and in subjects with sickle cell (Hb S) trait or disease, β -thalassaemia trait, β -thalassaemia major, Hb S/ β -thalassaemia or Hb S/Hb C disease and Hb H disease. (b) High performance liquid chromatography. The different haemoglobins elute at different times from the column and their concentrations are read automatically. In this example, the patient is a carrier of sickle cell disease.

the body iron stores are approximately 5–10 times normal. However, the serum ferritin correlates poorly with cardiac iron and is raised in relation to iron status in viral hepatitis and other inflammatory disorders and should therefore be interpreted in conjunction with other tests such as T_2^* MRI assessment of cardiac iron, liver biopsy iron (Fig. 6.11) and urine excretion of iron in response to deferoxamine or deferiprone. The function of the heart, liver and endocrine organs are also needed to determine the efficacy of chelation therapy (Table 6.4).

Treatment

1 Regular blood transfusions are needed to maintain the haemoglobin over 10 g/dL at all times. This usually requires 2–3 units every 4–6 weeks. Fresh blood, filtered to remove white cells, gives the best red cell survival with the fewest reactions. The patients should be genotyped at the start of the transfusion programme in case red cell antibodies against transfused red cells develop.

2 Regular folic acid (e.g. 5 mg/day) is given if the diet is poor.




Fig. 6.15 (a) Reduction in cardiac iron assessed by T_2^* MRI is greater in patients treated with deferiprone than with deferoxamine. (b) Improvement in left ventricular (LV) ejection factor is greater with deferiprone than with deferoxamine. (From Pennell *et al.* (2006) with permission.)

3 Iron chelation therapy is used to treat iron overload. The most established drug, deferoxamine, is inactive orally. It may be given by a separate infusion bag 1–2 g with each unit of blood transfused and by subcutaneous infusion 40 mg/kg over 8–12 h, 5–7 days weekly. It is commenced in infants after 10–15 units of blood have been transfused.

Iron chelated by deferoxamine is mainly excreted in the urine but up to one-third is also excreted in the stools. If patients comply with this intensive iron chelation regime, life expectancy for patients with thalassaemia major and other chronic refractory anaemias receiving regular blood transfusion (Table 6.3) improves considerably. In some cases, intensive continuous chelation therapy with intravenous deferoxamine can reverse heart damage caused by iron overload. However, lack of compliance is frequent and the drug is costly. In addition, deferoxamine may have side-effects, especially in children with relatively low serum ferritin levels. These include high tone deafness, retinal damage, bone abnormalities and growth retardation. Patients should have auditory and fundoscopic examinations at regular intervals.

Deferiprone is an orally active iron chelator which causes predominantly urine iron excretion. It is usually given in three doses daily. It may be used alone or in combination with deferoxamine. The drugs have an additive effect on iron excretion. Deferiprone is more effective than deferoxamine at removing cardiac iron (Fig. 6.15a). Compliance is also better. Side-effects include an arthropathy, agranulocytosis (in about 1%), neutropenia, gastrointestinal disturbance and zinc deficiency.

Deferasirox (ICL670, Exjade) is the newest oral chelator. It is given once daily and causes faecal iron excretion only. Skin rashes and transient changes in liver enzymes have been reported. The ease of administration and its lack of major side-effects are likely to result in its widespread use.

4 Vitamin C, 200 mg/day, increases excretion of iron produced by deferoxamine.

5 Splenectomy may be needed to reduce blood requirements. This should be delayed until the patient is over 6 years old because of the high risk of dangerous infections post-splenectomy. The vaccinations and antibiotics to be given are described in Chapter 9.

6 Endocrine therapy is given either as replacement because of end-organ failure or to stimulate the pituitary if puberty is delayed. Diabetics will require insulin therapy. Patients with osteoporosis may need additional therapy with increased calcium and vitamin D in their diet, together with a bisphosphonate and appropriate endocrine therapy.

7 Immunization against hepatitis B should be carried out in all non-immune patients. Treatment for transfusion-transmitted hepatitis C with α -interferon and ribavirin is needed if viral genomes are detected in plasma.

8 Allogeneic bone marrow transplantation offers the prospect of permanent cure. The success rate (long-term thalassaemia major-free survival) is over 80% in well-chelated younger patients without liver fibrosis or hepatomegaly. A human leucocyte antigen (HLA) matching sibling (or rarely other family member or matching unrelated donor) acts as donor. Failure is mainly a result of recurrence of thalassaemia, death (e.g. from infection) or severe chronic graft-versus-host disease.

β -Thalassaemia trait (minor)

This is a common, usually symptomless, abnormality characterized like α -thalassaemia trait by a hypochromic, microcytic blood picture (MCV and MCH very low) but high red cell count ($>5.5 \times 10^{12}/L$) and mild anaemia (haemoglobin 10–12 g/dL). It is usually more severe than α trait. A raised Hb A₂ ($>3.5\%$) confirms the diagnosis. One of the most important indications for making the diagnosis is that it allows the possibility of prenatal counselling to patients with a partner who also has a significant haemoglobin disorder. If both carry β -thalassaemia trait there is a 25% risk of a thalassaemia major child.

Thalassaemia intermedia

Cases of thalassaemia of moderate severity (haemoglobin 7.0–10.0 g/dL) who do not need regular transfusions are called thalassaemia intermedia (Table 6.5). This is a *clinical* syndrome which may be caused by a variety of genetic defects: homozygous β -thalassaemia with production of more Hb F than usual or with mild defects in β -chain synthesis, by β -thalassaemia trait alone of unusual severity ('dominant' β -thalassaemia) or β -thalassaemia trait in association with mild globin abnormalities such as Hb Lepore. The coexistence of α -thalassaemia trait improves the haemoglobin level in homozygous β -thalassaemia by reducing the degree of chain imbalance and thus of α -chain precipitation and ineffective erythropoiesis. Conversely, patients

Table 6.5 Thalassaemia intermedia.

Homozygous β -thalassaemia

Homozygous mild β^+ -thalassaemia

Coinheritance of α -thalassaemia

Enhanced ability to make fetal haemoglobin (γ -chain production)

Heterozygous β -thalassaemia

Coinheritance of additional α -globin genes ($\alpha\alpha\alpha/\alpha\alpha$ or $\alpha\alpha\alpha/\alpha\alpha\alpha$)

Dominant β -thalassaemia trait

$\delta\beta$ -Thalassaemia and hereditary persistence of fetal haemoglobin

Homozygous $\delta\beta$ -thalassaemia

Heterozygous $\delta\beta$ -thalassaemia/ β -thalassaemia

Homozygous Hb Lepore (some cases)

Haemoglobin H disease

with β -thalassaemia trait who also have excess (five or six) α genes tend to be more anaemic than usual. The patient with thalassaemia intermedia may show bone deformity, enlarged liver and spleen, extramedullary erythropoiesis (Fig. 6.16) and features of iron overload caused by increased iron absorption. Hb H disease (three-gene deletion α -thalassaemia) is a type of thalassaemia intermedia without iron overload or extramedullary haemopoiesis.


$\delta\beta$ -Thalassaemia

This involves failure of production of both β and δ chains. Fetal haemoglobin production is increased to 5–20% in the heterozygous state which resembles thalassaemia minor haematologically. In the homozygous state only Hb F is present and haematologically the picture is of thalassaemia intermedia.

Haemoglobin Lepore

This is an abnormal haemoglobin caused by unequal crossing-over of the β and δ genes to produce a polypeptide chain consisting of the δ chain at its amino end and β chain at its carboxyl end. The $\delta\beta$ -fusion chain is synthesized inefficiently and normal δ - and β -chain production is abolished. The homozygotes show thalassaemia intermedia and the heterozygotes thalassaemia trait.

Fig. 6.16 β -Thalassaemia intermedia: MRI scan showing masses of extramedullary haemopoietic tissue arising from the ribs and in the paravertebral region without encroachment of the spinal cord.



Hereditary persistence of fetal haemoglobin

These are a heterogeneous group of genetic conditions caused by deletions or cross-overs affecting the production of β and γ chains or, in non-deletion forms, by point mutations upstream from the γ -globin genes.

Association of β -thalassaemia trait with other genetic disorders of haemoglobin

The combination of β -thalassaemia trait with Hb E trait usually causes a transfusion-dependent thalassaemia major syndrome, but some cases are intermediate. β -Thalassaemia trait with Hb S trait produces the clinical picture of sickle cell anaemia rather than of thalassaemia (p. 90). β -Thalassaemia trait with Hb D trait causes a hypochromic, microcytic anaemia of varying severity.


Sickle cell anaemia

Sickle cell disease is a group of haemoglobin disorders in which the sickle β -globin gene is inherited. Homozygous sickle cell anaemia (Hb SS) is the most common while the doubly heterozygote conditions of Hb SC and Hb S β thal also cause


sickling disease. Hb S (Hb $\alpha_2\beta_2^S$) is insoluble and forms crystals when exposed to low oxygen tension. Deoxygenated sickle haemoglobin polymerizes into long fibres, each consisting of seven intertwined double strands with cross-linking. The red cells sickle and may block different areas of the microcirculation or large vessels causing infarcts of various organs. The sickle β -globin abnormality is caused by substitution of valine for glutamic acid in position 6 in the β chain (Fig. 6.17). It is very widespread and is found in up to one in four West Africans, maintained at this level because of the protection against malaria that is afforded by the carrier state.

Normal β - chain	Amino acid	pro	glu	glu
	Base composition	CCT	GAG	GAG
Sickle β - chain	Base composition	CCT	GUG	GAG
	Amino acid	pro	val	glu

Fig. 6.17 Molecular pathology of sickle cell anaemia. There is a single base change in the DNA coding for the amino acid in the sixth position in the β -globin chain (adenine is replaced by thymine). This leads to an amino acid change from glutamic acid to valine. A, adenine; C, cytosine; G, guanine; glu, glutamic acid; pro, proline; T, thymine; val, valine.



(a)



(b)

Fig. 6.18 Sickle cell anaemia. (a) Radiograph of the pelvis of a young man of West Indian origin which shows avascular necrosis with flattening of the femoral heads, more marked on the right, coarsening of the bone architecture and cystic areas in the right femoral neck caused by previous infarcts. (b) MRI scan of the hips of a 17-year-old female, showing a small area of high signal in

the anterior portion of the right hip (arrowed) with a low intensity rim. This is typical of early avascular necrosis. The irregular outline and signal in the left hip results from more advanced avascular necrosis. Joint fluid is shown as a high signal (white rim) surrounding the femoral head. (Courtesy of Dr L. Berger)




Fig. 6.19 Sick cell anaemia: (a) painful swollen fingers (dactylitis) in a child and (b) the hand of an 18-year-old Nigerian boy with the 'hand-foot' syndrome. There is marked shortening of the right middle finger because of dactylitis in childhood affecting the growth of the epiphysis.

Homozygous disease

Clinical features

Clinical features are of a severe haemolytic anaemia punctuated by crises. The symptoms of anaemia are often mild in relation to the severity of the anaemia because Hb S gives up oxygen (O_2) to tissues relatively easily compared with Hb A, its O_2 dissociation curve being shifted to the right (see Fig. 2.9). The clinical expression of Hb SS is very variable, some patients having an almost normal life, free of crises but others develop severe crises even as infants and may die in early childhood or as young adults. Crises may be vaso-occlusive, visceral, aplastic or haemolytic.

Painful vaso-occlusive crises

These are the most frequent and are precipitated by such factors as infection, acidosis, dehydration or deoxygenation (e.g. altitude, operations, obstetric delivery, stasis of the circulation, exposure to cold, violent exercise). Infarcts can occur in a variety of organs including the bones (hips, shoulders and vertebrae are commonly affected) (Fig. 6.18), the lungs and the spleen. The most serious vaso-occlusive crisis is of the brain (a stroke occurs in 7% of all patients) or spinal cord. Transcranial

Doppler ultrasonography detects abnormal blood flow indicative of arterial stenosis. This predicts for strokes in children. This can be largely prevented by regular blood transfusions in these cases. The 'hand-foot' syndrome (painful dactylitis caused by infarcts of the small bones) is frequently the first presentation of the disease and may lead to digits of varying lengths (Fig. 6.19).

Visceral sequestration crises

These are caused by sickling within organs and pooling of blood, often with a severe exacerbation of anaemia. The acute sickle chest syndrome is a feared complication and the most common cause of death after puberty. It presents with dyspnoea, falling arterial PO_2 , chest pain and pulmonary infiltrates on chest X-ray. Treatment is with analgesia, oxygen, exchange transfusion and ventilatory support if necessary. Hepatic and girdle sequestration crises and splenic sequestration may lead to severe illness requiring exchange transfusions. Splenic sequestration is typically seen in infants and presents with an enlarging spleen, falling haemoglobin and abdominal pain. Treatment is with transfusion and patients must be monitored at regular intervals as progression may be rapid. Attacks tend to be recurrent and splenectomy is often needed.



Fig. 6.20 Sickle cell anaemia: medial aspect of the ankle of a 15-year-old Nigerian boy showing necrosis and ulceration.

Aplastic crises

These occur as a result of infection with parvovirus or from folate deficiency and are characterized by a sudden fall in haemoglobin, usually requiring transfusion. They are characterized by a fall in reticulocytes as well as haemoglobin (see Fig. 20.4).

Haemolytic crises

These are characterized by an increased rate of haemolysis with a fall in haemoglobin but rise in reticulocytes and usually accompany a painful crisis.

Other clinical features

Ulcers of the lower legs are common, as a result of vascular stasis and local ischaemia (Fig. 6.20). The spleen is enlarged in infancy and early childhood but later is often reduced in size as a result of infarcts (autosplenectomy). Pulmonary hypertension detected by Doppler echocardiography and an




Fig. 6.21 *Salmonella* osteomyelitis: lateral radiograph of the lower femur and knee. The periosteum is irregularly raised in the lower third of the femur.

increased tricuspid regurgitant velocity are common and increases the risk of death. A proliferative retinopathy and priapism are other clinical complications. Chronic damage to the liver may occur through microinfarcts. Pigment (bilirubin) gallstones are frequent. The kidneys are vulnerable to infarctions of the medulla with papillary necrosis. Failure to concentrate urine aggravates the tendency to dehydration and crisis, and nocturnal enuresis is common. Osteomyelitis may also occur, usually from *Salmonella* spp. (Fig. 6.21).

Laboratory findings

- 1 The haemoglobin is usually 6–9 g/dL—low in comparison to symptoms of anaemia.
- 2 Sick cells and target cells occur in the blood (Fig. 6.22). Features of splenic atrophy (e.g. Howell-Jolly bodies) may also be present.
- 3 Screening tests for sickling are positive when the

Fig. 6.22 (a) Sick cell anaemia: peripheral blood film showing deeply staining sickle cells, target cells and polychromasia. (b) Homozygous Hb C disease: peripheral blood film showing many target cells, deeply staining rhomboidal and spherocytic cells.



blood is deoxygenated (e.g. with dithionate and Na_2HPO_4).

4 Haemoglobin electrophoresis (Fig. 6.14): in Hb SS, no Hb A is detected. The amount of Hb F is variable and is usually 5–15%, larger amounts are normally associated with a milder disorder.

Treatment

1 Prophylactic—avoid those factors known to precipitate crises, especially dehydration, anoxia, infections, stasis of the circulation and cooling of the skin surface.

2 Folic acid (e.g. 5 mg once weekly).

3 Good general nutrition and hygiene.

4 Pneumococcal, *Haemophilus* and meningococcal vaccination and regular oral penicillin are effective at reducing the infection rate with these organisms and should be strongly encouraged. Oral penicillin should start at diagnosis and continue at least until puberty. Hepatitis B vaccination is also given as transfusions may be needed.

5 Crises—treat by rest, warmth, rehydration by oral fluids and/or intravenous normal saline (3 L in 24 h) and antibiotics if infection is present. Analgesia at the appropriate level should be given. Suitable drugs are paracetamol, a non-steroidal anti-inflammatory agent and opiates (e.g. continuous subcutaneous diamorphine). Blood transfusion is given only if

there is very severe anaemia with symptoms. Exchange transfusion may be needed particularly if there is neurological damage, a visceral sequestration crisis or repeated painful crises. This is aimed at achieving an Hb S percentage of less than 30 in severe cases and after a stroke is continued for at least 2 years.

6 Particular care is needed in pregnancy and anaesthesia. There is debate as to whether or not patients need transfusions with normal blood to reduce Hb S levels during pregnancy or before delivery or for minor operations. Careful anaesthetic and recovery techniques must be used to avoid hypoxaemia or acidosis. Routine transfusions throughout pregnancy are given to those with a poor obstetric history or a history of frequent crises.

7 Transfusions—these are also sometimes given repeatedly as prophylaxis to patients having frequent crises or who have had major organ damage (e.g. of the brain) or show abnormal transcranial Doppler studies. The aim is to suppress Hb S production over a period of several months or even years. Iron overload, which may need iron chelation therapy, and alloimmunization against donated blood are common problems.

8 Hydroxyurea (15.0–20.0 mg/kg) can increase Hb F levels and has been shown to improve the clinical course of children or adults who are having three or

more painful crises each year. It should not be used during pregnancy.

9 Stem cell transplantation can cure the disease and many patients have now been successfully treated. The mortality rate is less than 10%. Transplantation is only indicated in the severest of cases whose quality of life or life expectancy are substantially impaired.

10 Research into other drugs (e.g. butyrates) to enhance Hb F synthesis or to increase the solubility of Hb S is taking place. 'Gene therapy' is a distant prospect not yet available (Chapter 21).

Sickle cell trait

This is a benign condition with no anaemia and normal appearance of red cells on a blood film. Haematuria is the most common symptom and is thought to be caused by minor infarcts of the renal papillae. Hb S varies from 25 to 45% of the total haemoglobin (Fig. 6.14a). Care must be taken with anaesthesia, pregnancy and at high altitudes.

Combination of haemoglobin S with other genetic defects of haemoglobin

The most common of these are Hb S/ β -thalassaemia, and sickle cell/C disease. In Hb S/ β -thalassaemia, the MCV and MCH are lower than in homozygous Hb SS. The clinical picture is of sickle cell anaemia; splenomegaly is usual. Patients with Hb SC disease have a particular tendency to thrombosis and pulmonary embolism, especially in pregnancy. In general, when compared with Hb SS disease, they have a higher incidence of retinal abnormalities, milder anaemia, splenomegaly and generally a longer life expectancy. Diagnosis is made by haemoglobin electrophoresis, particularly with family studies.

Haemoglobin C disease

This genetic defect of haemoglobin is frequent in West Africa and is caused by substitution of lysine for glutamic acid in the β -globin chain at the same point as the substitution in Hb S. Hb C tends to form rhomboidal crystals and in the homozygous state there is a mild haemolytic anaemia with marked target cell formation, cells with rhomboidal shape

and microspherocytes (Fig. 6.22b). The spleen is enlarged. The carriers show a few target cells only.

Haemoglobin D disease

This is a group of variants all with the same electrophoretic mobility. Heterozygotes show no haematological abnormality while homozygotes have a mild haemolytic anaemia.

Haemoglobin E disease

This is the most common haemoglobin variant in South-East Asia. In the homozygous state, there is a mild microcytic, hypochromic anaemia. Haemoglobin E/ β^0 -thalassaemia, however, resembles homozygous β^0 -thalassaemia both clinically and haematologically.

Prenatal diagnosis of genetic haemoglobin disorders

It is important to give genetic counselling to couples at risk of having a child with a major haemoglobin defect. If a pregnant woman is found to have a haemoglobin abnormality, her partner should be




Fig. 6.23 Polymerase chain reaction. The primers hybridize to DNA on either side of the piece of DNA to be analysed. Repeated cycles of denaturation, association with the primers, incubation with a DNA polymerase and deoxyribonucleotides (dNTPs) results in amplification of the DNA over a million times within a few hours.

tested to determine whether he also carries a defect. When both partners show an abnormality and there is a risk of a serious defect in the offspring, particularly β -thalassaemia major, it is important to offer antenatal diagnosis. Several techniques are available, the choice depending on the stage of pregnancy and the potential nature of the defect.

DNA diagnosis

The majority of samples are obtained by chorionic villus biopsy although amniotic fluid cells are sometimes used. Techniques to sample maternal blood for fetal cells or fetal DNA are being developed. The DNA is then analysed using one of the following methods.

Polymerase chain reaction (PCR) is the most commonly used technique (Fig. 6.23) and may be performed by using primer pairs that only amplify




Fig. 6.24 The rapid prenatal diagnosis of β -thalassaemia by amplification refractory mutation system (ARMS). The father has the common Mediterranean codon 39 (CD39) mutation, the mother the IVS1-110 G \rightarrow A mutation. The fetus is heterozygous for the CD39 mutation. CVS, fetal DNA from chorionic villus sampling; F, father; M, mother. (Courtesy of Dr J. Old and Professor D.J. Weatherall)

individual alleles ('allele-specific priming') (Fig. 6.24) or by using consensus primers that amplify all the alleles followed by restriction digestion to detect a particular allele. This is best illustrated by Hb S in which the enzyme DdeI detects the A-T change (Fig. 6.25).




Fig. 6.25 Sickle cell anaemia: antenatal diagnosis by DdeI-PCR analysis. The DNA is amplified by two primers that span the sickle cell gene mutation site and produces a product of 473 base pairs (bp) in size. The product is digested with the restriction enzyme DdeI and the resulting fragments analysed by agarose gel electrophoresis. The replacement of an adenine base in the normal β -globin gene by thymine results in Hb S and removes a normal restriction site for DdeI, producing a larger 376 bp fragment than the normal 175 and 201 bp fragments in the digested amplified product. In this case, the CVS DNA shows both the normal fragments and the larger sickle cell product and so is AS. The gel shows DNA from the mother (M), father (F), fetal DNA from a chorionic villus sample (CVS), a normal DNA control (AA) and a homozygous sickle cell DNA control (SS). (Courtesy of Dr. J. Old)




Fig. 6.26 α^0 -Thalassaemia: antenatal diagnosis by gap-PCR analysis. The common α^0 -thalassaemia deletion mutations are diagnosed using primers which bind to flanking sequences on either side of the deletion breakpoint. For the MED deletion, the primer pair (1 & 2) produce an amplified product of 650 base pairs (bp) in size. The primers are too far apart to amplify the normal DNA sequence, so a third primer (3) is included that is

complementary to the deleted sequence near one of the breakpoints. This produces a normal fragment of 1000 bp. The gel electrophoresis photograph shows the mother's DNA (lane 1, heterozygous), father's DNA (lane 2, heterozygous), CVS DNA (lane 3, normal) and two homozygous control DNAs (lanes 4 & 5). (Courtesy of Dr. J. Old)

Gap-PCR analysis is useful for detecting gene deletions in α -thalassaemia (Fig. 6.26), for $\delta\beta$ -thalassaemia, and Hb Lepore. Small deletions and point mutations are diagnosed by cycle-sequencing of PCR product using fluorescent labels and analysis of the fragments on a capillary-based automatic sequencer. This approach is useful for rare and unknown mutations, for confirming prenatal diagnosis of β -thalassaemia and sickle cell disorders by other PCR methods, and for the rare non-deletion α^+ -thalassaemia mutations that result in severe Hb H hydrops fetalis syndrome.

Pre-implantation genetic diagnosis which avoids the need for pregnancy termination involves performing conventional *in vitro* fertilization, followed by removing one or two cells from the blastomeres on day 3. PCR is used to detect thalassaemia mutations so that unaffected blastomeres can be selected for implantation. HLA typing can also be used to select an HLA matching blastomere matching a

previous thalassaemia major child. Ethical considerations are important in deciding to use these applications.

Fetal blood sampling

Fetal blood sampling may be performed in mid-second trimester and allows DNA study and protein synthesis studies.

Bibliography

- Adams R.J., Brambilla D., Optimizing Primary Stroke Prevention in Sickle Cell Anemia (STOP 2) Trial Investigators (2005) Discontinuing prophylactic transfusions used to prevent stroke in sickle cell disease. *N Engl J Med* 353,2769–78.
- Ahmed S., Saleem M., Modell B., Petrou M. (2002) Screening extended families for genetic hemoglobin disorders in Pakistan. *N Engl J Med* 347,1162–8.

- Anderson L.J., Holden S., Davis B. *et al.* (2001) Cardiovascular T2-star (T2*) magnetic resonance for the early diagnosis of myocardial iron overload. *Eur Heart J* 22,2171–9.
- Bain B.J. (2005) *Haemoglobinopathy Diagnosis*, 2nd edn. Blackwell Publishing, Oxford.
- Borgna-Pignatti C., Cappellini M.D., De Stefano P. *et al.* (2006) Cardiac morbidity and mortality in deferoxamine- or deferiprone-treated patients with thalassemia major. *Blood* 107,3733–7.
- Borgna-Pignatti C., Rugolotto S., De Stefano P. *et al.* (2004) Survival and complications in patients with thalassemia major treated with transfusion and deferoxamine. *Haematologica* 89,1187–93.
- Gladwin M.T., Sachdev V., Jison M.L. *et al.* (2004) Pulmonary hypertension as a risk factor for death in patients with sickle cell disease. *N Engl J Med* 350,886–95.
- Hankins J.S., Ware R.E., Rogers Z.R. *et al.* (2005) Long-term hydroxyurea therapy for infants with sickle cell anemia: the HUSOFT extension study. *Blood* 106,2269–75.
- Hoffbrand A.V. (2005) Deferiprone therapy for transfusional iron overload. *Best Pract Res Clin Haematol* 18,299–317.
- Kattamis A., Papassotiriou I., Palaiologu D. *et al.* (2006) The effects of erythropoietic activity and iron burden on hepcidin expression in patients with thalassemia major. *Haematologica* 91,809–12.
- Origa R., Bina P., Agus A. *et al.* (2005) Combined therapy with deferiprone and desferrioxamine in thalassemia major. *Haematologica* 90,1309–14.
- Papanikolaou G., Tzilianos M., Christakis J.I. *et al.* (2005) Hepcidin in iron overload disorders. *Blood* 105,4103–5.
- Pennell D.J., Berdoukas V., Karagiorga M. *et al.* (2006) Randomized controlled trial of deferiprone or deferoxamine in β -thalassemia major patients with asymptomatic myocardial siderosis. *Blood* 107,3738–44.
- Rund D. and Rachmilewitz E. (2005) Beta-thalassemia. *N Engl J Med* 353,1135–46.
- Steinberg M.H., Barton F., Castro O. *et al.* (2003) Effect of hydroxyurea on mortality and morbidity in adult sickle cell anemia: risks and benefits up to 9 years of treatment. *JAMA* 289,1645–51. Erratum in *JAMA* 290,756.
- Thein S.L. (2004) Genetic insights into the clinical diversity of beta thalassaemia. *Br J Haematol* 124,264–74.
- Weatherall D.J. and Clegg J.B. (2001) *The Thalassaemia Syndromes*, 4th edn. Blackwell Science, Oxford.

CHAPTER 7

7

The white cells 1: granulocytes, monocytes and their benign disorders

Granulocytes, 94
 Granulopoiesis, 96
 Clinical applications of myeloid growth factors, 97
 Monocytes, 99
 Disorders of neutrophil and monocyte function, 99

Causes of leucocytosis and monocytosis, 101
 Neutropenia, 104
 Histiocytic disorders, 106
 Bibliography, 107

The white blood cells (leucocytes) may be divided into two broad groups: the phagocytes and the immunocytes. Granulocytes, which include three types of cell—neutrophils (polymorphs), eosinophils and basophils—together with monocytes comprise the phagocytes. Their normal development and function, and benign disorders of white blood cells, are dealt with in this chapter. Only mature phagocytic cells and lymphocytes are found in normal peripheral blood (Table 7.1; Fig. 7.1). The lymphocytes, their precursor cells and plasma cells, which make up the immunocyte population, are considered in Chapter 8.

The function of phagocytes and immunocytes in protecting the body against infection is closely connected with two soluble protein systems of the body: immunoglobulins and complement. These proteins, which may also be involved in blood cell destruction in a number of diseases, are discussed together with the lymphocytes in Chapter 8.

Granulocytes

Neutrophil (polymorph)

This cell has a characteristic dense nucleus consisting of between two and five lobes, and a pale

Table 7.1 White cells: normal blood counts.

Adults	Blood count	Children	Blood count
<i>Total leucocytes</i>	$4.00\text{--}11.0 \times 10^9/\text{L}^*$	<i>Total leucocytes</i>	
Neutrophils	$2.5\text{--}7.5 \times 10^9/\text{L}^*$	Neonates	$10.0\text{--}25.0 \times 10^9/\text{L}$
Eosinophils	$0.04\text{--}0.4 \times 10^9/\text{L}$	1 year	$6.0\text{--}18.0 \times 10^9/\text{L}$
Monocytes	$0.2\text{--}0.8 \times 10^9/\text{L}$	4–7 years	$6.0\text{--}15.0 \times 10^9/\text{L}$
Basophils	$0.01\text{--}0.1 \times 10^9/\text{L}$	8–12 years	$4.5\text{--}13.5 \times 10^9/\text{L}$
Lymphocytes	$1.5\text{--}3.5 \times 10^9/\text{L}$		

* Normal black and Middle Eastern subjects may have lower counts. In normal pregnancy the upper limits are: total leucocytes $14.5 \times 10^9/\text{L}$, neutrophils $11 \times 10^9/\text{L}$.




Fig. 7.1 White blood cells (leucocytes): (a) neutrophil (polymorph); (b) eosinophil; (c) basophil; (d) monocyte; (e) lymphocyte.

cytoplasm with an irregular outline containing many fine pink-blue (azurophilic) or grey-blue granules (Fig. 7.1a). The granules are divided into primary, which appear at the promyelocyte stage, and secondary (specific) which appear at the myelocyte stage and predominate in the mature neutrophil. Both types of granule are lysosomal in origin: the primary contains myeloperoxidase, acid phosphatase and other acid hydrolases; the secondary contains collagenase, lactoferrin and lysozyme (see Fig. 7.7). The lifespan of neutrophils in the blood is only 6–10 h.

Neutrophil precursors

These do not normally appear in normal peripheral blood but are present in the marrow (Fig. 7.2). The earliest recognizable precursor is the myeloblast, a cell of variable size which has a large nucleus with fine chromatin and usually two to five nucleoli. The cytoplasm is basophilic and no cytoplasmic granules are present. The normal bone marrow contains up to 4% of myeloblasts. Myeloblasts give rise by

cell division to promyelocytes which are slightly larger cells and have developed primary granules in the cytoplasm. These cells then produce myelocytes which have specific or secondary granules. The nuclear chromatin is now more condensed and nucleoli are not visible. Separate myelocytes of the neutrophil, eosinophil and basophil series can be identified. The myelocytes give rise by cell division to metamyelocytes, non-dividing cells, which have an indented or horseshoe-shaped nucleus and a cytoplasm filled with primary and secondary granules. Neutrophil forms between the metamyelocyte and fully mature neutrophil are termed 'band', 'stab' or 'juvenile'. These cells may occur in normal peripheral blood. They do not contain the clear, fine filamentous distinction between nuclear lobes that is seen in mature neutrophils.

Monocytes

These are usually larger than other peripheral blood leucocytes and possess a large central oval or indented nucleus with clumped chromatin (Fig. 7.1d). The




Fig. 7.2. The formation of the neutrophil and monocyte phagocytes. Eosinophils and basophils are also formed in the marrow in a process similar to that for neutrophils.

abundant cytoplasm stains blue and contains many fine vacuoles, giving a ground-glass appearance. Cytoplasmic granules are also often present. The monocyte precursors in the marrow (monoblasts and promonocytes) are difficult to distinguish from myeloblasts and monocytes.

Eosinophils

These cells are similar to neutrophils, except that the cytoplasmic granules are coarser and more deeply red staining and there are rarely more than three nuclear lobes (Fig. 7.1b). Eosinophil myelocytes can be recognized but earlier stages are indistinguishable from neutrophil precursors. The blood transit time for eosinophils is longer than for neutrophils. They enter inflammatory exudates and have a special role in allergic responses, defence against parasites and removal of fibrin formed during inflammation.

Basophils

These are only occasionally seen in normal peripheral blood. They have many dark cytoplasmic granules

which overlie the nucleus and contain heparin and histamine (Fig. 7.1c). In the tissues they become mast cells. They have immunoglobulin E (IgE) attachment sites and their degranulation is associated with histamine release.

Granulopoiesis

The blood granulocytes and monocytes are formed in the bone marrow from a common precursor cell (see Fig. 1.2). In the granulopoietic series progenitor cells, myeloblasts, promyelocytes and myelocytes form a proliferative or mitotic pool of cells while the metamyelocytes, band and segmented granulocytes make up a post-mitotic maturation compartment (Fig. 7.3). Large numbers of band and segmented neutrophils are held in the marrow as a 'reserve pool' or storage compartment. The bone marrow normally contains more myeloid cells than erythroid cells in the ratio of 2 : 1 to 12 : 1, the largest proportion being neutrophils and metamyelocytes. In the stable or normal state, the bone marrow




Fig. 7.3 Neutrophil kinetics. CSF, colony-stimulating factor; G, granulocyte; IL, interleukin; M, monocyte; SCF, stem cell factor.

storage compartment contains 10–15 times the number of granulocytes found in the peripheral blood. Following their release from the bone marrow, granulocytes spend only 6–10 h in the circulation before moving into the tissues where they perform their phagocytic function. In the bloodstream there are two pools usually of about equal size: the circulating pool (included in the blood count) and the marginating pool (not included in the blood count). It has been estimated that they spend on average 4–5 days in the tissues before they are destroyed during defensive action or as the result of senescence.

Control of granulopoiesis: myeloid growth factors

The granulocyte series arises from bone marrow progenitor cells which are increasingly specialized. Many growth factors are involved in this maturation process including interleukin-1 (IL-1), IL-3, IL-5 (for eosinophils), IL-6, IL-11, granulocyte-macrophage colony-stimulating factor (GM-CSF), granulocyte CSF (G-CSF) and monocyte CSF (M-CSF) (see Fig. 1.7). The growth factors stimulate proliferation and differentiation and also affect the function of the mature cells on which they act (e.g. phagocytosis, superoxide generation and cytotoxicity in the case of neutrophils; phagocytosis,

cytotoxicity and production of other cytokines by monocytes).

Increased granulocyte and monocyte production in response to an infection is induced by increased production of growth factors from stromal cells and T lymphocytes, stimulated by endotoxin, IL-1 or tumour necrosis factor (TNF) (Fig. 7.4).

Clinical applications of myeloid growth factors

Clinical administration of G-CSF intravenously or subcutaneously has been found to produce a rise in neutrophils whereas administration of GM-CSF increases neutrophils, eosinophils and monocytes. G-CSF has become widely used in clinical practice and some of the indications are as follows. Short-acting G-CSF is given daily. There is also a longer acting pegolated (PEG) G-CSF which can be given once weekly.

Post-chemotherapy, radiotherapy or bone marrow transplantation In these situations, G-CSF accelerates haemopoietic recovery and shortens the period of neutropenia (Fig. 7.5). This may translate into a reduction of length of time in hospital, antibiotic usage and frequency of infection but periods of extreme neutropenia after intensive chemotherapy cannot be prevented.




Fig. 7.4 Regulation of haemopoiesis; pathways of stimulation of leucopoiesis by endotoxin, for example from infection. It is likely that endothelial and fibroblast cells release basal quantities of granulocyte-macrophage colony-stimulating factor (GM-CSF) and granulocyte colony-stimulating factor (G-CSF) in the normal resting state and that this is enhanced substantially by the monokine's tumour necrosis factor (TNF) and interleukin-1 (IL-1).




Fig. 7.5 Typical effect of granulocyte colony-stimulating factor (G-CSF) on recovery of neutrophils following autologous bone marrow transplantation.

Acute myeloid leukaemia G-CSF may be used after induction and consolidation chemotherapy to reduce hospital stay and antibiotic usage. In some protocols it is given with initial induction chemotherapy to mobilize blast cells into cell cycle, enhancing their sensitivity to chemotherapy.

Acute lymphoblastic leukaemia G-CSF is indicated to reduce the severity of neutropenia after intensive phases of therapy.

Myelodysplasia G-CSF has been given alone or in conjunction with erythropoietin in an attempt to

improve bone marrow function without accelerating leukaemic transformation. It is also used after intensive chemotherapy.

Severe neutropenia Both congenital and acquired neutropenia, including cyclical and drug-induced neutropenia, have been found to respond well to G-CSF.

Severe infection G-CSF has been used as an adjuvant to antimicrobial therapy.

Peripheral blood stem cell transplants G-CSF is used to increase the number of circulating multipotent progenitors, improving the harvest of sufficient peripheral blood stem cells for transplantation. It is also used to accelerate reconstitution after allogeneic or autologous transplantation.

Lymphomas G-CSF is given to reduce infection, delay in giving chemotherapy and hospitalization after chemotherapy. A single injection of pegolated G-CSF immediately after chemotherapy is often used.

Monocytes

Monocytes spend only a short time in the marrow and, after circulating for 20–40 h, leave the blood to enter the tissues where they mature and carry out their principal functions. Their extravascular lifespan after their transformation to macrophages may be as long as several months or even years. They may assume specific functions in different tissues (e.g. skin, gut, liver) (Fig. 7.6). One particularly important lineage is that of dendritic cells which are involved in antigen presentation to T cells (Chapter 8). GM-CSF and M-CSF are involved in their production and activation.

Disorders of neutrophil and monocyte function

The normal function of neutrophils and monocytes may be divided into three phases.

Chemotaxis (cell mobilization and migration) The phagocyte is attracted to bacteria or the site of inflammation by chemotactic substances released




Fig. 7.6 Reticuloendothelial system: distribution of macrophages.

from damaged tissues or by complement components and also by the interaction of leucocyte adhesion molecules with ligands on the damaged tissues. The leucocyte adhesion molecules also mediate recruitment, migration and interaction with other immune cells. They are also variously expressed on endothelial cells and platelets (Chapter 1).

Phagocytosis The foreign material (e.g. bacteria, fungi) or dead or damaged cells of the host are phagocytosed (Fig. 7.7). Recognition of a foreign particle is aided by opsonization with immunoglobulin or complement because both neutrophils and monocytes have Fc and C3b receptors. Opsonization of normal body cells (e.g. red cells or platelets) also makes them liable to destruction




Fig. 7.7 Phagocytosis and bacterial destruction. On entering the neutrophil, the bacterium is surrounded by an invaginated surface membrane and fuses with a primary lysosome to form a phagosome. Enzymes from the lysosome attack the bacterium. Secondary granules also fuse with the phagosomes, and new enzymes from these granules including lactoferrin attack the organism. Various types of activated oxygen, generated by glucose metabolism, also help to kill bacteria. Undigested residual bacterial products are excreted by exocytosis.

by macrophages of the reticuloendothelial system, as in autoimmune haemolysis, idiopathic (auto-immune) thrombocytopenic purpura or many of the drug-induced cytopenias.

Macrophages have a central role in antigen presentation: processing and presenting foreign antigens on human leucocyte antigen (HLA) molecules to the immune system. They also secrete a large number of growth factors and chemokines which regulate inflammation and immune responses.

Chemokines are chemotactic cytokines of which there are two main classes: CXC (α) chemokines, small (8–10 000 MW) pro-inflammatory cytokines which mainly act on neutrophils, and CC (β) chemokines such as macrophage inflammatory protein-1 α (MIP-1 α) and RANTES which act on monocytes, basophils, eosinophils and natural killer (NK) cells. Chemokines may be produced constitutively and control lymphocyte traffic under physiological conditions; inflammatory chemokines are induced or up-regulated by inflammatory stimuli. They bind to

and activate cells via chemokine receptors and play an important part in recruiting appropriate cells to the sites of inflammation. Chemokine receptors have been identified as co-receptors for human immunodeficiency virus (HIV) entry into cells.

Killing and digestion This occurs by oxygen-dependent and oxygen-independent pathways. In the oxygen-dependent reactions, superoxide (O_2^-), hydrogen peroxide (H_2O_2) and other activated oxygen (O_2) species, are generated from O_2 and reduced nicotinamide adenine dinucleotide phosphate (NADPH). In neutrophils, H_2O_2 reacts with myeloperoxidase and intracellular halide to kill bacteria; activated oxygen may also be involved. The non-oxidative microbicidal mechanisms involve microbicidal proteins. These may act alone (e.g. cathepsin G) or in conjunction with H_2O_2 (e.g. lysozyme, elastase). They may also act with a fall in pH within phagocytic vacuoles into which lysosomal enzymes are released. An additional factor,

lactoferrin—an iron-binding protein present in neutrophil granules—is bacteriostatic by depriving bacteria of iron and generating free radicals (Fig. 7.7). Finally, nitric oxide (NO) generated through NO synthase from L-arginine is another mechanism by which phagocytes kill microbes.

Defects of phagocytic cell function

Chemotaxis These defects occur in rare congenital abnormalities (e.g. 'lazy leucocyte' syndrome) and in more common acquired abnormalities either of the environment (e.g. corticosteroid therapy) or of the leucocytes themselves (e.g. in acute or chronic myeloid leukaemia, myelodysplasia and the myeloproliferative syndromes).

Phagocytosis These defects usually arise because of a lack of opsonization which may be caused by congenital or acquired causes of hypogammaglobulinaemia or lack of complement components.

Killing This abnormality is clearly illustrated by the rare X-linked or autosomal recessive chronic granulomatous disease that results from abnormal leucocyte oxidative metabolism. There is an abnormality affecting different elements of the respiratory burst oxidase or its activating mechanism. The patients have recurring infections, usually bacterial but sometimes fungal, which present in infancy or early childhood in most cases.

Other rare congenital abnormalities may also result in defects of bacterial killing (e.g. myeloperoxidase deficiency and the Chédiak–Higashi syndrome; see below). Acute or chronic myeloid leukaemia and myelodysplastic syndromes may also be associated with defective killing of ingested microorganisms.

Benign disorders

A number of the hereditary conditions may give rise to changes in granulocyte morphology.

Pelger–Huët anomaly In this uncommon condition bilobed neutrophils are found in the peripheral blood. Occasional unsegmented neutrophils are also seen. Inheritance is autosomal dominant.

May–Hegglin anomaly In this rare condition the neutrophils contain basophilic inclusions of RNA (resembling Döhle bodies) in the cytoplasm. There is an associated mild thrombocytopenia with giant platelets. Inheritance is autosomal dominant.

Other rare disorders In contrast to these two relatively benign anomalies, other rare congenital leucocyte disorders may be associated with severe disease. The Chédiak–Higashi syndrome is inherited in an autosomal recessive manner, and there are giant granules in the neutrophils, eosinophils, monocytes and lymphocytes accompanied by neutropenia, thrombocytopenia and marked hepatosplenomegaly. Abnormal leucocyte granulation or vacuolation is also seen in patients with rare mucopolysaccharide disorders (e.g. Hurler's syndrome).

Common morphological abnormalities Figure 7.8 shows some of the more common abnormalities of neutrophil morphology that can be seen in peripheral blood. Hypersegmented forms occur in megaloblastic anaemia, Döhle bodies and toxic changes in infection. The 'drumstick' appears on the nucleus of a proportion of the neutrophils in normal females and is caused by the presence of two X chromosomes. Pelger cells are seen in the benign congenital abnormality but also in patients with acute myeloid leukaemia or myelodysplasia.

Causes of leucocytosis and monocytosis

Neutrophil leucocytosis

An increase in circulating neutrophils to levels greater than $7.5 \times 10^9/L$ is one of the most frequently observed blood count changes. The causes of neutrophil leucocytosis are given in Table 7.2. Neutrophil leucocytosis is sometimes accompanied by fever as a result of the release of leucocyte pyrogens. Other characteristic features of reactive neutrophilia may include: (a) a 'shift to the left' in the peripheral blood differential white cell count (i.e. an increase in the number of band forms) and the occasional presence of more primitive cells such as metamyelocytes and myelocytes; (b) the presence of cytoplasmic toxic granulation




Fig. 7.8 Abnormal white blood cells. (a) Neutrophil leucocytosis: toxic changes shown by the presence of red-purple granules in the band form neutrophils. (b) Neutrophil leucocytosis: a Döhle body can be seen in the cytoplasm of the neutrophil. (c) Megaloblastic anaemia: hypersegmented oversized neutrophil in peripheral blood. (d) May-Hegglin anomaly: the

neutrophils contain basophilic inclusions 2–5 µm in diameter; there is an associated mild thrombocytopenia with giant platelets. (e) Pelger-Huët anomaly: coarse clumping of the chromatin in *pince nez* configuration. (f) Chédiak-Higashi syndrome: bizarre giant granules in the cytoplasm of a monocyte. (g) Alder's anomaly: coarse violet granules in the cytoplasm of a neutrophil.

Table 7.2 Causes of neutrophil leucocytosis.

Bacterial infections (especially pyogenic bacterial, localized or generalized)
Inflammation and tissue necrosis (e.g. myositis, vasculitis, cardiac infarct, trauma)
Metabolic disorders (e.g. uraemia, eclampsia, acidosis, gout)
Neoplasms of all types (e.g. carcinoma, lymphoma, melanoma)
Acute haemorrhage or haemolysis
Drugs (e.g. corticosteroid therapy (inhibits margination): lithium, tetracycline)
Chronic myeloid leukaemia, myeloproliferative disease, polycythaemia vera, myelofibrosis, essential thrombocythaemia
Treatment with myeloid growth factors (e.g. G-CSF, GM-CSF)
Rare inherited disorders
Asplenia

G- and GM-CSF, granulocyte and granulocyte-macrophage colony-stimulating factor.

and Döhle bodies (Fig. 7.8a,b); and (c) an elevated neutrophil alkaline phosphatase (NAP) score. For this the strength of the staining of each of 100 neutrophils is scored between 0 and 4. The maximum score is therefore 400; a normal score is between 20 and 100.

The leukaemoid reaction

The leukaemoid reaction is a reactive and excessive leucocytosis usually characterized by the presence of immature cells (e.g. myeloblasts, promyelocytes and myelocytes) in the peripheral blood. Occasionally,




Fig. 7.9 Eosinophilia.

Table 7.3 Causes of eosinophilia.

Allergic diseases, especially hypersensitivity of the atopic type (e.g. bronchial asthma, hay fever, urticaria and food sensitivity)
Parasitic diseases (e.g. amoebiasis, hookworm, ascariasis, tapeworm infestation, filariasis, schistosomiasis and trichinosis)
Recovery from acute infection
Certain skin diseases (e.g. psoriasis, pemphigus and dermatitis herpetiformis, urticaria and angioedema, atopic dermatitis)
Drug sensitivity
Polyarteritis nodosa, vasculitis, serum sickness
Graft-versus-host disease
Hodgkin's disease and some other tumours, especially clonal T-cell disorders
Metastatic malignancy with tumour necrosis
Hypereosinophilic syndrome
Chronic eosinophilic leukaemia
Myeloproliferative including systemic mastocytosis
Treatment with GM-CSF
Pulmonary syndromes
Eosinophilic pneumonia, transient pulmonary infiltrates (Loeffler's syndrome), allergic granulomatosis (Churg–Strauss syndrome), tropical pulmonary eosinophilia

GM-CSF, granulocyte–macrophage colony-stimulating factor.

lymphocytic reactions occur. Associated disorders include severe or chronic infections, severe haemolysis or metastatic cancer. Leukaemoid reactions are often particularly marked in children. Granulocyte changes such as toxic granulation and Döhle bodies and a high NAP score help to differentiate the leukaemoid reaction from chronic myeloid leukaemia (in which the NAP score is low).

Eosinophilic leucocytosis (eosinophilia)

The causes of an increase in blood eosinophils (Fig. 7.9) above $0.4 \times 10^9/L$ are listed in Table 7.3. Sometimes no underlying cause is found and if the eosinophil count is elevated ($>1.5 \times 10^9/L$) for over 6 months and associated with tissue damage then the hypereosinophilic syndrome is diagnosed. The

Table 7.4 Causes of monocytosis.

Chronic bacterial infections: tuberculosis, brucellosis, bacterial endocarditis, typhoid
 Connective tissue diseases—SLE, temporal arteritis, rheumatoid arthritis
 Protozoan infections
 Chronic neutropenia
 Hodgkin's disease, AML, and other malignancies
 Myelodysplasia (especially chronic myelomonocytic leukaemia)
 Treatment with GM-CSF or M-CSF

AML, acute myeloblastic leukemia; GM- and M-CSF, granulocyte-macrophage and macrophage colony-stimulating factor; SLE, systemic lupus erythematosus.

heart valves, skin and lungs may be affected and treatment is usually with steroids or cytotoxic drugs. In 25% of cases a clonal T-cell population is present. In other cases of chronic eosinophilia, a clonal cytogenetic abnormality is present in the bone marrow and the term chronic eosinophilic leukaemia is used. In 50% of cases there is a novel fusion gene on chromosome 4q caused by deletion of part of the long arm causing joining of the fragments of FIP1L1 and PDGFRA. This creates an active tyrosine kinase and the condition responds to imatinib (p. 181).

Basophil leucocytosis (basophilia)

An increase in blood basophils above $0.1 \times 10^9/L$ is uncommon. The usual cause is a myeloproliferative disorder such as chronic myeloid leukaemia or polycythaemia vera. Reactive basophil increases are sometimes seen in myxoedema, during smallpox or chickenpox infection and in ulcerative colitis.

Monocytosis

A rise in blood monocyte count above $0.8 \times 10^9/L$ is infrequent. The conditions listed in Table 7.4 may be responsible.

Neutropenia

The lower limit of the normal neutrophil count is $2.5 \times 10^9/L$ except in black people and in the Middle East where $1.5 \times 10^9/L$ is normal. When the absolute

Table 7.5 Causes of neutropenia.

Selective neutropenia

Congenital

Kostmann's syndrome

Acquired

Drug-induced

Anti-inflammatory drugs (phenylbutazone)
 Antibacterial drugs (chloramphenicol, co-trimoxazole, sulfasalazine, imipenem)
 Anticonvulsants (phenytoin, carbamazepine)
 Antithyroids (carbimazole)
 Hypoglycaemics (tolbutamide)
 Phenothiazines (chlorpromazine, thioridazine)
 Psychotropics and antidepressants (clozapine, mianserin, imipramine)
 Miscellaneous (gold, penicillamine, mepacrine, frusemide, deferiprone)

Benign (racial or familial)

Cyclical

Immune

Autoimmune
 Systemic lupus erythematosus
 Felty's syndrome
 Hypersensitivity and anaphylaxis

Large granular lymphocytic leukaemia (p. 195)

Infections

Viral (e.g. hepatitis, influenza, HIV)
 Fulminant bacterial infection (e.g. typhoid, miliary tuberculosis)

Part of general pancytopenia (see Table 20.1)

Bone marrow failure
 Splenomegaly

HIV, human immunodeficiency virus.

neutrophil level falls below $0.5 \times 10^9/L$ the patient is likely to have recurrent infections and when the count falls to less than $0.2 \times 10^9/L$ the risks are very serious, particularly if there is also a functional defect. Neutropenia may be selective or part of a general pancytopenia (Table 7.5).

Congenital neutropenia

Kostmann's syndrome is an autosomal recessive disease presenting in the first year of life with life-threatening infections. Most cases are caused by

mutations of the gene coding for neutrophil elastase. G-CSF produces a clinical response although marrow fibrosis and acute myeloid leukaemia may supervene.

Drug-induced neutropenia

A large number of drugs have been implicated (Table 7.5) and may induce neutropenia either by direct toxicity or immune-mediated damage.

Cyclical neutropenia

This is a rare syndrome with 3–4-week periodicity. Severe but temporary neutropenia occurs. Monocytes tend to rise as the neutrophils fall. Mutation of the gene for neutrophil elastase underlies some cases.

Autoimmune neutropenia

In some cases of chronic neutropenia an autoimmune mechanism can be demonstrated. The antibody may be directed against one of the neutrophil-specific antigens (e.g. NA, NB).

Idiopathic benign neutropenia

An increase in the marginating fraction of blood neutrophils and a corresponding reduction in the circulating fraction is one cause of benign neutropenia. Many normal Africans and other races, especially in the Middle East, have a low peripheral blood neutrophil count without excess margination. These subjects have no increased susceptibility to infection and the bone marrow appears normal although there is diminished neutrophil production.

Finally, the term chronic idiopathic neutropenia is used for unexplained acquired neutropenia (neutrophil count below normal for the ethnic group), without phasic variations or underlying disease. It is more common in females and thought to be brought about by immune cells causing inhibition of myelopoiesis in the bone marrow.

Clinical features

Severe neutropenia is particularly associated with infections of the mouth and throat. Painful and often intractable ulceration may occur at these sites (Fig. 7.10), on the skin or at the anus. Septicaemia rapidly supervenes. Organisms carried as commensals by normal individuals, such as *Staphylococcus*




Fig. 7.10 Ulceration of the tongue in severe neutropenia.

epidermidis or Gram-negative organisms in the bowel, may become pathogens. Other features of infections associated with severe neutropenia are described on p. 149.

Diagnosis

Bone marrow examination is useful in determining the level of damage in granulopoiesis (i.e. whether there is reduction in early precursors or whether there is reduction only of circulating and marrow neutrophils with late precursors remaining in the marrow). Marrow aspiration and trephine biopsy may also provide evidence of leukaemia, myelodysplasia or other infiltration.

Management

The treatment of patients with acute severe neutropenia is described on p. 150. In many patients with drug-induced neutropenia spontaneous recovery occurs within 1–2 weeks after stopping the drug. Patients with chronic neutropenia have recurrent infections which are mainly bacterial in origin although fungal and viral infections (especially

herpes) also occur. Early recognition and vigorous treatment with antibiotics, antifungal or antiviral agents, as appropriate, is essential. Prophylactic antibacterial agents (e.g. oral co-trimoxazole or ciprofloxacin and colistin) and antifungal agents (e.g. oral amphotericin and fluconazole or itraconazole) may be of value in reducing the incidence and severity of infections caused by severe neutropenia. The haemopoietic growth factor G-CSF may be used to stimulate neutrophil production and is effective in a variety of benign chronic neutropenic states. Corticosteroid therapy or splenectomy has been associated with good results in some patients with autoimmune neutropenia. Rituximab (anti-CD20) may also be effective. Conversely, corticosteroids impair neutrophil function and should not be used indiscriminately in patients with neutropenia.

Histiocytic disorders

A classification of the histiocytic disorders is given in Table 7.6.

Dendritic cells

These are specialized antigen-presenting cells found mainly in the skin, lymph nodes, spleen and

thymus. They comprise myeloid- and monocyte-derived cells including Langerhans' cells and a lymphocyte-derived subset. Their primary role is in antigen presentation to T and B lymphocytes (p. 115).

Langerhans' cell histiocytes

Langerhans' cell histiocytosis (LCH) includes diseases previously called histiocytosis X, Letterer-Siwe disease, Hand-Schüller-Christian disease and eosinophilic granuloma. The disease may be single organ or multisystem. There is a clonal proliferation of CD1a-positive cells. The multisystem disease affects children in the first 3 years of life with hepatosplenomegaly, lymphadenopathy and eczematous skin symptoms. Localized lesions may occur especially in the skull, ribs and long bones, the posterior pituitary causing diabetes insipidus, the central nervous system, gastrointestinal tract and lungs. The lesions include Langerhans' cells (characterized by the presence of tennis racquet-shaped Birbeck granules in electron-microscopy sections), eosinophils, lymphocytes, neutrophils and macrophages.

Haemophagocytic lymphohistiocytosis (haemophagocytic syndrome)

This is a rare, recessively inherited or more frequently acquired disease, usually precipitated by a viral (especially Epstein-Barr or herpes viruses), bacterial or fungal infection or occurring in association with tumours. Often the patient is immunocompromised. Patients present with fever and pancytopenia, often with splenomegaly and liver dysfunction. There are increased numbers of histiocytes in the bone marrow which ingest red cells, white cells and platelets (Fig. 7.11). Clinical features are fever, pancytopenia and multiorgan dysfunction often with lymphadenopathy, hepatic and splenic enlargement, coagulopathy and CNS signs. Treatment is of the underlying infection, if known, with support care. T-cell activation is implicated in the aetiology. Chemotherapy with etoposide, corticosteroids, ciclosporin or Rituximab (anti-CD20) may be tried. The condition is often fatal.

Table 7.6 Classification of histiocytic disorders.

Dendritic cell-related

Langerhans' cell histiocytosis

Solitary dendritic cell histiocytoma

Macrophage-related

Haemophagocytic lymphohistiocytosis

primary (familial)

secondary—infection, drug, tumour

Sinus histiocytosis with massive lymphadenopathy

Malignancies


AML FAB type M4 and M5 (p. 158)

Chronic myelomonocytic leukaemia (p. 186)


Dendritic and macrophage-related sarcomas (localized or disseminated)

AML, acute myeloid leukaemia;

FAB, French-American-British.



(a)



(b)

Fig. 7.11 Haemophagocytic lymphohistiocytosis: bone marrow aspirates showing histiocytes that have ingested red cells, erythroblasts and neutrophils.

Sinus histiocytosis with massive lymphadenopathy

This is also known as the Rosai–Dorfman syndrome. There is painless, chronic cervical lymphadenopathy. There may be fever and weight loss. The histology is typical and the condition subsides over months or years.

Bibliography

- Brito-Babapulle F. (2003) The eosinophilias, including the idiopathic hypereosinophilic syndrome. *Br J Haematol* 121,203–23.
- Charo I.F. and Ransohoff R.M. (2006) The many roles of chemokines and chemokine receptors in inflammation. *N Engl J Med* 354,610–21.
- Dale D.C. (ed.) (2002) Severe chronic neutropenia. *Semin Hematol* 39,73–141.
- Duffield J.S. (2003) The inflammatory macrophage: a story of Jekyll and Hyde. *Clin Sci (Lond)* 104,27–38.
- Kyono W. and Coates T.D. (2002) A practical approach to neutrophil disorders. *Pediatr Clin North Am* 49,929–71.
- Lakshman R. and Finn A. (2001) Neutrophil disorders and their management. *J Clin Pathol* 54,7–19.
- Ozer H., Armitage J.O., Bennett C.L. *et al.* (2000) 2000 update of recommendations for the use of hematopoietic colony-stimulating factors: evidence-based, clinical practice guidelines. American Society of Clinical Oncology Growth Factors Expert Panel. *J Clin Oncol* 18,3558–85.
- Stock W. and Hoffman R. (2000) White blood cells 1: non-malignant disorders. *Lancet* 355,1351–7.
- Tefferi A. (ed.) (2005) Eosinophilic and mast cell disorders. *Acta Haematol* 114,52–60.
- Tefferi A., Patnaik M.M. and Pardanani A. (2006) Eosinophilia: secondary, clonal and idiopathic. *Br J Haematol* 133,468–92.
- Zeidler C. and Welte K. (2002) Kostmann syndrome and severe congenital neutropenia. *Semin Hematol* 39,82–8.

CHAPTER 8

8

The white cells 2: lymphocytes and their benign disorders

Lymphocytes, 108
Immunoglobulins, 109
Antigen–receptor gene arrangements, 112
Complement, 113
The immune response, 115

Lymphocytosis, 117
Immunodeficiency, 119
Differential diagnosis of lymphadenopathy, 121
Bibliography, 122

Lymphocytes are the immunologically competent cells that assist the phagocytes in defence of the body against infection and other foreign invasion (Fig. 8.1). Two unique features characteristic of the immune system are the ability to generate *antigenic specificity* and the phenomenon of *immunological memory*. A complete description of the functions of lymphocytes is beyond the scope of this book, but information essential to an understanding of the diseases of the lymphoid system, and of the role of lymphocytes in haematological diseases, is included here.

Lymphocytes

In postnatal life, the bone marrow and thymus are the *primary lymphoid organs* in which lymphocytes develop (Fig. 8.2). The *secondary lymphoid organs* in which specific immune responses are generated are the lymph nodes, spleen and lymphoid tissues of the alimentary and respiratory tracts.

B and T lymphocytes

The immune response depends upon two types of lymphocytes, B and T cells (Table 8.1), which derive from the haemopoietic stem cell. B cells mature in the bone marrow and circulate in the peripheral blood until they undergo recognition of antigen. The B-cell

receptor is membrane-bound immunoglobulin and after activation this is secreted as free soluble immunoglobulin. At this point they mature into memory B cells or plasma cells. The latter home to the bone marrow and have a characteristic morphology with an eccentric round nucleus with a 'clock-face' chromatin pattern and strongly basophilic cytoplasm (Fig. 8.1d).

T cells develop from cells that have migrated to the thymus where they differentiate into mature T cells during passage from the cortex to the medulla. During this process, self-reactive T cells are deleted (negative selection) whereas T cells with some specificity for host human leucocyte antigen (HLA) molecules are selected (positive selection). The mature helper cells express CD4 and cytotoxic cells express CD8 (Table 8.1). The cells also express one of two T-cell antigen receptor heterodimers, $\alpha\beta$ (>90%) or $\gamma\delta$ (<10%), and recognize antigen only when it is presented at a cell surface (see below).

Natural killer cells

Natural killer (NK) cells are cytotoxic CD8⁺ cells that lack the T-cell receptor (TCR). They are large cells with cytoplasmic granules and typically express surface molecules CD16 (Fc receptor), CD56 and CD57. NK cells are designed to kill target cells that have a low level of expression of HLA class I




Fig. 8.1 Lymphocytes:
(a) small lymphocyte;
(b) activated lymphocyte;
(c) large granular lymphocyte;
(d) plasma cell.

molecules such as may occur during viral infection or on a malignant cell. NK cells do this by displaying a number of receptors for HLA molecules on their surface. When HLA is expressed on the target cell these deliver an inhibitory signal into the NK cell. When HLA molecules are absent on the target cell this inhibitory signal is lost and the NK cell can then kill its target. In addition, NK cells display antibody-dependent cell-mediated cytotoxicity (ADCC). In this, antibody binds to antigen on the surface of the target cell and then NK cells bind to the Fc portion of the bound antibody and kill the target cell.

Lymphocyte circulation

Lymphocytes in the peripheral blood migrate through *post-capillary venules* into the substance of the lymph nodes or into the spleen or bone marrow. T cells home to the perifollicular zones of the

cortical areas of lymph nodes (paracortical areas) (Fig. 8.2) and to the periarteriolar sheaths surrounding the central arterioles of the spleen. B cells selectively accumulate in follicles of the lymph nodes and spleen. Lymphocytes return to the peripheral blood via the efferent lymphatic stream and the thoracic duct. CD4 helper cells predominate in normal peripheral blood and germinal centres, but in the marrow and gut the major T-cell subpopulation is CD8 positive.

Immunoglobulins

These are a group of proteins produced by plasma cells and B lymphocytes that bind to antigen. They are divided into five subclasses or *isotypes*: immunoglobulin G (IgG), IgA, IgM, IgD and IgE. IgG, the most common, contributes approximately 80% of normal serum immunoglobulin and is further




Fig. 8.2 Primary and secondary lymphoid organs and blood.

subdivided into four *subclasses*: IgG₁, IgG₂, IgG₃ and IgG₄. IgA is subdivided into two types. IgM is usually produced first in response to antigen, IgG subsequently and for a more prolonged period. The same cell can switch from IgM to IgG, or to IgA or

IgE synthesis. IgA is the main immunoglobulin in secretions, particularly of the gastrointestinal tract. IgD and IgE (involved in delayed hypersensitivity reactions) are minor fractions. Some important biochemical and biological properties of the three

Table 8.1 Functional aspects of T and B cells.

	T cells	B cells
Origin	Thymus	Bone marrow
Tissue distribution	Parafollicular areas of cortex in nodes, peri-arteriolar in spleen	Germinal centres of lymph nodes, spleen, gut, respiratory tract; also subcapsular and medullary cords of lymph nodes
Blood	80% of lymphocytes; CD4 > CD8	20% of lymphocytes
Membrane receptors	TCR for antigen	BCR (= immunoglobulin) for antigen
Function	CD8 ⁺ : CMI against intracellular organisms CD4 ⁺ : T-cell help for antibody production and generation of CMI	Humoral immunity by generation of antibodies
Characteristic surface markers	CD1 CD2 CD3 CD4 or 8 CD5 CD6 CD7 MHC class I MHC class II when activated	CD19 CD20 CD22 CD9 (pre B cells) CD10 (precursor B cells) CD79 MHC class I and II
Genes rearranged	TCR α , β , γ , δ	IgH, Ig κ , Ig λ

BCR, B-cell receptor; C, complement; CMI, cell-mediated immunity; IFN, interferon; Ig, immunoglobulin; MHC, major histocompatibility complex; TCR, T-cell receptor; TNF, tumour necrosis factor.

Table 8.2 Some properties of the three main classes of immunoglobulin (Ig).

	IgG	IgA	IgM
Molecular weight	140 000	140 000	900 000
Sedimentation constant	7S	7S	19S
Normal serum level (g/L)	6.0–16.0	1.5–4.5	0.5–1.5
Present in	Serum and extracellular fluid	Serum and other body fluids (e.g. of bronchi and gut)	Serum only
Complement fixation	Usual	Yes (alternative pathway)	Usual and very efficient
Placental transfer	Yes	No	No
Heavy chain	(γ_{1-4})	α (α_1 or α_2)	μ

main immunoglobulin subclasses are summarized in Table 8.2.

The immunoglobulins are all made up of the same basic structure (Fig. 8.3) consisting of two heavy chains which are called gamma (γ) in IgG, alpha (α) in IgA, mu (μ) in IgM, delta (δ) in IgD and epsilon (ϵ) in IgE, and two light chains—kappa (κ) or lambda (λ)—which are common to all five immunoglobulins. The heavy and light chains each have highly

variable regions which give the immunoglobulin specificity, and constant regions in which there is virtual complete correspondence in amino acid sequence in all antibodies of a given isotype (e.g. IgA, IgG) or isotype subclass (e.g. IgG₁, IgG₂). IgG antibody can be broken into a constant Fc fragment and two highly variable Fab fragments. IgM molecules are much larger because they consist of five subunits.




Fig. 8.3 Basic structure of an immunoglobulin molecule. Each molecule is made up of two light (κ or λ) (blue areas) and two heavy (purple) chains, and each chain is made up of variable (V) and constant (C) portions, the V portions including the antigen-binding site. The heavy chain (μ , δ , γ , ϵ or α) varies according to the immunoglobulin class. IgA molecules form dimers, while IgM forms a ring of five molecules. Papain cleaves the molecules into an Fc fragment and two Fab fragments.

The main role of immunoglobulins is defence of the body against foreign organisms. However, they also have a vital role in the pathogenesis of a number of haematological disorders. Secretion of a specific

immunoglobulin from a monoclonal population of lymphocytes or plasma cells causes *paraproteinaemia* (p. 216). Bence-Jones protein found in the urine in some cases of myeloma consists of a monoclonal secretion of light chains or light-chain fragments (either κ or λ). Immunoglobulins may bind to blood cells in a variety of immune disorders and cause their agglutination (e.g. in cold agglutinin disease; p. 67) or destruction following direct complement lysis or after elimination by the reticuloendothelial system.

Antigen-receptor gene arrangements

Immunoglobulin gene rearrangements

The immunoglobulin heavy-chain and κ and λ light-chain genes occur on chromosomes 14, 2 and 22, respectively. In the germline state, the heavy-chain gene consists of separate segments for variable (V), diversity (D), joining (J) and constant (C) regions. Each of the V, D and J regions contain a number (n) of different *gene segments* (Fig. 8.4). In cells not committed to immunoglobulin synthesis these gene segments remain in their separate germline state. During early differentiation of B cells there is rearrangement of heavy-chain genes so that one of the V heavy-chain segments combines with one of the D segments, which has itself already combined with one of the J segments. Thus, they form a transcriptionally active gene for the heavy chain. The protein coding segments of the C region mRNA




Fig. 8.4 Rearrangement of a heavy-chain immunoglobulin gene. One of the V segments is brought into contact with a D, a J and a C (in this case C_μ) segment, forming an active transcriptional gene from which the corresponding mRNA is produced. The DJ rearrangement precedes VDJ joining. The class of immunoglobulin depends on which of the nine constant regions (1μ , 1δ , 4γ , 2α , 1ϵ) is used.




Fig. 8.5 The sequence of immunoglobulin gene rearrangement, antigen and immunoglobulin expression during early B-cell development. Intracytoplasmic CD22 is a feature of very early B cells. HLA, human leucocyte antigen; TdT, terminal deoxynucleotidyl transferase.

are joined to the V region after splicing out intervening RNA. The class of immunoglobulin that is secreted depends on which of the nine (4γ , 2α , 1μ , 1δ and 1ϵ) constant regions is used. Diversity is introduced by the variability of which V segment joins with which D and with which J segment. In the arbitrary example shown in Fig. 8.4, V_2 joins with D_1 and J_2 . Additional diversity is generated by the enzyme terminal deoxynucleotidyl transferase (TdT), which inserts a variable number of new bases into the DNA of the D region at the time of gene rearrangement.

Similar rearrangements occur during generation of the light-chain gene (Fig. 8.5). Enzymes called *recombinases* are needed both in B and T cells to join up the adjacent pieces of DNA after excision of intervening sequences. These recognize certain heptamer- and nonamer-conserved sequences flanking the various gene segments. Mistakes in recombinase activity play an important part in the chromosome translocations of B- or T-cell malignancy.

T-cell receptor gene rearrangements

The vast majority of T cells contain a TCR composed of a heterodimer of α and β chain. In a minority of T

cells, the TCR is composed of γ and δ chains. The α , β , γ and δ genes of the TCRs each include V, D, J and C regions. During T-cell ontogeny, rearrangements of these gene segments occur in a similar fashion to those for immunoglobulin genes, thus creating T cells expressing a wide variety (10^8 or more) of TCR structures (Fig. 8.6). TdT is involved in creating additional diversity and the same recombinase enzymes used in B cells are involved in joining up TCR gene segments.

Complement

This consists of a series of plasma proteins constituting an *amplification enzyme system* which is capable of lysis of bacteria (or of blood cells) or can 'opsonize' (coat) bacteria or cells so that they are phagocytosed. The complement sequence consists of nine major components—C1, C2, etc.—which are activated in turn (denoted thus C1—) and form a cascade, resembling the coagulation sequence (Fig. 8.7). The most abundant and pivotal protein is C3, which is present in plasma at a level of approximately 1.2 g/L. The early (opsonizing) stages leading to coating of the cells with C3b can occur by two different pathways:




Fig. 8.6 The sequence of events during early T-cell development. The earliest events appear to be the expression of surface CD7, intranuclear terminal deoxynucleotidyl transferase (TdT) and intracytoplasmic CD3 followed by T-cell receptor (TCR) gene rearrangement. Early medullary thymocytes may express both CD4 and CD8, but they then lose one or other of these structures.




Fig. 8.7 The complement (C) sequence. The activated factors are denoted by a bar over the number. Both pathways generate a C3 convertase. In the classic pathway, the convertase is the major (b) component of C4 and C2 ($\bar{C}4b2b$). In the alternate pathway, it is the combination of C3b and the major fragment (b) of factor B ($\bar{C}3bBb$).

- 1 The *classical pathway* usually activated by IgG or IgM coating of cells; or
- 2 The *alternate pathway*, which is more rapid and activated by IgA, endotoxin (from Gram-negative bacteria) and other factors (Fig. 8.7).

Macrophages and neutrophils have C3b receptors and they phagocytose C3b-coated cells. C3b is degraded to C3d detected in the direct antiglobulin (Coombs') test using an anticomplement agent (p. 343). If the complement sequence goes to

completion there is generation of an active phospholipase that punches holes in the cell membrane (e.g. of the red cell or bacterium), causing direct lysis. The complement pathway also generates the biologically active fragments C3a and C5a which act directly on phagocytes to stimulate the respiratory burst (p. 100). Both may trigger anaphylaxis by release of mediators from tissue mast cells and basophils which causes vasodilatation and increased permeability.




Fig. 8.8 Antigen receptors on lymphocytes and their interaction with antigen. (a) The B-cell antigen receptor is membrane-bound immunoglobulin. Two heavy chains (H) are covalently bonded to two light chains (L). This antigen-binding unit is associated with the CD79 heterodimer which acts as a signal transduction unit. (b) The T-cell receptor consists of a number of components that together constitute the CD3 complex.

The immune response

One of the most striking features of the immune system is its capacity to produce a highly *specific* response. For both T and B cells this specificity is achieved by the presence of a particular receptor on the lymphocyte surface (Fig. 8.8). Naïve (or virgin) B and T lymphocytes which leave the bone marrow and thymus are resting cells that are not in cell division. They recirculate in the lymphatic system. Specialized macrophages called dendritic cells (DCs; p. 99) process antigens before presenting them to B and T lymphocytes—they are therefore known as *antigen-presenting cells* (APCs). The immune system contains many different lymphocytes. Each of these lymphocytes has a receptor that shows differences in structure from that of any other lymphocyte, and consequently will bind to only a restricted number of antigens. T and B cells undergo clonal expansion if they meet an APC that is presenting an antigen that can trigger their antigen receptor molecules. At this stage, lymphocytes may develop into effector cells (such as plasma cells or cytotoxic T cells) or memory cells.

Two antigen-binding chains (α , β) are associated with several proteins (γ , δ , ϵ , ζ) that mediate signal transduction. Antigen is recognized in the form of short peptides held on the surface of HLA molecules. CD8⁺ T cells interact with peptide on a class I HLA molecule and the CD8 heterodimer interacts with the α_3 domain of the class I protein.


DC precursors constitutively migrate at low levels from blood into tissues but their rate of migration is increased at the site of inflammation. Immature DCs are efficient at macropinocytosis which allows them to capture antigens from the environment. DCs can be matured by a variety of stimuli such as inflammatory cytokines—tumour necrosis factor- α (TNF- α) and interleukin-1 (IL-1)—and viral and bacterial products such as lipopolysaccharide (LPS) or double-stranded (ds) RNA. Mature DCs express high levels of co-stimulatory molecules and can efficiently present antigen to naïve antigen-specific T cells.

T cells are unable to bind antigen free in solution and require it to be presented on APCs in the form of peptides held on the surface of HLA molecules (Fig. 8.8b). T cells therefore recognize not only the antigen, but also 'self' HLA molecules and are known as *HLA-restricted*. The CD4 molecule on helper cells recognizes class II (HLA-DP, -DQ and -DR) molecules, whereas the CD8 molecule recognizes class I (HLA-A, -B and -C) molecules (Fig. 21.5). The antigen recognition site of the TCR is joined to several other subunits in the CD3 complex


which together mediate signal transduction. During these structural interactions the cells release cytokines such as IL-1, -2, -4 and -10 which act to modify expansion of activated cells. Depending on their cytokine production, CD4⁺ T cells can be broadly subdivided into T helper type 1 (Th1) and Th2 cells. Th1 cells produce mainly IL-2, TNF- β and γ -interferon (IFN- γ), and are important in boosting cell-mediated immunity

(and granuloma formation), whereas Th2 cells produce IL-4 and IL-10 and are mainly responsible for providing help for antibody production. An imbalance in the ratio of these two subsets may be partly responsible for some forms of immunodeficiency.

Antigen-specific immune responses are generated in *secondary lymphoid organs* and commence when antigen is carried into a lymph node (Fig. 8.9)



(a)



(b)

Fig. 8.9 (a) Structure of a lymph node. (b) Lymph node showing germinal follicles surrounded by a darker mantle zone rim and lighter, more diffuse marginal and T-zone areas.




Fig. 8.10 Generation of a germinal centre. B cells activated by antigen migrate from the T zone to the follicle where they undergo massive proliferation. Cells enter the dark zone as centroblasts and accumulate mutations in their immunoglobulin V genes. Cells then pass back into the

light zone (Fig. 8.8a) as centrocytes. Only those cells that can interact with antigen on follicular dendritic cells and receive signals from antigen-specific T cells (Fig. 8.8) are selected and migrate out as plasma cells and memory cells. Cells not selected die by apoptosis.

on dendritic cells. B cells recognize antigen through their surface immunoglobulin and although most antibody responses require help from antigen-specific T cells, some antigens such as polysaccharides can lead to T-cell independent antibody production. T cells are screened for recognition of antigen and if a T cell makes an interaction it migrates into the follicle. In the follicle, germinal centres arise as a result of continuing response to antigenic stimulation (Fig. 8.10). These consist of follicular dendritic cells (FDCs), which are loaded with antigen, B cells and activated T cells which have migrated up from the T zone. Proliferating B cells move to the dark zone of the germinal centre as *centroblasts* where they undergo somatic mutation of their immunoglobulin variable-region genes. Their progeny are known as *centrocytes* and these must be selected by antigen on FDCs otherwise they will undergo apoptosis. If selected they become memory B cells or plasma cells (Fig. 8.10). Plasma cells

migrate to the bone marrow and produce high affinity antibody. Although they contain intracellular immunoglobulin they do not express surface immunoglobulin.

Lymphocytosis

Lymphocytosis often occurs in infants and young children in response to infections that produce a neutrophil reaction in adults. Conditions particularly associated with lymphocytosis are listed in Table 8.3.

Glandular fever is a general term for a disease characterized by fever, sore throat, lymphadenopathy and atypical lymphocytes in the blood. It may be caused by primary infection with Epstein-Barr virus (EBV), cytomegalovirus, human immunodeficiency virus (HIV) or toxoplasma. EBV infection, otherwise known as infectious mononucleosis, is the most common cause.

Table 8.3 Causes of lymphocytosis.

Infections

acute: infectious mononucleosis, rubella, pertussis, mumps, acute infectious lymphocytosis, infectious hepatitis, cytomegalovirus, HIV, herpes simplex or zoster

chronic: tuberculosis, toxoplasmosis, brucellosis, syphilis

Chronic lymphoid leukaemias (Chapter 15)

Acute lymphoblastic leukaemia

Non-Hodgkin's lymphoma (some)

Thyrototoxicosis

HIV, human immunodeficiency virus.

Infectious mononucleosis

This is caused by primary infection with EBV and occurs only in a minority of infected individuals—in most cases infection is subclinical. The disease is characterized by a lymphocytosis caused by clonal expansions of T cells reacting against B lymphocytes infected with EBV. The disease is associated with a high titre of heterophile ('reacting with cells of another species') antibody which reacts with sheep, horse or ox red cells.

Clinical features

The majority of patients are between the ages of 15 and 40 years. A prodromal period of a few days occurs with lethargy, malaise, headaches, stiff neck and a dry cough. In established disease the following features may be found.

- 1 Bilateral cervical lymphadenopathy is present in 75% of cases. Symmetrical generalized lymphadenopathy occurs in 50% of cases. The nodes are discrete and may be tender.
- 2 Over half of patients have a sore throat with inflamed oral and pharyngeal surfaces. Follicular tonsillitis is frequently seen.
- 3 Fever may be mild or severe.
- 4 A morbilliform rash, severe headache and eye signs (e.g. photophobia, conjunctivitis and periorbital oedema) are not uncommon. The rash may follow therapy with amoxicillin or ampicillin.
- 5 Palpable splenomegaly occurs in over half of patients and hepatomegaly in approximately 15%. Approximately 5% of patients are jaundiced.
- 6 Peripheral neuropathy, severe anaemia (caused

by autoimmune haemolysis) or purpura (caused by thrombocytopenia) are less frequent complications.

Diagnosis

Pleomorphic atypical lymphocytosis A moderate rise in white cell count (e.g. $10\text{--}20 \times 10^9/\text{L}$) with an absolute lymphocytosis is usual, and some patients have even higher counts. Large numbers of atypical lymphocytes are seen in the peripheral blood film (Fig. 8.11). These T cells are variable in appearance but most have nuclear and cytoplasmic features similar to those seen during reactive lymphocyte transformation. The greatest number of atypical lymphocytes are usually found between the seventh and 10th day of the illness.

Heterophile antibodies Heterophile antibodies against sheep red cells may be found in the serum at high titres. These form the basis of the Paul-Bunnell test—the antibodies are not absorbed by guinea-pig kidney cells but are absorbed by ox red cells (Fig. 8.12). Modern slide screening tests, such as the *monospot test*, substitute formalinized horse red cells for the sheep cells used in the Paul-Bunnell test. Highest titres occur during the second and third week and the antibody persists in most patients for 6 weeks.

EBV antibody If viral diagnostic facilities are available, a rise in the titre of antibody against the EBV capsid antigen may be demonstrated during the first 2–3 weeks. Specific antibody to the EBV nuclear antigen develops later and persists for life.

Haematological abnormalities other than the atypical lymphocytosis are frequent. Occasional patients develop an autoimmune haemolytic anaemia. The IgM autoantibody is typically of the 'cold' type and usually shows 'i' blood group specificity. Thrombocytopenia is frequent and an autoimmune thrombocytopenic purpura occurs in a smaller number of patients.

Differential diagnosis

The differential diagnosis of infectious mononucleosis includes cytomegalovirus, HIV or toxoplasmosis infection; acute leukaemia; influenza; rubella; bacterial tonsillitis; and infectious hepatitis.




Fig. 8.11 Infectious mononucleosis: representative 'reactive' T lymphocytes in the peripheral blood film of a 21-year-old man (see also Fig. 8.1b). (c)

Treatment

In the great majority of patients only symptomatic treatment is required. Corticosteroids are sometimes given to those with severe systemic symptoms. Patients characteristically develop an erythematous rash if given ampicillin therapy. Most patients recover fully 4–6 weeks after initial symptoms. However, convalescence may be slow and associated with severe malaise and lethargy.

Lymphopenia

Lymphopenia may occur in severe bone marrow failure, with corticosteroid and other immunosuppressive therapy, in Hodgkin's disease and with

widespread irradiation. It also occurs in a variety of immunodeficiency syndromes, the most important of which is HIV infection (p. 328).

Immunodeficiency

A large number of inherited or acquired deficits in any of the components of the immune system can cause an impaired immune response with increased susceptibility to infection (Table 8.4). A primary lack of T cells (as in AIDS) leads not only to bacterial infections, but also to viral, protozoal, fungal and mycobacterial infections. In some cases, however, lack of specific subsets of T cells which control B-cell maturation may lead to a secondary lack of B-cell




Fig. 8.12 Serological diagnosis of acute Epstein-Barr virus (EBV) infection. (a) IgM heterophile antibodies (shown here as antibody monomers for ease of illustration) against sheep red cells are used in the Paul-Bunnell test. These are not absorbed out by guinea-pig kidney but do react with ox red cells. A positive Paul-Bunnell test should show a titre of sheep red cell agglutination that is at least fourfold lower after absorption with ox red cells but not greater than threefold lower after absorption with guinea-pig kidney. (b) The commonly used monospot test detects agglutination of sheep red cells or formalized horse red cells only. (c) After acute infection heterophile and EBV-specific IgM are present for approximately 3 months. IgG antibodies to viral capsid antigen (VCA) and EBV nuclear antigen (EBNA) remain elevated for years.

Table 8.4 Classification of immunodeficiencies.

<i>Primary</i>	
B cell (antibody deficiency)	X-linked agammaglobulinaemia, acquired common variable hypogammaglobulinaemia, selective IgA or IgG subclass deficiencies
T cell	Thymic aplasia (DiGeorge's syndrome), PNP deficiency
Mixed B and T cell	Severe combined immune deficiency (as a result of ADA deficiency or other causes); Bloom's syndrome; ataxia-telangiectasia; Wiskott-Aldrich syndrome
<i>Secondary</i>	
B cell (antibody deficiency)	Myeloma; nephrotic syndrome, protein-losing enteropathy
T cell	AIDS; Hodgkin's lymphoma, non-Hodgkin's lymphoma; drugs: steroids, ciclosporin, azathioprine, fludarabine, etc.
T and B cell	Chronic lymphocytic leukaemia, post-bone marrow transplantation and post-chemotherapy/radiotherapy

ADA, adenosine deaminase; AIDS, acquired immune deficiency syndrome; Ig, immunoglobulin; PNP, purine nucleoside phosphorylase.

function as in many cases of common variable immunodeficiency, which may develop in children or adults of either sex. In others, a primary defect of B cells or of APCs is present. X-linked agammaglobulinaemia is caused by a failure of B-cell development and pyogenic bacterial infections dominate the clinical course. Immunoglobulin replacement therapy can be given by monthly courses of intravenous immunoglobulin. Rare syndromes include aplasia of the thymus, severe combined (T and B) immunodeficiency as a result of adenosine deaminase deficiency and selective deficiencies of IgA or IgM. Acquired immune deficiency occurs after cytotoxic chemotherapy or radiotherapy and is particularly pronounced after stem cell transplantation where dysregulation of the immune system persists for 1 year or more and is responsible for a high incidence of serious viral infections (e.g. with cytomegalovirus or herpes zoster). Immunodeficiency is

also frequently associated with tumours of the lymphoid system including chronic lymphocytic leukaemia and myeloma.

Differential diagnosis of lymphadenopathy

The principal causes of lymphadenopathy are listed in Fig. 8.13. The clinical history and examination give essential information. The age of the patient, length of history, associated symptoms of possible infectious or malignant disease, whether the nodes are painful or tender, consistency of the nodes and whether there is generalized or local lymphadenopathy are all important. The size of the liver and spleen are assessed. In the case of local node enlargement, inflammatory or malignant disease in the associated lymphatic drainage area are particularly considered.




Fig. 8.13 Causes of lymphadenopathy. ALL, acute lymphoblastic leukaemia; CLL, chronic lymphocytic leukaemia; SLE, systemic lupus erythematosus. Malignancies are listed in red.

Further investigations will depend on the initial clinical diagnosis but it is usual to include a full blood count, blood film and erythrocyte sedimentation rate (ESR). Chest X-ray, monospot test, cytomegalovirus and *Toxoplasma* titres, and anti-HIV and Mantoux (Heaf) tests are frequently needed. In many cases, it will be essential to make a histological diagnosis by node biopsy but a fine needle aspirate may sometimes avoid the need for this. Computed tomography (CT) scanning is valuable in determining the presence and extent of deep node enlargement. Subsequent investigations will depend on the diagnosis made and the patient's particular features. In some cases of deep node enlargement, where enlarged superficial nodes are not available for biopsy, bone marrow, liver biopsies, CT or ultrasound guided trucut node biopsy may be needed in an attempt to reach a histological diagnosis and avoid the need for a diagnostic laparotomy.

Bibliography

- Hsu K.C. and Dupont B. (2005) Natural killer cell receptors: regulating innate immune responses to hematologic malignancy. *Semin Hematol* 42,91–103.
- Janeway C.A., Travers P., Walport M., Shlomchik M.J. (eds) (2004) *Immunobiology*, 6th edn. Garland Science, New York and Oxford.
- Krogsgaard M. and Davis M.M. (2005) How T cells 'see' antigen. *Nat Immunol* 3,239–45.
- Randolph G.J., Angeli V., Swartz M.A. (2005) Dendritic-cell trafficking to lymph nodes through lymphatic vessels. *Nat Rev Immunol* 8,617–28.
- Webster A.D.B. (2005) Immunodeficiency diseases. In: Hoffbrand A.V., Catovsky D., Tuddenham E.G.D. (eds) *Postgraduate Haematology*, 5th edn. Blackwell Publishing, Oxford: 370–9.

CHAPTER 9

9

The spleen

The anatomy and circulation of the spleen, 123
The functions of the spleen, 123
Extramedullary haemopoiesis, 124
Imaging the spleen, 125
Splénomegaly, 125

Hypersplenism, 126
Hyposplenism, 127
Splenectomy, 127
Prevention of infection in hyposplenic patients, 127
Bibliography, 128

The spleen has an important and unique role in the function of the haemopoietic and immune systems. As well as being directly involved in many diseases of these systems, a number of important clinical features are associated with hypersplenic and hyposplenic states.

The anatomy and circulation of the spleen

The spleen lies under the left costal margin, has a normal weight of 150–250 g and a length of between 5 and 13 cm. It is normally not palpable but becomes palpable when the size is increased to over 14 cm.

Blood enters the spleen through the splenic artery which then divides into *trabecular arteries* which permeate the organ and give rise to *central arterioles* (Fig. 9.1). The majority of the arterioles end in *cords* which lack an endothelial lining and form an open blood system unique to the spleen with a loose reticular connective tissue network lined by fibroblasts and many macrophages. The blood re-enters the circulation by passing across the endothelium into venous *sinuses*. Blood then passes into the splenic vein and so back into the general circulation. The cords and sinuses form the *red pulp* which forms 75% of the spleen and has an essential role in monitoring the integrity of red blood cells (see below). A minority of the splenic vasculature is

closed in which the arterial and venous systems are connected by a continuous endothelial layer.

The central arterioles are surrounded by a core of lymphatic tissue known as *white pulp* which has an organization similar to lymph nodes (Fig. 9.1). The *periarteriolar lymphatic sheath* (PALS) lies directly around the arteriole and is equivalent to the T zone of the lymph node (p. 116). B cell follicles are found adjacent to the PALS and these are surrounded by the *marginal zone* and *perifollicular zone* which are rich in macrophages and dendritic cells. Lymphocytes migrate into white pulp from the sinuses of the red pulp or from vessels that end directly in the marginal and perifollicular zones.

There are both rapid (1–2 min) and slow (30–60 min) blood circulations through the spleen. The slow circulation becomes increasingly important in splénomegaly.

The functions of the spleen

The spleen is the largest filter of the blood in the body and several of its functions are derived from this.

Control of red cell integrity

The spleen has an essential role in the 'quality control' of red cells. Excess DNA, nuclear remnants (Howell-Jolly bodies) and siderotic granules are




Fig. 9.1 Schematic representation of the blood circulation in the spleen. Most blood flows in an 'open' circulation through splenic cords and regains entry into the circulation through the venous sinuses.




Fig. 9.2 Splenic atrophy: peripheral blood film showing Howell-Jolly bodies, Pappenheimer bodies (siderotic granules; see p. 25) and misshapen cells.

removed (Fig. 9.2). Aged or abnormal red cells are also removed from the circulation. In the relatively hypoxic environment of the red pulp, and because of plasma skimming in the cords, the membrane flexibility of aged and abnormal red cells is impaired and they are retained within the sinus where they are ingested by macrophages.

Immune function


The lymphoid tissue in the spleen is in a unique position to respond to antigens filtered from the blood and entering the white pulp. Macrophages and dendritic cells in the marginal zone initiate an immune response and then present antigen to B and T cells to start adaptive immune responses. This arrangement is highly efficient at initiating immune responses to encapsulated bacteria and explains the susceptibility of hyposplenic patients to these organisms.

Extramedullary haemopoiesis

The spleen, like the liver, undergoes a transient period of haemopoiesis at around 3–7 months of fetal life but is not a site of erythropoiesis in the adult. However, haemopoiesis may be re-established in both organs as *extramedullary haemopoiesis*, in disorders such as myelofibrosis or in chronic severe haemolytic and megaloblastic anaemias. Extramedullary haemopoiesis may result either from reactivation of dormant stem cells within the spleen or homing of stem cells from the bone marrow to the spleen.



(a)



(b)



(c)

Imaging the spleen

Ultrasound is the most frequently used technique to image the spleen (Fig. 9.3). This can also detect whether or not blood flow in the splenic, portal and hepatic veins is normal, as well as liver size and consistency. Computed tomography (CT) is preferable for detecting structural detail and any associated lymphadenopathy (e.g. for lymphoma staging). Magnetic resonance imaging (MRI) also gives improved fine detail structure. Positron emission tomography (PET) is used particularly for detecting residual disease after treatment of lymphoma.

Splenomegaly

Splenic size is increased in a wide range of conditions (Table 9.1). Splenomegaly is usually apparent under the left costal margin but massive splenomegaly may be felt in the right iliac fossa (see Fig. 19.4). The spleen moves with respiration and a medial splenic notch may be palpable in some cases. In developed countries the most common causes of splenomegaly are infectious mononucleosis, haematological malignancy and portal hypertension, whereas malaria and schistosomiasis are more prevalent on a global scale (Table 9.1). Chronic myeloid leukaemia, myelofibrosis, lymphoma, Gaucher's disease, malaria, leishmaniasis and schistosomiasis are potential causes of massive splenomegaly.

Tropical splenomegaly syndrome

A syndrome of massive splenomegaly of uncertain aetiology has been found frequently in many malarious zones of the tropics including Uganda, Nigeria, New Guinea and the Congo. Smaller numbers of patients with this disorder are seen in southern

Fig. 9.3 Imaging the spleen. (a) Ultrasound of spleen showing splenomegaly (15.3 cm). (b) Normal spleen (10 cm) on computed tomography (CT) scan. (c) CT scan: The spleen is enlarged and shows multiple low density areas. A diagnosis of diffuse large cell B lymphoma was made histologically after splenectomy. (Figs (a) and (b) courtesy of Dr T. Ogunremi)

Table 9.1 Causes of splenomegaly.

Haematological

Chronic myeloid leukaemia*
 Chronic lymphocytic leukaemia
 Acute leukaemia
 Malignant lymphoma*
 Chronic myelofibrosis*
 Polycythaemia vera
 Hairy cell leukaemia
 Thalassaemia major or intermedia*
 Sickle cell anaemia (before splenic infarction)
 Haemolytic anaemias
 Megaloblastic anaemia

Portal hypertension

Cirrhosis
 Hepatic, portal, splenic vein thrombosis

Storage diseases

Gaucher's disease*
 Niemann–Pick disease
 Histiocytosis X

Systemic diseases

Sarcoidosis
 Amyloidosis
 Collagen diseases—systemic lupus erythematosus,
 rheumatoid arthritis
 Systemic mastocytosis

Infections

Acute: septicaemia, bacterial endocarditis, typhoid,
 infectious mononucleosis
 Chronic: tuberculosis, brucellosis, syphilis, malaria,
 leishmaniasis,* schistosomiasis*

*Tropical**

Possibly caused by malaria

* Possible causes of massive (>20 cm) splenomegaly.

Arabia, the Sudan and Zambia. Previously, such terms as 'big spleen disease', 'cryptogenic splenomegaly' and 'African macroglobulinaemia' have been used to describe this syndrome.

While it seems probable that malaria is the fundamental cause of tropical splenomegaly syndrome, this disease is not the result of active malarial infection as parasitaemia is usually scanty and malarial pigment is not found in biopsy material from the liver and spleen. The available evidence

suggests that an abnormal host response to the continual presence of malarial antigen results in a reactive and relatively benign lymphoproliferative disorder that predominantly affects the liver and spleen.

Splenomegaly is usually gross and the liver is also enlarged. Portal hypertension may be a feature. The anaemia is often severe and the lowest haemoglobin levels are found in subjects with the largest spleens. While leucopenia is usual, some patients develop a marked lymphocytosis. The moderate degree of thrombocytopenia present does not often cause spontaneous bleeding. Serum IgM levels are high and fluorescent techniques reveal high titres of malarial antibody.

Although splenectomy corrects the pancytopenia, there is an increased risk of fulminant malarial infection. Trials of antimalarial prophylaxis (e.g. proguanil and other antimalarial drugs) have proved successful in the management of many affected patients, supporting the view that a continuing presence of malarial antigen is needed for the perpetuation of the lymphoproliferation associated with this syndrome. Resistant cases have also been treated successfully with chemotherapy.

Hypersplenism

Normally, only approximately 5% (30–70 mL) of the total red cell mass is present in the spleen although up to half of the total marginating neutrophil pool and 30% of the platelet mass may be located there. As the spleen enlarges, the proportion of haemopoietic cells within the organ increases such that up to 40% of the red cell mass, and 90% of platelets (Fig. 23.9), may be pooled in an enlarged spleen. *Hypersplenism* is a clinical syndrome that can be seen in any form of splenomegaly. It is characterized by:

- Enlargement of the spleen;
- Reduction of at least one cell line in the blood in the presence of normal bone marrow function; and
- Evidence of increased release of premature cells, such as reticulocytes or immature platelets, from the bone marrow into the blood.

Splenectomy may be indicated if the hypersplenism is symptomatic. It is followed by a rapid improvement in the peripheral blood count.

Table 9.2 Causes of hyposplenism and blood film features.

Causes	Blood film features
Splenectomy	<i>Red cells</i>
Sickle cell disease	Target cells
Essential thrombocythaemia	Acanthocytes
Adult gluten-induced enteropathy	Irregularly contracted or crenated cells
Dermatitis herpetiformis	Howell–Jolly bodies (DNA remnants)
Rarely	Siderotic (iron) granules
inflammatory bowel disease	(Pappenheimer bodies)
splenic arterial venous thrombosis	
	<i>White cells</i>
	± Mild lymphocytosis, monocytosis
	<i>Platelets</i>
	± Thrombocytosis

Hyposplenism

Functional hyposplenism is revealed by the blood film findings of Howell–Jolly bodies or Pappenheimer bodies (siderotic granules on iron staining; Fig. 9.2). The most obvious cause is surgical removal of the spleen but hyposplenism can also occur in sickle cell anaemia, gluten-induced enteropathy, inflammatory bowel disease and splenic arterial thrombosis (Table 9.2).

Splenectomy

Surgical removal of the spleen may be indicated for treatment of haematological disorders as well as after splenic rupture or for splenic tumours or cysts (Table 9.3). Splenectomy is usually performed by

open abdominal laparotomy but it is also possible to remove the spleen by laparoscopic surgery.

The platelet count can often rise dramatically in the early postoperative period, reaching levels of up to $1000 \times 10^9/L$ and peaking at 1–2 weeks. Thrombotic complications are seen in some patients and prophylactic aspirin or heparin are often required during this period. Long-term alterations in the peripheral blood cell count may also be seen, including a persistent thrombocytosis, lymphocytosis or monocytosis.

Prevention of infection in hyposplenic patients

Patients with hyposplenism are at lifelong increased risk of infection from a variety of organisms. This is seen particularly in children under the age of 5 years and those with sickle cell anaemia. The most characteristic susceptibility is to encapsulated bacteria such as *Streptococcus pneumoniae*, *Haemophilus influenzae* type B and *Neisseria meningitidis*. *Streptococcus pneumoniae* is a particular concern and can cause a rapid and fulminant disease. Malaria tends to be more severe in splenectomized individuals. Measures to reduce the risk of serious infection include the following:

- 1 The patient should be informed about the increased susceptibility to infection and advised to carry

Table 9.3 Indications for splenectomy.

Splenic rupture
Chronic immune thrombocytopenia
Haemolytic anaemia (some cases), e.g. hereditary spherocytosis, autoimmune haemolytic anaemia, thalassaemia major or intermedia
Chronic lymphocytic leukaemia and lymphomas
Myelofibrosis
Tropical splenomegaly

Table 9.4 Recommendations for vaccination of patients with hyposplenism.

Vaccine	Time of vaccination	Revaccination schedule	Comments
1 Pneumococcal polyvalent vaccine	{ If possible, at least 2 weeks prior to splenectomy. Alternatively, 2 weeks post-splenectomy for all 3 vaccines	5 yearly	Assessment of antibody response may be useful
2 Haemophilus influenzae Type b conjugate		Not required	
3 Meningococcal C conjugate		Not required	Not required if previously vaccinated
4 Influenza	As soon as available for seasonal protection	Annual	Standard killed vaccine

a card about their condition. They should be counselled about the increased risk of infection on foreign travel, including that from malaria and tick bites.

2 Prophylactic oral phenoxymethylpenicillin is recommended for life. Erythromycin may be prescribed for patients allergic to penicillin. A supply of tablets may also be given to the patient to take in the event of onset of fever before medical care is available.

3 Vaccination against pneumococcus, haemophilus, meningococcus and influenza infection is recommended (Table 9.4). All types of vaccine, including live vaccines, can be given safely to hyposplenic individuals although the immune response to vaccination may be impaired.

Bibliography

- Bowdler A.J. (ed.) (2002) *The Complete Spleen: Structure, Function and Clinical Disorders*, 2nd edn. Humana Press, Towota, NJ.
- Davies J.M., Barnes R., Milligan D. *et al.* for the British Committee for Standards in Haematology. Working Party of the Haematology/Oncology Task Force. (2002) Update of guidelines for the prevention and treatment of infection in patients with an absent or dysfunctional spleen. *Clin Med* 2,440–3.
- Giebink G.S. (2001) The prevention of pneumococcal disease in children. *N Engl J Med* 345,1177–83.
- Mebius R.E. and Kraal G. (2005) Structure and function of the spleen. *Nat Rev Immunol* 5,606–16.
- Shadbolt C. and Peters A.M. (2003) The spleen. In: Peters A.M. (ed.) *Nuclear Medicine in Radiological Diagnosis*. Martin Dunitz, London.

CHAPTER 10

10

The aetiology and genetics of haematological malignancies

The aetiology of haemopoietic malignancy, 129

The genetics of haemopoietic malignancy, 131

Chromosome nomenclature, 134

Genetic abnormalities associated with haematological malignancies, 135

Specific examples of translocations associated with haematological malignancy, 138

Diagnostic methods used to study malignant cells, 138

Value of genetic markers in management of haematological malignancy, 143

Bibliography, 146

The haemopoietic malignancies are clonal diseases that derive from a single cell in the marrow or peripheral lymphoid tissue which has undergone a genetic alteration (Fig. 10.1). In this chapter we discuss the aetiology and genetic basis of haematological malignancy and subsequent chapters discuss the diagnosis and management of these conditions.

The aetiology of haemopoietic malignancy

Exactly how genetic mutations accumulate in haemopoietic malignancies is largely unknown. As in most diseases it is the combination of genetic background and environmental influence that determines the risk of developing a malignancy. However, in the majority of individual cases neither a genetic susceptibility nor an environmental agent is apparent.

Inherited factors

The incidence of leukaemia is greatly increased in some genetic diseases, particularly Down's syndrome (where acute leukaemia occurs with a 20–30-fold increased frequency), Bloom's syndrome, Fanconi's anaemia, ataxia telangiectasia, Klinefelter's

syndrome and Wiskott–Aldrich syndrome. There is also a weak familial tendency in diseases such as acute myeloid leukaemia (AML), B-cell chronic lymphocytic leukaemia (CLL), Hodgkin's lymphoma and non-Hodgkin's lymphoma (NHL) although the genes predisposing to this risk are largely unknown.

Environmental influences

Chemicals

Chronic exposure to benzene may cause bone marrow hypoplasia, dysplasia and chromosome abnormalities and is an unusual cause of myelodysplasia or AML. Other industrial solvents and chemicals less commonly cause leukaemia.

Drugs

The alkylating agents (e.g. chlorambucil, mustine (chlormethine), melphalan, procarbazine and nitrosoureas—BCNU, CCNU) predispose to AML, especially if combined with radiotherapy or if used to treat patients with lymphocytic or plasmacytic disorders. Epipodophyllotoxins such as etoposide are powerful antileukaemic agents but their use is associated with a risk of the development of




Fig. 10.1 Theoretical graph to show the replacement of normal bone marrow cells by a clonal population of malignant cells arising by successive mitotic divisions from a single cell with an acquired genetic alteration.

secondary leukaemias associated with balanced translocations including that of the *MLL* gene at 11q23.

Radiation

Radiation, especially to the marrow, is leukemogenic. This is illustrated by an increased incidence of all types of leukaemia (except CLL) in survivors of the atom bomb explosions in Japan.

Infection

A proportion of cases of childhood acute lymphoblastic leukaemia (ALL) are initiated by genetic mutations that occur during development *in utero* (Fig. 10.2). Studies in identical twins have shown that both may be born with the same chromosomal abnormality (e.g. t(12; 21)). This has presumably arisen spontaneously in a progenitor cell that has




Fig. 10.2 Prenatal origin of acute lymphoblastic leukaemia (ALL) in a pair of identical twins. ALL was diagnosed in the first twin at age 5 years and in the second at age 14 years. Both tumours had an identical t(12; 21) translocation indicating probable origin of the leukaemic clone *in utero* and dissemination to both twins via a shared placental blood supply. Because of the prolonged latency

of the ALL it is presumed that a secondary event is required to initiate the development of frank leukaemia. At the time of the diagnosis of ALL in twin 1 the t(12; 21) translocation could be detected in the bone marrow of twin 2. It is likely that such a 'fetal origin' of childhood ALL occurs in a significant number of sporadic ALL cases. (After J.L. Wiemels *et al.* 1999)

Table 10.1 Infections associated with haemopoietic malignancies.

Infection	Tumour
<i>Virus</i>	
HTLV-1	Adult T-cell leukaemia/lymphoma
Epstein–Barr virus	Burkitt's and Hodgkin's lymphomas; PTLD
HHV-8	Primary effusion lymphoma; multicentric Castleman's disease
HIV-1	High-grade B-cell lymphoma
<i>Bacteria</i>	
<i>Helicobacter pylori</i>	Gastric lymphoma (MALT)
<i>Protozoa</i>	
Malaria	Burkitt's lymphoma

HHV-8, human herpes virus 8; HIV, human immunodeficiency virus; HTLV-1, human T-lymphotropic virus type 1; MALT, mucosa-associated lymphoid tissue; PTLD, post-transplant lymphoproliferative disease.

passed from one twin to the other as a result of the shared placental circulation. Environmental exposure during pregnancy may be important for this first event. One twin may develop ALL early (e.g. at age 4) because of a second transforming event while the other remains well or develops ALL later. The *TEL-AML1* translocation is present in the blood of approximately 10% of newborn infants but only 1 in 100 of these go on to develop ALL at a later date. The mechanism of the 'second genetic hit' within the tumour cell is unclear but an abnormal response of the immune system to infection is suggested by epidemiological studies. Children with a high level of social activity, notably those attending early nursery daycare, have a reduced incidence of ALL, whereas those living in more isolated communities and who have a reduced exposure to common infections in the first years of life have a higher risk.

Viruses (Table 10.1)

Viral infection is associated with several types of haemopoietic malignancy. The retrovirus human T-lymphotropic virus type 1 (HTLV-1) is the cause of adult T-cell leukaemia/lymphoma (ATLL) (p. 214) although most people infected with this virus do not develop the tumour. Epstein–Barr virus (EBV) DNA is integrated into the genome of endemic (African) Burkitt's lymphoma cells but rarely in sporadic Burkitt's lymphoma cells. It is also the cause of post-transplant lymphoproliferative disease (PTLD) which

develops during immunosuppressive therapy after solid organ transplantation, of many cases of lymphoma associated with HIV infection and is present in a proportion of patients with Hodgkin's disease. Human herpes virus 8 (HHV-8; Kaposi's sarcoma-associated virus (KSHV)) is associated with Kaposi's sarcoma and primary effusion lymphoma (PEL).

HIV infection is associated with an increased incidence of lymphomas at unusual sites such as the central nervous system. The HIV-associated lymphomas are usually of B-cell origin and of high-grade histology.

Bacteria

Helicobacter pylori infection has been implicated in the pathogenesis of gastric mucosa B-cell (MALT) lymphoma (p. 212).

Protozoa

Endemic Burkitt's lymphoma occurs in the tropics, particularly in malarial areas. It is thought that malaria may alter host immunity and predispose to tumour formation as a result of EBV infection.

The genetics of haemopoietic malignancy

Malignant transformation occurs as a result of the accumulation of genetic mutations in cellular genes. The genes that are involved in the development of




Fig. 10.3 Proliferation of normal cells depends on a balance between the action of proto-oncogenes (a) and tumour-suppressor genes (b). In a malignant cell this balance is disturbed leading to uncontrolled cell division.

cancer can be divided broadly into two groups: *oncogenes* and *tumour-suppressor genes*.

Oncogenes

Oncogenes arise because of gain-of-function mutations in normal cellular genes called *proto-oncogenes* (Fig. 10.3). Proto-oncogenes are involved in a variety of important cellular processes, often in the pathway by which external signals are transduced to the cell nucleus to activate genes. Oncogenic versions are generated when the activity of proto-oncogenes is increased or they acquire a novel function. This can occur in a number of ways including translocation, mutation or duplication. One of the striking features of haematological malignancies (in contrast to most solid tumours) is their high frequency of chromosomal translocations. A subset of proto-oncogenes are involved in control of apoptosis (e.g. *BCL-2* which is overexpressed in follicular lymphoma) (p. 209).

Tumour-suppressor genes

Tumour-suppressor genes may acquire loss-of-function mutations, usually by point mutation or deletion, which lead to malignant transformation

(Fig. 10.3). Tumour-suppressor genes commonly act as components of control mechanisms which regulate entry of the cell from the G_1 phase of the cell cycle into the S phase or passage through the S phase to G_2 and mitosis (Fig. 1.8). Examples of oncogenes and tumour-suppressor genes involved in haemopoietic malignancies are shown in Table 10.2. The most significant tumour-suppressor gene in human cancer is p53 which is mutated or inactivated in over 50% of cases of malignant disease, including many haemopoietic tumours.

Clonal progression

Malignant cells appear to arise as a multistep process with acquisition of mutations in different intracellular pathways (Fig. 10.4). Another feature of malignancy is clonal progression. In many cases the disease develops new characteristics during its clinical course and this may be accompanied by new chromosome changes. Selection of subclones may occur during treatment or reflect disease acceleration. Drug resistance can arise through a variety of molecular mechanisms. In one example the cells express a protein that actively pumps a number of different drugs to the outside of the cells (multidrug resistance, MDR).

Table 10.2 Some of the more frequent genetic abnormalities in leukaemia, lymphoma and myeloproliferative diseases.

Disease	Genetic abnormality*	Oncogene(s) involved
<i>Myeloid</i>		
AML M ₂	t(8; 21)	ETO and CBF α (AML1)
	t(6; 9)	DEK, CAN
AML M ₃	t(15; 17)	RAR α , PML
AML M ₄	inv(16), del(16q)	CBF β , MYH11
AML M ₅	del(11q); t(9; 11); t(11; 19)	MLL
AML (all types)	Nucleotide insertion Mutation, ITD	Nucleophosmin Flt3
MDS	-5/del(5q) -7/del(7q) Point mutation	Unclear N-RAS
Secondary myeloid leukaemia	11q23 translocations	MLL gene
CML	t(9; 22)	ABL, BCR
Myeloproliferative disease	Point mutation	JAK2
Chronic eosinophilic syndrome	20q- 4q12 del	FIP1L1-PDGFR α fusion
<i>Lymphoid</i>		
Precursor B lineage ALL	t(12; 21) t(4; 11) t(9; 22) t(1; 19) Hyperdiploidy Hypodiploidy	TEL, AML1 AF4, MLL (ALL1, HRX) ABL, BCR PBX-1, E2A
Burkitt's lymphoma, B-ALL	t(8; 14) [†] t(2; 8) t(8; 22)	MYC to IgH locus MYC to IgK locus MYC to Ig λ locus
T-ALL	t(1; 14) t(8; 14) [†] t(11; 14) t(7; 9)	TAL-1 to TCR δ locus MYC RBTN-1 or RBTN-2 TAL-2 to TCR β locus
Follicular lymphoma	t(14; 18)	BCL-2 to IgH
Anaplastic lymphoma	t(2; 5)	ALK to NPM
Mantle cell lymphoma	t(11; 14) [†] Mutation	BCL-1 (cyclin D1) to IgH p53 and ATM
CLL	Trisomy 12 13q14 deletion 11q22-23 deletion or mutation 17p deletion or mutation 6q21 deletion	Unknown Unknown ATM p53 Unknown
MALT B-cell lymphoma	t(1; 14)	BCL10 to IgH locus
T-PLL	inv(14q) or t(14q) Mutations	ATM

ALL, acute lymphoblastic leukaemia; AML, acute myeloid leukaemia; CLL, chronic lymphocytic leukaemia; CML, chronic myeloid leukaemia; ITD, internal tandem duplication; MALT, mucosa-associated lymphoid tissue; MDS, myelodysplasia; PLL, prolymphocytic leukaemia.

* Several chromosomal abnormalities may occur in patients with many of the conditions listed (e.g. Flt-3 mutation occurs in 30% of AML cases with or without chromosome translocations).

[†] The breakpoints on chromosome 14 are in different positions in T-ALL, B-ALL, mantle cell lymphoma and Burkitt's lymphoma.




Fig. 10.4 Multistep origin of a malignant tumour. Successive mutations lead to a growth advantage of one subclone.

Chromosome nomenclature

The normal somatic cell has 46 chromosomes and is called *diploid*; ova or sperm have 23 chromosomes and are called *haploid*. The chromosomes occur in pairs and are numbered 1–22 in decreasing size order; there are two sex chromosomes, XX in females, XY in males. *Karyotype* is the term used to describe the chromosomes derived from a mitotic cell which have been set out in numerical order (see Fig. 13.1d). A somatic cell with more or less than 46 chromosomes is termed *aneuploid*; more than 46 is *hyperdiploid*, less than 46 *hypodiploid*; 46 but with chromosome rearrangements, *pseudodiploid*.

Each chromosome has two arms, the shorter called 'p', the longer called 'q'. These meet at the *centromere* and the ends of the chromosomes are called *telomeres*. On staining each arm divides into regions




Fig. 10.5 A schematic representation of a chromosome. The bands may be divided into subbands according to staining pattern.

numbered outwards from the centromere and each region divides into bands (Fig. 10.5).

When a whole chromosome is lost or gained, a – or + is put in front of the chromosome number. If part of the chromosome is lost it is prefixed with *del* for deletion. If there is extra material replacing part of a chromosome the prefix *add* for additional material is used. Chromosome translocations are denoted by *t*, the chromosomes involved placed in brackets with the lower numbered chromosome first. The prefix *inv* describes an inversion where part of the chromosome has been inverted to run in the opposite direction. An *isochromosome*, denoted by *i*, describes a chromosome with identical chromosome arms at each end; for example, *i*(17q) would consist of two copies of 17q joined at the centromere.

Telomeres

Telomeres are repetitive sequences at the ends of chromosomes. They decrease by approximately 200 base pairs of DNA with every round of replication. When they decrease to a critical length, the cell exits from cell cycle. Germ cells and stem cells, which need to self-renew and maintain a high proliferative potential, contain the enzyme *telomerase* which can

add extensions to the telomeric repeats and compensate for loss at replication and so enable the cells to continue proliferation. Telomerase is also often expressed in malignant cells but this is probably a consequence of the malignant transformation rather than an initiating factor.

Genetic abnormalities associated with haematological malignancies

Some of the genetic abnormalities commonly associated with different types of leukaemia and lymphoma are listed in Table 10.2. The types of gene abnormality include the following (Fig. 10.6).

Point mutation

This is best illustrated by the Val617Phe mutation in the *JAK2* gene which leads to constitutive activation of the JAK2 protein in most cases of myeloproliferative disease (Chapter 19). Mutations within the *RAS* oncogenes or p53 tumour-suppressor gene are common in many haemopoietic malignancies. The point mutation may involve several base pairs. In 35% of cases of AML the *nucleophosmin* gene shows an insertion of 4 base pairs resulting in a frameshift change. Internal tandem duplication or point mutations occur in the *Flt-3* gene in 30% of cases of AML.

Translocations

These are a characteristic feature of haematological malignancies and there are two main mechanisms whereby they may contribute to malignant change (Fig. 10.7).

1 Fusion of parts of two genes to generate a chimeric fusion gene that encodes a novel '*fusion protein*', e.g. *BCR-ABL* in t(9; 22) in chronic myeloid leukaemia (CML) (see Fig. 13.1), *RAR α -PML* in t(15; 17) in AML M₃ (Fig. 10.8) or *TEL-AML1* in t(12; 21) in precursor B-ALL (Fig. 10.9).

2 Overexpression of a normal cellular gene; for example, overexpression of *BCL-2* in the t(14; 18) translocation of follicular lymphoma or of *MYC* in Burkitt's lymphoma (Fig. 10.10). Interestingly, this class of translocation nearly always involves a *TCR* or immunoglobulin gene locus, presumably




Fig. 10.6 Types of genetic abnormality which may lead to haemopoietic malignancy. (a) Point mutation; (b) chromosomal translocation; (c) chromosomal deletion or loss; (d) chromosomal duplication; (e) DNA methylation or deacetylation of histone tails suppresses gene transcription.

as a result of aberrant activity of the recombinase enzyme which is involved in immunoglobulin or *TCR* gene rearrangement in immature B or T cells.

Deletions

Chromosomal deletions may involve a small part of a chromosome, the short or long arm (e.g. 5q-) (Fig. 10.12b) or the entire chromosome (e.g. monosomy 7). Losses most commonly affect chromosomes 5, 6, 7, 11, 20 and Y. The critical event is probably loss of a tumour suppressor gene or of a microRNA as in the 13q14 deletion in CLL (see below).




Fig. 10.7 The two possible mechanisms by which chromosomal translocations can lead to expression of an oncogene.




Fig. 10.8 Generation of the t(15; 17) translocation. The *PML* gene at 15q22 may break at one of three different breakpoint cluster regions (BCR-1, -2 and -3) and joins with exons 3–9 of the *RARα* gene at 17q12. Three different fusion mRNAs are generated (termed long (L), variable (V) or short (S)) and these give rise to fusion proteins of different size. In this diagram only the long version resulting from a break at BCR-1 is shown.

Duplication or amplification

In chromosomal duplication (e.g. trisomy 12 in CLL) or gene amplification, gains are common in chromosomes 8, 12, 19, 21 and Y. Gene amplification is not a common feature in haemopoietic malignancy but has been described involving the *MLL* gene.

Epigenetic alterations

Gene expression may be dysregulated in cancer, not only by structural changes to the genes themselves

but also by alterations in the mechanism by which genes are transcribed. These changes are called *epigenetic* and are stably inherited with each cell division so they are passed on as the malignant cell divides. The most important mechanisms are methylation of cytosine residues in DNA and enzymatic alterations, such as acetylation or methylation, of the histone proteins that package DNA within the cell (Fig. 10.6e). Demethylating agents such as azacytidine increase gene transcription and are finding a role in the management of haemopoietic malignant disease.




Fig. 10.9 Mechanism of action of the core binding factor (CBF) transcription factor and its disruption in acute myeloid leukaemia. CBF consists of two subunits, *CBFβ* and *CBFα* (or *AML1*), which together form a heterodimer (a). This complex binds to the DNA sequence TGTGGT in the regulatory region of certain target genes. This binding allows recruitment of coactivators which lead to transcription from these genes. (b) The t(8;21)

translocation leads to a fusion protein of *CBFα* with *ETO*. Although the CBF subunits can still form heterodimers, their binding to DNA leads to recruitment of a corepressor complex which blocks transcription. (c) In the inv(16) mutation, a *CBFα*-*MYH10* fusion protein is generated which again can form CBF heterodimers but these do not gain access to DNA.




Fig. 10.10 The genetic events in one of the three translocations found in Burkitt's lymphoma and B-cell acute lymphoblastic leukaemia. The oncogene *c-MYC* is normally located on the long arm (q) of chromosome 8. In the (8;14) translocation, *c-MYC* is translocated into close

proximity to the immunoglobulin heavy-chain gene on the long arm of chromosome 14. Part of the heavy-chain gene (the V region) is reciprocally translocated to chromosome 8. C, constant region; *IgH*, immunoglobulin heavy-chain gene; J, joining region; V, variable region.

MicroRNAs

Chromosomal abnormalities, both deletions and amplifications, can result in loss or gain of short (micro) RNA sequences. These are normally transcribed but

not translated. MicroRNAs (miRNAs) control expression of adjacent or distally located genes. Deletion of the miR15a/miR16-1 locus may be relevant to CLL development with the common 13q deletion.

Specific examples of translocations associated with haematological malignancy

Philadelphia (Ph) chromosome

The translocation t(9;22) which underlies Philadelphia chromosome positive chronic myeloid leukaemia (CML) and some cases of ALL is discussed in detail in Chapter 13.

Retinoic acid receptor

In the t(15;17) translocation associated with AML M₃ (p. 158), the promyelocytic leukaemia gene *PML* on chromosome 15 is fused to the retinoic acid receptor α gene, *RAR α* , on chromosome 17 (Fig. 10.8). The resultant PML-RAR α fusion protein functions as a transcriptional repressor whereas normal (wild-type) *RAR α* is an activator. Normally, the PML protein forms homodimers with itself whereas the RAR α protein forms heterodimers with the retinoid X receptor protein, RXR. The PML-RAR α fusion protein binds to PML and RXR, preventing them from linking with their natural partners. This results in the cellular phenotype of arrested differentiation. Cases of AML M₃ associated with the t(15;17) translocation respond to treatment with high doses of all-*trans* retinoic acid (ATRA) which causes differentiation of the abnormal promyelocytes and results in improved prognosis (p. 170). Interestingly, in rare variants of AML M₃, *RAR α* is fused to other genes and in these cases ATRA treatment is not successful.

Translocations involving the core binding factor genes

Core binding factor (CBF) is a heterodimeric transcription factor and is important in regulating expression of a number of genes such as *IL-3* and *GM-CSF*. Genes encoding the two components of CBF, *CBF α* and *CBF β* , are involved in a number of chromosomal translocations associated with leukaemia (Fig. 10.9). These include t(8;21) in which the *CBF α* gene, also known as *AML1*, is translocated to the *ETO* gene on chromosome 8. Another common rearrangement in AML is inv(16) in which the *CBF β*

gene is fused to the *SMMHC* (*MYH11*) gene. In the t(12;21) translocation associated with pre-B-ALL the *TEL* gene is fused to the *CBF α* gene to generate a novel fusion protein. All three translocations appear to act as dominant inhibitors of normal wild-type CBF activity. They are all associated with a relatively good prognosis.

Translocation of *MYC*

In Burkitt's lymphoma and B-ALL one of three translocations is normally found. All of these bring the *MYC* oncogene into close proximity with one of the immunoglobulin genes (Fig. 10.10); the most frequent is translocation to the heavy-chain locus, t(8;14). As a result, expression of the *MYC* gene is deregulated and the gene is expressed in parts of the cell cycle during which it should normally be switched off.

Translocation of the *BCL-2* gene

This oncogene is translocated from chromosome 18 to chromosome 14 in the (14;18) translocation found in approximately 85% of cases of follicular lymphoma and in some cases of diffuse large cell lymphoma. The translocation leads to constitutive expression of the *BCL-2* gene with increased survival of cells because of reduced apoptosis.

Diagnostic methods used to study malignant cells

Karyotype analysis

Karyotype analysis involves direct morphological analysis of chromosomes from tumour cells under the microscope (Fig. 10.11; see Fig. 13.1). This requires tumour cells to be in metaphase and so cells are cultured to encourage cell division prior to chromosomal preparation.

Fluorescent *in situ* hybridization analysis

Fluorescent *in situ* hybridization (FISH) analysis involves the use of fluorescent-labelled genetic probes which hybridize to specific parts of the genome. It is possible to label each chromosome with a different




Fig. 10.11 A colour-banded karyotype from a normal male. Each chromosome pair shows an individual colour-banding pattern. This involves a cross-species multiple colour chromosome banding technique. Probe sets developed from the chromosomes of gibbons are combinatorially labelled and hybridized to human

chromosomes. The success of cross-species colour banding depends on a close homology between host and human conserved DNA, divergence of repetitive DNA and a high degree of chromosomal rearrangement in the host relative to the human karyotype. (Courtesy of Dr C.J. Harrison)

combination of fluorescent labels (Fig. 10.12a). This is a sensitive technique that can detect extra copies of genetic material in both metaphase and interphase (non-dividing) cells (e.g. trisomy 12 in CLL) or, by using two different probes, reveal chromosomal translocations (Fig. 10.12b); for example, t(9;22) in CML (Fig. 13.1e) or reduced chromosome numbers or segments (e.g. monosomy 7 or 5 in myelodysplasia) (Fig. 10.12c).


Southern blot analysis

Southern blot analysis involves the extraction of DNA from leukaemic cells followed by restriction enzyme digestion, gel electrophoresis and transfer by 'blotting' to a suitable membrane. The DNA is then hybridized to a probe complementary to the gene of interest. When the probe recognizes a segment within the boundaries of a single restriction

fragment one band is identified but if the gene has been translocated to a new area in the genome a novel band of different electrophoretic mobility is seen. Although time consuming and relatively insensitive, this remains a powerful technique.


Polymerase chain reaction

Polymerase chain reaction (PCR) (see Fig. 6.23) can be performed on blood or bone marrow for a number of specific translocations such as t(9;22) and t(15;17) (Fig. 10.13). It can also be used to detect 'clonal' cells of B- or T-cell lineage by immunoglobulin or T-cell receptor (TCR) gene rearrangement analysis. As it is relatively straightforward and extremely sensitive (detecting one abnormal cell in 10^5 – 10^6 normal cells), it has become of great value in the diagnosis and monitoring of minimal residual disease (p. 144).



Missing red FISH signal (arrow) shows deletion of the long arm of chromosome 5, while the normal homologue 5 is marked by red & green signals

(b)



(c)

Fig. 10.12 (a) The principles of fluorescent *in situ* hybridization (FISH). A particular strength is that it may be performed on both dividing and non-dividing (quiescent) cells. (b) Detection of long arm (q) of chromosome 5. (Courtesy of Dr Ellie Nacheva) (c) An example of FISH analysis showing the t(12;21) translocation. The green probe hybridizes to the region of the *TEL* gene on chromosome 12 and the red probe hybridizes to the region of the *AML1* gene on chromosome 21. The arrows point to the two derived chromosomes resulting from the reciprocal translocation. (Courtesy of Dr C.J. Harrison)

DNA microarray platforms

DNA microarrays allow a rapid and comprehensive analysis of cellular transcription by hybridizing labelled cellular mRNA to DNA probes which are immobilized on a solid support (Fig. 10.14a).

Oligonucleotides or complementary DNA (cDNA) arrays may be immobilized on the array and RNA from the tissue of interest is used to generate fluorescent cDNA or RNA which is then annealed to the nucleic acid matrix. This approach can rapidly determine mRNA expression from a large number



Fig. 10.13 Reverse transcription polymerase chain reaction (PCR) analysis of bone marrow from a patient with AML M_3 (acute promyelocytic leukaemia) at diagnosis. The PML-RAR α fusion product (cDNA) formed by the t(15; 17) translocation has been amplified by PCR using oligonucleotide primers from the PML and RAR α genes. Lane 1, water control; lane 2, low molecular weight DNA marker; lane 3, patient sample. A single 355 base pair fusion message has been amplified showing the presence of the fusion gene.

of genes and may be used to determine the mRNA expression pattern of different leukaemia or lymphoma subtypes (Fig. 10.14b). The technique cannot be used to detect (minimal) residual disease and is not yet in routine diagnostic use but it is clear that it can give valuable information for subclassification of large cell non-Hodgkin's lymphoma diffuse (Fig. 17.5), or AML without cytogenetic changes.


Flow cytometry

In this technique, antibodies labelled with different fluorochromes recognize the pattern and intensity of expression of different antigens on the surface of normal and leukaemic cells (see Appendix 1). Normal cells each have a characteristic profile but malignant cells often express an aberrant phenotype that can be useful in allowing their detection. In the case of B cell malignancies (e.g. CLL), expression of

only one light chain, κ or λ , by the tumour cells distinguishes them from a normal polyclonal population which express both κ and λ chains, usually in a κ : λ ratio of 2:1 (Fig. 10.15).

Immunohistology


Antibodies can also be used to stain tissue sections with fluorescent markers and this is known as *immunohistology* or *immunohistochemistry*. The presence and architecture of tumour cells can be identified by visualization of stained tissue sections under the microscope. The clonal nature of B-cell malignancies can be shown in tissue sections by staining for κ or λ chains. A malignant clonal population (e.g. in B-cell NHL) will express one or other light chain but not both (Fig. 17.4).



(a)

Fig. 10.14 (a) Principle of transcriptional profiling of RNA expression in leukaemia samples using DNA microarrays.

(Continued)



(b)

Fig. 10.14 (Continued) (b) Microarray analysis of genes distinguishing acute lymphoblastic leukaemia (ALL) from acute myeloid leukaemia (AML). The 50 genes most highly correlated on gene-expression microarrays with each of these leukaemias are shown. Each row corresponds to a gene; each column corresponds to the expression value in a particular sample. Expression for each gene is normalized across the samples such that the

mean is 0 and the SD is 1. Expression greater than the mean is shaded in red, and that below the mean is shaded in blue. Although the genes as a group appear correlated with the type of leukaemia under study, no single gene is uniformly expressed across the class, illustrating the value of a multigene prediction method. (Reproduced courtesy of Golub and colleagues)




Fig. 10.15 Flow cytometric analysis of B-cell chronic lymphocytic leukaemia. The five plots show the tumour to have a phenotype of: (a) CD5⁺ CD19⁺; (b) FMC7⁻ CD2⁻; (c) CD20⁺ CD79b^{low}; (d) CD23⁺ and (e) $\lambda^+ \kappa^-$.

Value of genetic markers in management of haematological malignancy

The detection of genetic abnormalities may be important in several aspects of the management of patients with leukaemia or lymphoma.

Initial diagnosis

Many genetic abnormalities are so specific for a particular disease that their presence determines that diagnosis. An example is the t(15; 17) translocation

which classifies an AML as promyelocytic leukaemia. Clonal immunoglobulin or TCR gene rearrangements are useful in establishing clonality and determining the lineage of a lymphoid malignancy.

For establishing a treatment protocol

Each major type of haematological malignancy can be further subdivided on the basis of detailed genetic information. For instance, AML is a diverse group of disorders with characteristic genotypes. Individual subtypes respond differently to standard treatment.




Fig. 10.16 Sensitivity of detection of leukaemic cells using five different techniques. 10^1 to 10^6 = 1 cell in 10 to 1 cell in 10^6 detected.

The t(15; 17), inv(16) and t(8; 21) subgroups do well, whereas monosomy 7 carries a poor prognosis. Treatment strategies are now tailored for the individual and in some instances knowledge of the underlying genetic abnormality can lead to more rational treatment. The best examples are the use of retinoic acid in the treatment of acute promyelocytic leukaemia associated with the t(15; 17) translocation and the use of imatinib for treatment of Ph+ CML. Genetic information is also valuable for giving a prognosis. For instance, Ph+ ALL has a particularly poor prognosis, whereas hyperdiploidy in ALL is a favourable finding. DNA microarrays are valuable for prognosis (e.g. in diffuse large B-cell lymphoma) which can be divided into good and bad prognostic groups according to the pattern of gene expression (Fig. 17.5).

Monitoring the response to therapy

The detection of minimal residual disease (MRD) (disease that cannot be seen by conventional microscopy of the blood or bone marrow) in AML, ALL or CML when the patient is in remission after chemotherapy or stem cell transplantation is possible using the following techniques (in increasing order of sensitivity; Fig. 10.16).




Fig. 10.17 Detection of minimal residual disease (MRD) by four-colour flow cytometry in: normal bone marrow mononuclear cells (BM), BM from a patient with B lineage ALL at diagnosis and in remission 6 weeks after diagnosis. The cells were detected with four different antibodies (anti-CD10, anti-CD19, anti-CD34, anti-CD38) attached to fluorescent labels abbreviated as PE, APC, PerCP and FITC, respectively. The tridimensional plot shows the immunophenotype of CD19⁺ lymphoid cells in the three samples. MRD of 0.03% of cells expressing the leukaemia-associated phenotype (CD10⁺, CD34⁺, CD38⁺) were detected at 6 weeks, confirmed by PCR analysis. (From D. Campana and E. Coustan-Smith (1999). Cytometry. *Commun Clin Cytometry* 38,139–52, with permission)




Fig. 10.18 Real-time quantitative PCR in acute B-lineage lymphoblastic leukaemia for minimal residual disease using the immunoglobulin heavy chain as target. Primers are designed based on DNA from sequence analysis of the presenting leukaemic clone. Bone marrow samples taken in clinical remission are amplified by PCR using these primers and fluorescent labelled using Sybergreen. The intensity of the signal measures the total DNA


molecules amplified in successive cycles. In this example, the intensities of amplification of DNA from two follow-up bone marrow samples (FU1 and FU2) are compared with serial deletions of (10^{-1} to 10^{-4}) of the DNA from the presentation bone marrow. FU1 shows a level of residual disease approximately 1 in 5000 (0.02%) and FU2 of 1 in 12 000 (0.008%). (Courtesy of Dr L. Foroni)

- 1 Cytogenetic analysis.
- 2 Southern blot analysis to look for a tumour-specific DNA rearrangement.
- 3 Fluorescence-activated cell sorting (FACS) to detect tumour cells using immunological markers that detect 'leukaemia-specific' combinations of antigens (Fig. 10.17).
- 4 PCR to amplify tumour-specific translocations

or immunoglobulin/TCR sequences specific to the original clone (Fig. 10.18).

These approaches are being evaluated but they already have an important role in determining the treatment of individual patients (e.g. persistence of MRD in childhood ALL after the initial 1–3 months of therapy predicts probable relapse; Fig. 10.19) while persistence of the *BCR-ABL* fusion gene after

Fig. 10.19 Cumulative incidence of relapse according to minimal residual disease (MRD) levels at the end of remission induction in children with acute lymphoblastic leukaemia (ALL) treated at St Jude Children's Research Hospital. (Courtesy of Dr D. Campana)



allogeneic stem cell transplantation for CML suggests the need for early treatment with donor leucocytes (p. 262).

Bibliography

- Abramson J.S. and Shipp M.A. (2005) Advances in the biology and therapy of diffuse large B-cell lymphoma: moving toward a molecularly targeted approach. *Blood* 106,1164–74.
- Bullinger L. (2006) Gene expression profiling in acute myeloid leukemia. *Haematologica* 91,733–8.
- Chou W.C., Tang J.L., Lin L.I. *et al.* (2006) Nucleophosmin mutations in *de novo* acute myeloid leukemia: the age-dependent incidences and the stability during disease evolution. *Cancer Res* 66,3310–6.
- Foroni, L., Gameiro, P.M., Hoffbrand, A.V. (2005) Minimal residual disease in acute leukaemia. In: Hoffbrand A.V., Catovsky D., Tuddenham E.G.D. (eds) *Postgraduate Haematology*, 5th edn. Blackwell Science, Oxford: pp. 561–74.
- Gilliland D.G. and Griffin J.D. (2002) The roles of FLT3 in hematopoiesis and leukemia. *Blood* 100,1532–42.
- Haferlach T., Kohlmann A., Schnittger S. *et al.* (2005) General approach to the diagnosis of leukemia using gene expression profiling. *Blood* 106,1189–98.
- Hubank M. (2004) Gene expression profiling and its application in studies of haematological malignancy. *Br J Haematol* 124,577–94.
- Kottaridis P.D., Gale R.F. and Linch D.C. (2003) FLT3 mutations and leukaemia. *Br J Haematol* 122,523–38.
- Krause D.S. and Van Etten R.A. (2005) Mechanisms of disease: tyrosine kinases as targets for cancer therapy. *N Engl J Med* 353,172–87.
- Lossos I.S., Czerwinski D.K., Alizadeh A.A. *et al.* (2004) Prediction of survival in diffuse large-B-cell lymphoma based on the expression of six genes. *N Engl J Med* 350,1828–37.
- Lotem J. and Sachs L. (2006) Epigenetics and the plasticity of differentiation in normal and stem cancer cells. *Oncogene*: advanced on line publication, doi:10.1038/sj.onc.1209816.
- Speck N.A. and Gilliland D.G. (2002) Core-binding factors in hematopoiesis and leukaemia. *Nat Rev Cancer* 2,502–13.
- Staudt L.M. and Dave S. (2005) The biology of human lymphoid malignancies revealed by gene expression profiling. *Adv Immunol* 87,163–208.
- Wiemels J.L., Ford A.M., Van Wering E.R., Postma A. and Greaves M. (1999) Protracted latency of acute lymphoblastic leukaemia after TEL-AML gene fusion *in utero*. *Blood* 94,1057–62.

CHAPTER 11

11

Management of haematological malignancy

General support therapy, 147

Specific therapies for haematological malignancy, 151

Bibliography, 156

The treatment of haematological malignancy has improved greatly over the last 40 years. This has resulted from developments in *support therapy* and in *specific treatment*. Details of specific treatment are discussed in relation to individual diseases in the appropriate chapter. Support care and general aspects of the agents used in the treatment of haematological malignancy are described here.

General support therapy

Patients with haematological malignancies often present with medical problems related to suppression of normal haemopoiesis and this problem is compounded by the treatments that are given to eradicate the tumour. General supportive therapy for bone marrow failure includes the following.

1 Insertion of a central venous catheter A central venous catheter is usually inserted prior to intensive treatment via a skin tunnel from the chest into the superior vena cava (Fig. 11.1). This gives ease of access for administering chemotherapy, blood products, antibiotics and intravenous feeding. In addition, blood may be taken for laboratory tests.

2 Blood product support (Chapter 27). Red cell and platelet transfusions are used to treat anaemia and thrombocytopenia. A number of particular issues apply to the support of patients with haematological malignancy:

- (i) The threshold haemoglobin for transfusion will depend on clinical factors such as symptoms and speed of onset of anaemia but most units give

red cell support for a Hb <8 g/dL, with a higher threshold in older patients. In patients needing both red cells and platelets, platelets are given first to reduce the risk of a further fall in the platelet count.

- (ii) The trigger for platelet transfusion is typically a platelet count $<10 \times 10^9/L$ but this should be doubled in the presence of active bleeding or infection.

- (iii) Fresh frozen plasma (FFP) may be needed to reverse coagulation defects.

- (iv) Cytomegalovirus (CMV) negative blood should be given to all patients until it has been shown that they are either CMV seropositive or that they will never be candidates for stem cell transplantation. This is to prevent transmission of CMV to uninfected patients as the virus is a significant problem in stem cell transplant recipients (p. 258).

- (v) Red cell transfusions should be avoided if at all possible in patients with a very high white cell count ($>100 \times 10^9/L$) because of the hyperviscosity and the risk of precipitating thrombotic episodes as a result of white cell stasis.

- (vi) Large volume transfusions, such as 3 units of blood or more, can precipitate pulmonary oedema in older patients and should be given slowly and with clinical monitoring. Diuretics such as frusemide (furosemide) are often given.

- (vii) Febrile reactions with blood products are not uncommon and should be managed by slowing the infusion and administration of drugs such as




Fig. 11.1 (a) A central venous line in a patient undergoing intensive chemotherapy. (b) Chest X-ray showing correct placement of a central venous line. (Courtesy of Dr D. Tattersall)

antihistamines, pethidine or hydrocortisone. The dosage of steroids should be limited because of concerns with immunosuppression.

(viii) Blood products given to highly immunosuppressed patients (e.g. from chemotherapy, such as fludarabine, or with aplastic anaemia or Hodgkin's disease) should be irradiated prior to administration to prevent graft-versus-host disease (p. 256).

(ix) The use of recombinant erythropoietin to reduce the need for blood transfusion and improve patient well being (e.g. in myeloma or myelodysplasia) is discussed on p. 14.

3 Haemostatic support A coagulation screen should be performed regularly on patients undergoing intensive chemotherapy and support with vitamin K or FFP may be required. Cryoprecipitate may be needed for fibrinogen deficiency (e.g. precipitated by asparaginase in the management of acute lymphoblastic leukaemia (ALL)). Antiplatelet drugs such as aspirin or clopidogrel are usually discontinued in patients undergoing intensive chemotherapy and patients on long-term warfarin can be switched to low molecular weight heparin, which can then itself be stopped if the platelet count falls below $50 \times 10^9/L$. Progesterones are given to premenopausal women undergoing intensive chemotherapy to pre-

vent menstruation. Tranexamic acid can be given to reduce haemorrhage in patients with chronic low-grade blood loss.

4 Antiemetic therapy Nausea and vomiting are common side-effects of chemotherapy. A key objective is to try to prevent nausea occurring early in the treatment as it is more difficult to control once problems have arisen. The 5-HT₃ (serotonin) receptor antagonists such as ondansetron or granisetron can control nausea from intensive chemotherapy in over 60% of cases and the addition of dexamethasone can increase this by approximately 20%. Metoclopramide, prochlorperazine or cyclizine, benzodiazepines (e.g. lorazepam), domperidone or cannabinoids (e.g. nabilone) can all have a role.

5 Tumour lysis syndrome Chemotherapy may trigger an acute rise in plasma uric acid, potassium and phosphate and cause hypocalcaemia because of rapid lysis of tumour cells. This syndrome is seen most commonly with rapidly dividing tumours such as lymphoblastic lymphoma or acute leukaemia and can cause acute renal failure. Allopurinol, intravenous fluids and electrolyte replacement are the mainstay of prevention and alkalization of the urine is sometimes used. Rasburicase, the oxidation of uric acid to allantoin, is highly effective in controlling hyperuricaemia.

6 Psychological support Patients with a diagnosis of malignant disease commonly feel concerns about such issues as the discomfort of treatment, finance, sexuality and fear of mortality. Even when patients achieve a clinical remission there is understandable concern about the chance of disease relapse. Psychological support should be an integral part of the relationship between physician and patient, and patients should be allowed to express their fears and concerns at the earliest opportunity. Most patients value the opportunity to read more about their disorder and many excellent booklets or websites are now available. Teamwork is also crucial and the nursing staff and trained counsellors have a vital role in offering support and information during inpatient and outpatient care. Many units have specialist input from clinical psychologists and psychiatric help may occasionally be required. Inadequate communication is perhaps the most common failing of medical teams. The immediate family should be kept informed of the patient's progress whenever possible and appropriate.

7 Reproductive issues Men who are to receive cytotoxic drugs should be offered sperm storage, ideally before treatment commences or, if impossible, within a short period of time thereafter. Ethical issues relating to storage or potential usage of tissue in the event of treatment failure will need to be addressed. Permanent infertility in women is less common after chemotherapy although premature menopause may occur. Storage of fertilized ova is usually impractical and storage of unfertilized ova is currently very difficult and, despite some recent progress, is not offered as a routine service.

8 Nutritional support Some degree of weight loss is virtually inevitable in patients undergoing inpatient chemotherapy because of the combination of a poor nutritional intake, malabsorption caused by drugs and a catabolic disease state. If a weight loss of >10% occurs, support with total nutrition is often given, either enterally via a nasogastric tube or parenterally through a central venous catheter.

9 Pain is rarely a major problem in haematological malignancies except myeloma although bone pain can be a presenting feature. The mucositis that follows intensive chemotherapy can cause severe discomfort and continuous infusions of opiate analgesia are often required. Pain is often a considerable issue in

patients with multiple myeloma and can be managed by a combination of analgesia and chemotherapy/radiotherapy. Advice from palliative care teams or specialist pain management practitioners should be sought when required.

10 Prophylaxis and treatment of infection Patients with haematological malignancy are at great risk of infection which remains the major cause of morbidity and mortality. Immunosuppression may result from neutropenia, hypogammaglobulinaemia and impaired cellular function. These can be secondary to the primary disease or its treatment. Neutropenia is a particular concern and in many patients neutrophils are totally absent from the blood for periods of 2 weeks or more. The use of granulocyte colony-stimulating factor (G-CSF) to reduce periods of neutropenia is discussed on p. 97. The likely sequence of infection following stem cell transplantation is illustrated in Fig. 11.2.

Bacterial infection

This is the most common problem and usually arises from the patient's own commensal bacterial flora. Gram-positive skin organisms (e.g. *Staphylococcus* and *Streptococcus*) commonly colonize central venous lines, whereas Gram-negative gut bacteria (e.g. *Pseudomonas aeruginosa*, *Escherichia coli*, *Proteus*, *Klebsiella* and anaerobes) can cause overwhelming septicaemia. Even organisms not normally considered pathogenic, such as *Staphylococcus epidermidis*, may cause life-threatening infection. In the absence of neutrophils, local superficial lesions can rapidly cause severe septicaemia.

Prophylaxis of bacterial infection

Protocols used to limit bacterial infection vary from unit to unit and may include the use of a prophylactic antibiotic such as ciprofloxacin. During periods of neutropenia, topical antiseptics for bathing and chlorhexidine mouthwashes and a 'clean diet' are recommended. The patient is nursed in a reverse-barrier room. The severity and length of mucositis may be reduced by treatment with recombinant human keratinocyte growth factor (Palifermin) which reduces the severity of oral mucositis. Oral non-absorbed antimicrobial agents such as neomycin and colistin reduce gut commensal flora but




Fig. 11.2 A protocol for the management of fever in the neutropenic patient. CRP, C-reactive protein; CXR, chest X-ray.

their value is unclear. Regular surveillance cultures are taken to document the patient's bacterial flora and its sensitivity.

Treatment of bacterial infection

Fever is the main indication that infection is present because if neutropenia is present pus will not be formed and infections are often not localized. Fever may be caused by blood products or drugs, but infection is the most common cause and fever of over 38°C in neutropenic patients should be investigated and treated within hours. Cultures should be taken from any likely focus of infection including blood from central venous lines and peripheral veins, from urine and mouth swabs. The mouth and throat, intravenous catheter site, and perineal and perianal areas are particularly likely foci. A chest X-ray is indicated as chest infections are frequent.

Antibiotic therapy must be started immediately after blood and other cultures have been taken; in many febrile episodes no organisms are isolated.

There are many different antibiotic regimes in use and a close link with the microbiology team

is essential. A typical regimen might be based on a single agent such as a broad-spectrum penicillin (e.g. Tazocin), meropenem or a broad-spectrum cephalosporin. An aminoglycoside such as gentamicin or vancomycin is often added. *Staphylococcus epidermidis* is a common source of fever in patients with intravenous lines and an agent such as teicoplanin, vancomycin or linezolid may be needed. If an infective agent and its antibiotic sensitivities become known, appropriate changes in the regimen are made. If no response occurs within 48–72 h, changing the antibiotics or treating a fungal or viral infection are considered.

Viral infection

Prophylaxis and treatment of viral infection

Herpes viruses, such as herpes simplex, varicella zoster, CMV and Epstein-Barr virus (EBV), undergo latency following primary infection and are never eradicated from the host. Most patients with haematological malignancy have already been infected with these agents and viral reactivation is therefore

the most common problem. Aciclovir is frequently given prophylactically. Herpes simplex is a common cause of oral ulcers but is usually controlled easily by aciclovir. Varicella zoster frequently reactivates in patients with lymphoproliferative diseases to cause shingles which requires treatment with high doses of aciclovir or valaciclovir. Primary infection, usually in children, can be very serious and immunoglobulin can be used to prevent infection following recent exposure. Reactivation of CMV infection is particularly important following stem cell transplantation (Chapter 21) but may occur following intensive chemotherapy. Failure of immune control of EBV following allogeneic transplantation can lead to outgrowth of a B-cell tumour known as post-transplant lymphoproliferative disease (PTLD).

Fungal infection

Prophylaxis and treatment of fungal infection

Because of the intensity of current chemotherapy, fungal infections are a major cause of morbidity and mortality. The two major subtypes are yeasts such as *Candida species* and moulds of which *Aspergillus fumigatus* is the most common.

Invasive aspergillosis is now the most common cause of infectious death in intensively immunocompromised patients. Infection occurs through inhalation of *Aspergillus* spores (conidia) (Fig. 11.3) and air filtration systems are used in many haematology wards. The diagnosis of invasive aspergillosis can be difficult. Definitive diagnosis requires demonstration of invasive growth on a biopsy specimen but such evidence is rarely available. Polymerase chain reaction (PCR) for fungal DNA or enzyme-linked immunosorbent assay (ELISA) for *Aspergillus* galactomannan are often used and the detection of β 1-3 D-glucan is also showing promise. High resolution computed tomography (HRCT) scan is a valuable tool and early features are nodular lesions with a 'ground glass' appearance. Later on, wedge lesions, the halo sign and the air crescent sign are seen (Fig. 11.4). A high index of suspicion for fungal infection should be maintained and treatment is often started empirically for a fever that has failed to resolve after 3–4 days of antibiotic treatment.

Treatment of established *Aspergillus* infection is



Fig. 11.3 Sporing heads of *Aspergillus fumigatus*. (Courtesy of Dr Elizabeth Johnson)


with amphotericin (usually liposomal), voriconazole or caspofungin. Surgery to remove lung lesions may be needed.

Candida species are a common hospital pathogen and frequently cause oral infection. *Candida* can be significant when isolated from normally sterile body fluids such as blood or urine. Prophylaxis or treatment is usually with fluconazole or itraconazole.

Pneumocystis carinii (also known as *Pneumocystis jiroveci*) is an important cause of pneumonitis. Prophylaxis with co-trimoxazole or nebulized pentamidine is highly effective and is given to those who have received intensive (combination) chemotherapy. Treatment is with high dose co-trimoxazole.

Specific therapies for haematological malignancy


Specific therapy is aimed at reducing the tumour cell burden by the use of drugs or radiotherapy. The hope in some diseases is to eradicate the tumour completely and cure rates for haematological malignancy are gradually improving. However, cure is often not achievable so palliation can also be an important aim.




(a)



(b)



(c)



(d)

Fig. 11.4 (a) Chest X-ray of patient with pulmonary aspergillosis which shows an area of cavitation containing a central fungal ball (arrow) leading to the typical 'air-crescent' sign. Hazy ground-glass shadowing with bronchiolar dilatation (b) and (c) nodules are seen in early aspergillosis, whereas a fungal ball with surrounding air is typical of more advanced disease (d).

A wide variety of drugs are used in the management of haemopoietic malignancies and several drugs are often combined together in regimens that minimize the potential for resistance to occur against a single agent. Many act specifically on dividing cells and their selectivity is dependent on the high proliferation rate within the tumour. Not all tumour cells will be killed by a single course of treatment and it is usual to give several courses of treatment which gradually eradicate the tumour burden. This 'log kill' hypothesis also gives the residual normal haemopoietic cells the opportunity to recover between treatment courses.

Drugs used in the treatment of haemopoietic malignancies

Cytotoxic drugs (Table 11.1)

Alkylating agents such as chlorambucil, cyclophosphamide and melphalan are activated to expose reactive alkyl groups which make covalent bonds to molecules within the cell. These have a particular affinity for purines and are thus able to cross-link DNA strands and impair DNA replication, resulting in a block at G₂ (Fig. 1.8) and death of the cell by apoptosis (Fig. 1.11).

Table 11.1 Drugs used in the treatment of leukaemia and lymphoma.

	Mechanism of action	Particular side-effects*
<i>Alkylating agents</i>		
Cyclophosphamide	Cross-link DNA, impede RNA formation	Haemorrhagic cystitis, cardiomyopathy, loss of hair
Chlorambucil		Marrow aplasia, hepatic toxicity, dermatitis
Busulfan		Marrow aplasia, pulmonary fibrosis, hyperpigmentation
Melphalan		Marrow aplasia
Nitrosoureas BCNU, CCNU		Renal and pulmonary toxicity
Cisplatin	Intrastrand DNA linkage	Renal dysfunction, neurotoxicity, ototoxicity
<i>Antimetabolites</i>		
Methotrexate	Inhibit pyrimidine or purine synthesis or incorporation into DNA	Mouth ulcers, gut toxicity
6-Mercaptopurine [†]		Jaundice
6-Tioguanine [†]		Gut toxicity
Cytosine arabinoside		CNS especially cerebellar toxicity and conjunctivitis at high doses
<i>Cytotoxic antibiotics</i>		
Anthracyclines (e.g. daunorubicin)	Bind to DNA and interfere with mitosis	Cardiac toxicity, hair loss
Hydroxodaunorubicin (Adriamycin)		
Mitoxantrone		
Idarubicin		
Bleomycin		
	DNA breaks	Pulmonary fibrosis, skin pigmentation
<i>Plant derivatives</i>		
Vincristine (Oncovin)	Spindle damage	Neuropathy (peripheral or bladder or gut)
Vinblastine		
Vindesine	Mitotic inhibitor	Hair loss, oral ulceration
Epipodophyllotoxin (etoposide, VP-16)		
<i>Purine analogues</i>		
Fludarabine	Inhibits adenosine deaminase or other purine pathways	Immunosuppression (low CD4 counts); renal and neurotoxicity (at high doses)
2-Chlorodeoxyadenosine		
Deoxycoformycin		
<i>Miscellaneous</i>		
Hydroxyurea	Inhibits ribonucleotide reductase	Pigmentation, nail dystrophy, skin ulceration
Signal transduction inhibitors (imatinib)	Inhibits tyrosine kinase activity	Myelosuppression, fluid retention
Corticosteroids	Lymphoblast lysis	Peptic ulcer, obesity, diabetes, osteoporosis, psychosis, hypertension
Trans-retinoic acid	Induces differentiation	Liver dysfunction, skin hyperkeratosis, leucocytosis and hyperviscosity, pleural or pericardial effusion

(Continued on p. 154)

Table 11.1 (Continued) Drugs used in the treatment of leukaemia and lymphoma.

	Mechanism of action	Particular side-effects*
Arsenic	Induces differentiation or apoptosis	Hyperleucocytosis, cardiac
α -Interferon	Activation of RNAase and natural killer activity	Flu-like symptoms, thrombocytopenia, leucopenia, weight loss
<i>Monoclonal antibodies</i>		
Rituximab (anti-CD20)	Induction of apoptosis	Infusion reactions, immunosuppression
CamPATH (anti-CD52)	Lysis of target cell by complement fixation	
Ibritumomab (Zevalin) (anti-CD20 ⁺ ⁹⁰ Y)	Toxicity to bound cell	Myelosuppression, nausea
Tositumomab (Bexxar) (anti-CD20 ⁺ ¹³¹ I)	Toxicity to bound cell	Myelosuppression
L-Asparaginase	Deprive cells of asparagine	Hypersensitivity, low albumin and coagulation factors, pancreatitis

* Most of the drugs cause nausea, vomiting, mucositis and bone marrow toxicity, and in large doses infertility.

Tissue necrosis is a problem if the drugs are extravasated during infusion.

† Allopurinol potentiates the action and side-effects of 6-mercaptopurine.

Antimetabolites block metabolic pathways used in DNA synthesis. There are three major groups:

1 **Folate antagonists**, such as methotrexate (Fig. 4.5). Methotrexate is widely used alone or in combination with cytosine arabinoside as intrathecal prophylaxis of CNS disease in patients with ALL, AML or high grade non-Hodgkin lymphomas. High systemic doses may also penetrate the CNS. Folinic acid (formyl THF) is able to overcome the activity of methotrexate and is sometimes administered to 'rescue' normal cells after high-dose methotrexate therapy.

2 **Pyrimidine analogues** include cytosine arabinoside (cytarabine; ara-C) which is an analogue of 2'-deoxycytidine and is incorporated into DNA where it inhibits DNA polymerase and blocks replication.

3 **Purine analogues** include fludarabine (which inhibits DNA synthesis in a manner similar to ara-C), mercaptopurine, azathioprine and deoxycoformycin.

Cytotoxic antibiotic drugs include the anthracyclines such as doxorubicin, hydroxodaunorubicin, epirubicin and mitoxantrone. These are able to intercalate into DNA and then bind strongly to enzymes called topoisomerases. These enzymes are critical for relieving torsional stress in replicating

DNA by nicking and resealing DNA strands. If their activity is blocked, DNA replication cannot take place.

Bleomycin is a metal chelating antibiotic that generates superoxide radicals within cells that degrade preformed DNA. It is active on non-cycling cells.

Plant derivatives include the vinca alkaloids such as vincristine which is derived from the periwinkle plant. It binds to tubulin and prevents its polymerization to microtubules. This blocks cell division in metaphase (Fig. 1.8). Etoposide inhibits topoisomerase action.

Other agents

Hydroxyurea (hydroxycarbamide) is used widely in the treatment of myeloproliferative disorders. It inhibits the enzyme ribonucleotide reductase which converts ribonucleotides to deoxyribonucleotides. It is not thought to damage DNA and is used in non-malignant disorders such as sickle cell anaemia (p. 89).

Imatinib is a new class of drug that binds to the bcr-abl fusion protein generated from the Philadelphia chromosome translocation in chronic




Fig. 11.5 The site of action of drugs used in the management of haemopoietic malignancies. ATRA, all-trans retinoic acid.

myeloid leukaemia (CML; p. 177). It blocks binding of adenosine triphosphate (ATP) and thus prevents phosphorylation of substrate proteins leading to apoptosis of the cell (Fig. 13.4).

Corticosteroids have a potent lymphocytotoxic activity and have an important role in many chemotherapeutic regimens used in the treatment of lymphoid malignancy and myeloma.

All-trans retinoic acid (ATRA) is a vitamin A derivative that acts as a differentiation agent in acute promyelocytic leukaemia (APML). Tumour cells in APML are arrested at the promyelocyte stage as a

result of transcriptional repression resulting from the PML-RARA fusion protein (p. 136). ATRA relieves this block and may lead to a brisk neutrophilia within a few days of treatment with other side-effects known as the 'ATRA syndrome' (p. 170).

Demethylation agents (e.g. azacytidine) act to increase transcription by reducing methylation on cytosine residues within DNA.

Interferon- α is an antiviral and antimitotic substance produced in response to viral infection and inflammation. It has proven useful in CML, myeloma and myeloproliferative diseases.

Monoclonal antibodies are highly effective against B-cell malignancies. Rituximab binds to CD20 on B cells and appears to mediate cell death, primarily through direct induction of apoptosis and opsonization (p. 212). Alemtuzumab (Campath) binds to CD52 and is highly efficient at fixing complement which lyses the target B and T cells. Antibodies may also carry attached toxins (e.g. Mylotarg) (anti-CD33) or radioactive isotopes (e.g. Zevalin or Bexaar).

Asparaginase is an enzyme derived from bacteria which breaks down the amino acid asparagine within the circulation. ALL cells lack asparagine synthase and thus need a supply of exogenous asparagine for protein synthesis. Intramuscular asparaginase is an important agent in the treatment of ALL, although hypersensitivity reactions are not uncommon and blood clotting may be disturbed.

Platinum derivatives (e.g. cisplatin) are used in combinations for treating lymphoma.

Arsenic is useful in treatment of relapsed AML M₃. It induces differentiation and apoptosis.

Bibliography

- Cairo M.S. and Bishop M. (2004) Tumour lysis syndrome: new therapeutic strategies and classification. *Br J Haematol* 127,3–11.
- Donnelly J.P. (2001) Prophylaxis of infections. In: Rolston K.V.I. and Rubenstein E.B. (eds) *Textbook of Febrile Neutropenia*. Martin Dunitz, London: pp. 215–43.
- Furno P., Bucaneve G., Del Favero A. (2002) Monotherapy or aminoglycoside-containing combinations for empirical antibiotic treatment of febrile neutropenic patients: a meta-analysis. *Lancet Infect Dis* 2,231–42.
- Glasmach A., Prentice A., Gorschluter M. *et al.* (2003) Itraconazole prevents invasive fungal infections in neutropenic patients treated for hematologic malignancies: evidence from a meta-analysis of 3597 patients. *J Clin Oncol* 21,4615–26.
- Prentice A.G. and Donnelly J.P. (2005) Supportive care in the management of leukaemia. In: Hoffbrand A.V., Catovsky D., Tuddenham E.G.D. (eds) *Postgraduate Haematology*, 5th edn. Blackwell Publishing, Oxford: pp. 586–602.
- Rebulla P., Finazza G., Marangoni F. *et al.* (1997) The threshold for prophylactic platelet transfusions in adults with acute myeloid leukemia. The Gruppo Italiano Malattie Ematologiche Maligne dell'Adulto. *N Engl J Med* 337,1870–5.
- Shoham S. and Levitz S.M. (2005) The immune response to fungal infections. *Br J Haematol* 129,569–82.
- Spielberger R., Stiff P., Bensinger W. *et al.* (2004) Palifermin for oral mucositis after intensive therapy for hematologic cancers. *N Engl J Med* 351,2590–8.
- Stewart A.F. (2005) Hypercalcemia associated with cancer. *N Engl J Med* 352,373–9.
- Swanton C. (2004) Cell-cycle targeted therapies. *Lancet Oncol* 5,27–34.
- Walsh T.J., Teppler H., Donowitz G.R. *et al.* (2004) Caspofungin versus liposomal amphotericin B for empirical antifungal therapy in patients with persistent fever and neutropenia. *N Engl J Med* 351,1391–402.

Classification of leukaemia, 157

Classification and pathogenesis of acute leukaemia, 157

Acute lymphoblastic leukaemia, 158

Acute myeloid leukaemia, 167

Bibliography, 172

The leukaemias are a group of disorders characterized by the accumulation of malignant white cells in the bone marrow and blood. These abnormal cells cause symptoms because of: (i) bone marrow failure (i.e. anaemia, neutropenia, thrombocytopenia) and (ii) infiltration of organs (e.g. liver, spleen, lymph nodes, meninges, brain, skin or testes).

Classification of leukaemia

The main classification is into four types: acute and chronic leukaemias, which are further subdivided into lymphoid or myeloid (Table 12.1).

Classification and pathogenesis of acute leukaemia

Acute leukaemias are usually aggressive diseases in which malignant transformation occurs in the haemopoietic stem cell or early progenitors. Genetic damage is believed to involve several key biochemical

steps resulting in: (i) an increased rate of proliferation; (ii) reduced apoptosis and (iii) a block in cellular differentiation. Together these events cause accumulation of the early bone marrow haemopoietic cells which are known as *blast cells*. The dominant clinical feature of these diseases is usually bone marrow failure caused by accumulation of blast cells although organ infiltration also occurs. If untreated these diseases are usually rapidly fatal but, paradoxically, they are also easier to cure than chronic leukaemias.

Acute leukaemia is defined as the presence of over 20% of blast cells in the blood or bone marrow at clinical presentation. It can be diagnosed with even less than 20% blasts if specific leukaemia-associated cytogenetic or molecular genetic abnormalities are present. It is further subdivided into acute myeloid leukaemia (AML) and acute lymphoblastic leukaemia (ALL) on the basis of whether the blasts are shown to be myeloblasts or lymphoblasts (Table 12.2).

Differentiation of ALL from AML

In most cases the clinical features and morphology on routine staining distinguish ALL from AML. In ALL the blasts show no differentiation (with the exception of B-cell ALL (B-ALL)) whereas in AML some evidence of differentiation to granulocytes or monocytes is usually seen in the blasts or their progeny. Specialized tests are needed to confirm the diagnosis of AML or ALL and to subdivide

Table 12.1 Classification of leukaemias.

Acute (see Table 12.2)

Acute myeloid leukaemia: M₀–M₇

Acute lymphoblastic leukaemia: L₁–L₃

Chronic (see Tables 13.1 and 15.1)

Chronic myeloid leukaemias

Chronic lymphoid leukaemias

Table 12.2 Classification of acute myeloid (AML) and acute lymphoblastic (ALL) leukaemia according to the French–American–British (FAB) groups. For more detailed WHO classification, see page 366.

AML	Cytogenetic abnormality
M ₀ undifferentiated	
M ₁ without maturation	
M ₂ with granulocytic maturation	t(8; 21)
M ₃ acute promyelocytic	t(15; 17)
M ₄ granulocytic and monocytic maturation	inv (16)
M ₅ monoblastic (M _{5a}) or monocytic (M _{5b})	
M ₆ erythroleukaemia	
M ₇ megakaryoblastic	
ALL	
L ₁ blast cells small, uniform high nuclear to cytoplasmic ratio	
L ₂ blast cells larger, heterogeneous, lower nuclear to cytoplasmic ratio	
L ₃ vacuolated blasts, basophilic cytoplasm (usually B-ALL)*	

* In WHO classification, L₃ cases with mature immunophenotype and Burkitt's lymphoma translocations are not classified as ALL (p. 367).

cases of AML or ALL into their different subtypes (Tables 12.3 and 12.4).

In a minority of cases of acute leukaemia the blast cells show features of both AML and ALL.

These features may be on the same cell (biphenotypic) or on separate populations (bilineal) and they include inappropriate expression of immunological markers or inappropriate gene rearrangements. This is termed hybrid acute leukaemia and treatment is usually given on the basis of the dominant pattern.

Acute lymphoblastic leukaemia

This is caused by an accumulation of lymphoblasts in the bone marrow and is the most common malignancy of childhood.

Incidence and pathogenesis

ALL is the most common form of leukaemia in children; its incidence is highest at 3–7 years, falling off by 10 years with a secondary rise after the age of 40 years. The common (CD10⁺) precursor B type which is most usual in children has an equal sex incidence; there is a male predominance for T-cell ALL (T-ALL).

The pathogenesis is varied. In a proportion of cases the first event occurs in the fetus *in utero*, with a secondary event possibly precipitated by infection in childhood. This is discussed further on p. 130. In other cases, the disease seems to arise as a postnatal mutation in an early lymphoid progenitor cell.

Table 12.3 Specialized tests for acute lymphoblastic leukaemia (ALL) and acute myeloid leukaemia (AML).

	ALL	AML
<i>Cytochemistry</i>		
Myeloperoxidase	–	+ (including Auer rods)
Sudan black	–	+ (including Auer rods)
Non-specific esterase	–	+ in M ₄ , M ₅
Periodic acid–Schiff	+ (coarse block positivity in ALL)	+ (fine blocks in M ₆)
Acid phosphatase	+ in T-ALL (Golgi staining)	+ in M ₆ (diffuse)
<i>Immunoglobulin and TCR genes</i>	Precursor B-ALL: clonal rearrangement of immunoglobulin genes T-ALL: clonal rearrangement of TCR genes	Germline configuration of immunoglobulin and TCR genes

Chromosomes and genetic analysis (see Table 10.2)

Immunological markers (flow cytometry) (Table 12.4)

B-ALL, B-cell acute lymphoblastic leukaemia; T-ALL, T-cell acute lymphoblastic leukaemia; TCR, T-cell receptor.

Table 12.4 Immunological markers for classification of acute myeloid (AML) and acute lymphoblastic (ALL) leukaemia.

Marker	AML	ALL	
		Precursor B*	T
<i>Myeloid</i>			
CD13	+	–	–
CD33	+	–	–
CD117	+	–	–
Glycophorin	+(M ₆)	–	–
Platelet antigens (e.g. CD41)	+(M ₇)	–	–
Myeloperoxidase	+(M ₀)		
<i>B lineage</i>			
CD19	–	+	–
cCD22	–	+	–
cCD79a	–	+	–
CD10	–	+ or –	–
cIg	–	+(pre-B)	–
SIg	–	– (early pre-B)	–
TdT	–	+	+
<i>T lineage</i>			
CD7	–	–	+
cCD3	–	–	+
CD2	–	–	+
TdT	–	+	+

c, Cytoplasmic; S, surface.

* B-ALL resembles precursor B-ALL immunologically but has surface immunoglobulin (Ig) and is terminal deoxynucleotidyl transferase negative (TdT[–]).

Classification

This may be on the basis of morphology or immunological markers.

Morphology The French–American–British (FAB) group subclassifies ALL into three subtypes (Table 12.2 and Fig. 12.1):

- 1 The L₁ type show uniform, small blast cells with scanty cytoplasm;
- 2 The L₂ type comprise larger blast cells with more prominent nucleoli and cytoplasm and with more heterogeneity; and
- 3 The L₃ blasts are large with prominent nucleoli, strongly basophilic cytoplasm and cytoplasmic vacuoles.

Immunophenotype Immunological markers can be used to divide ALL cases into early pre-B, pre-B, B- and T-cell subtypes (Table 12.4).

Clinical features

Clinical features are a result of the following.

1 Bone marrow failure Anaemia (pallor, lethargy and dyspnoea); neutropenia (fever, malaise, features of mouth, throat, skin, respiratory, perianal or other infections; Fig. 12.3); and thrombocytopenia (spontaneous bruises, purpura, bleeding gums and menorrhagia; Fig. 12.4).

2 Organ infiltration Tender bones, lymphadenopathy (Fig. 12.5a), moderate splenomegaly, hepatomegaly and meningeal syndrome (headache, nausea and vomiting, blurring of vision and diplopia). Fundal examination may reveal papilloedema and sometimes haemorrhage. Many patients have a fever which usually resolves after starting chemotherapy. Less common manifestations include testicular swelling (Fig. 12.5b) or signs of mediastinal compression in T-ALL (Fig. 12.6).

Investigations

Haematological investigations may reveal a normochromic, normocytic anaemia with thrombocytopenia in most cases. The total white cell count may be decreased, normal or increased to $200 \times 10^9/L$ or more. Blood film examination typically shows a variable numbers of blast cells. The bone marrow is hypercellular with >20% leukaemic blasts. The blast cells are characterized by morphology, immunological tests and cytogenetic analysis. Identification of the immunoglobulin/T-cell receptor (TCR) gene rearrangement, (aberrant) immunophenotype and molecular genetics of the tumour cells is important to determine treatment and to detect minimal residual disease during follow-up.

Lumbar puncture for cerebrospinal fluid examination should be performed and may show that the spinal fluid has an increased pressure and contains leukaemic cells. Biochemical tests may reveal a raised serum uric acid, serum lactate dehydrogenase or, less commonly, hypercalcaemia. Liver and renal function tests are performed as a baseline before treatment begins. Radiography may reveal




Fig. 12.1 Acute lymphoblastic leukaemia. (a) L_1 subtype—blasts show scanty cytoplasm without granules. (b) L_2 subtype—blasts are larger and heterogeneous with more abundant cytoplasm. (c) L_3 subtype—blasts are deeply basophilic with cytoplasmic vacuolation.

(d) Periodic acid–Schiff (PAS) staining reveals coarse granules. (e) Indirect immunofluorescence reveals nuclear terminal deoxynucleotidyl transferase (TdT) (green) and membrane CD10 (orange). (Courtesy of Professor G. Janossy)




Fig. 12.2 Development of three cell lineages from pluripotential stem cells giving rise to the three main immunological subclasses of acute leukaemia. The immunological characterization using pairs of markers is shown, as well as the three markers characterizing the early 'stem' cells. ALL, acute lymphoblastic leukaemia; AML, acute myeloid leukaemia; c, cytoplasmic.



(a)



(c)

Fig. 12.3 (a) An orbital infection in a female patient (aged 68 years) with acute myeloid leukaemia and severe neutropenia (haemoglobin 8.3 g/dL, white cells $15.3 \times 10^9/\text{L}$, blasts 96%, neutrophils 1%, platelets $30 \times 10^9/\text{L}$). (b) Acute myeloid leukaemia: top plaque *Candida albicans* on soft palate; lower: plaque *Candida albicans* in the mouth, with lesion of herpes simplex on the upper lip.



(b)

(c) Skin infection (*Pseudomonas aeruginosa*) in a female patient (aged 33 years) with acute lymphoblastic leukaemia receiving chemotherapy and with severe neutropenia (haemoglobin 10.1 g/dL, white cells $0.7 \times 10^9/\text{L}$, neutrophils $<0.1 \times 10^9/\text{L}$, lymphocytes $0.6 \times 10^9/\text{L}$, platelets $20 \times 10^9/\text{L}$).

lytic bone lesions and a mediastinal mass caused by enlargement of the thymus and/or mediastinal lymph nodes characteristic of T-ALL (Fig. 12.6).


The differential diagnosis includes AML, aplastic anaemia (with which ALL sometimes presents), marrow infiltration by other malignancies (e.g. rhabdomyosarcoma, neuroblastoma and Ewing's sarcoma), infections such as infectious mononucleosis and pertussis, juvenile rheumatoid arthritis and immune thrombocytopenic purpura.

Cytogenetics and molecular genetics

Cytogenetic analysis shows differing patterns in infants, children and adults which partly explains the different prognoses of these groups (Fig. 12.7). Cases are stratified according to the number of chromosomes in the tumour cell (*ploidy*) or by specific molecular abnormalities. Hyperdiploid cells have >50 chromosomes and generally have a good prognosis whereas hypodiploid cases carry a poor



Fig. 12.4 Purpura over the lower limbs in a male patient (aged 53 years) with acute leukaemia.



(a)




(b)


Fig. 12.5 Acute lymphoblastic leukaemia. (a) Marked cervical lymphadenopathy in a boy. (b) Testicular swelling and erythema on the left-hand side of the scrotum caused by testicular infiltration. (Courtesy of Professor J.M. Chessels)

prognosis. The most common specific abnormality in childhood ALL is the $t(12; 21)(p13; q22)$ *TEL-AML1* translocation (p. 133). The AML1 protein plays an important part in transcriptional control of haemopoiesis and this is repressed by the *TEL-AML1* fusion protein. The frequency of the Philadelphia chromosome translocation $t(9; 22)$ increases with age and carries a poor prognosis. Translocations of chromosome 11q23 involve the

MLL gene and are seen particularly in cases of infant leukaemia. Using more sensitive molecular genetic tests, as well as fluorescence *in situ* hybridization (FISH) analysis, some cases normal by conventional cytogenetic testing are found to have fusion genes or other genetic abnormalities. These molecular genetic changes carry prognostic significance whether or not the corresponding chromosomal change is present.



(a)



(b)

Fig. 12.6 Chest X-ray of a boy aged 16 years with acute lymphoblastic leukaemia (T-ALL). (a) There is a large mediastinal mass caused by thymic enlargement at presentation. (b) After 1 week of therapy with prednisolone, vincristine and daunorubicin the mass has resolved.




Fig. 12.7 Cytogenetic subsets of acute lymphoblastic leukaemia (ALL). The incidence of different cytogenetic abnormalities in infants, children and adults.

Treatment

This may be conveniently divided into supportive and specific treatments.

General supportive therapy

General supportive therapy for bone marrow failure is described in Chapter 11 and includes the insertion of a central venous cannula, blood product support and prevention of tumour lysis syndrome. Any episode of fever must be treated promptly.

Specific therapy

Specific therapy of ALL is with chemotherapy and sometimes radiotherapy (Fig. 12.8). These are used in various phases in a treatment course which usually has four components (Fig. 12.9). The protocols differ in infants, children and adults and are also *risk adjusted* to reduce the treatment given to patients with good prognosis. The rare B-ALL is treated by a different protocol involving short and intensive courses of drugs.

Remission induction

At presentation, the patient with acute leukaemia has a very high tumour burden and is at great risk from the complications of bone marrow failure and leukaemic infiltration. The aim of *remission induction* is to rapidly kill most of the tumour cells and get the patient into remission. This is defined as less than 5% blasts in the bone marrow, normal peripheral blood count and no other symptoms or signs of the disease. Prednisolone or dexamethasone (in children), vincristine and asparaginase are the drugs

usually used and they are very effective—achieving remission in over 90% of children and in 80–90% of adults (in whom daunorubicin is also usually added). However, it should be remembered that remission is not the same as cure. In remission a patient may still be harbouring large numbers of tumour cells and without further chemotherapy virtually all patients will relapse. Nevertheless, achievement of remission is a valuable first step in the treatment course. Patients who fail to achieve remission have a poor prognosis.

Intensification (consolidation)

These courses use high doses of multidrug chemotherapy in order to completely reduce or eliminate the tumour burden to very low levels. The doses of chemotherapy are near the limit of patient tolerability and during intensification blocks patients may need a great deal of support. Typical protocols involve the use of vincristine, cyclophosphamide, cytosine arabinoside, daunorubicin, etoposide, thioguanine or mercaptopurine given as blocks in different combinations. The optimal number of intensification blocks is under trial but two or three is typical in children, with more blocks given in adults.

Central nervous system directed therapy

Few of the drugs given systemically reach the cerebrospinal fluid (CSF) and specific treatment is required to prevent or treat central nervous system (CNS) disease. Options are high-dose methotrexate given intravenously, intrathecal methotrexate or cytosine arabinoside or cranial irradiation. Cranial irradiation is now avoided as far as possible in




Fig. 12.8 Acute leukaemia: principles of therapy. ALL, acute lymphoblastic leukaemia; chemo \pm TBI, chemotherapy \pm total-body irradiation; SCT, stem cell transplantation.

children because of substantial side-effects. CNS relapses still occur and present with headache, vomiting, papilloedema and blast cells in the CSF. Treatment is with intrathecal methotrexate, cytosine arabinoside and hydrocortisone, with or without cranial irradiation and systemic reinduction because bone marrow disease is usually also present.

Maintenance

This is given for 2 years in girls and adults and for 3 years in boys, with daily oral mercaptopurine and once-weekly oral methotrexate. Intravenous vincristine with a short course (5 days) of oral corticosteroids is added at monthly or 3-monthly (in adults) intervals. The value of tests for minimal residual disease (MRD) at the end of induction or during consolidation is being explored in trials in which the intensity of consolidation or maintenance therapy is reduced in those who rapidly become MRD

negative, whereas more intensified therapy, or even stem cell transplantation, is given to those with persistent MRD. There is a high risk of varicella or measles during maintenance therapy in children who lack immunity to these viruses. If exposure to these infections occurs, prophylactic immunoglobulin should be given. In addition, oral co-trimoxazole is given to reduce the risk of *Pneumocystis carinii*.

Other treatments

Allogeneic stem cell transplantation is indicated for patients with Ph (BCR-ABL) positive ALL, for primary refractory disease or for early relapsed disease. Imatinib mesylate is included into treatment protocols for patients with BCR-ABL positive ALL.

Prognosis

There is a great variation in the chance of individual

Induction
e.g. vincristine, asparaginase,
dexamethasone (or prednisolone) ± daunorubicin

Consolidation
e.g. daunorubicin, cytosine arabinoside, vincristine,
etoposide, thioguanine or mercaptopurine,
cyclophosphamide in one to four courses


Possible stem cell
transplantation

Cranial prophylaxis
e.g. high dose systemic methotrexate
+ multiple intrathecal methotrexate
(or cytosine arabinoside)
or multiple intrathecal methotrexate
cranial irradiation (1800–2400 rad)
+ intrathecal methotrexate

Maintenance therapy
e.g. mercaptopurine, methotrexate, vincristine,
dexamethasone (or prednisolone)

Late intensification (as consolidation)

Maintenance therapy as above (2–3 years)



(a)

(b)

Fig. 12.9 Acute lymphoblastic leukaemia (ALL). (a) Flow chart illustrating typical treatment regimen. (b) Kaplan-Meier analyses of overall survival in 2628 children with newly diagnosed ALL. (From Pui CH and Evans WE (2006) Treatment of acute lymphoblastic leukaemia. *New Engl J Med* 354,169)

Table 12.5 Prognosis in acute lymphoblastic leukaemia (ALL).

	Good	Poor
WBC	Low	High (e.g. $>50 \times 10^9/L$)
Sex	Girls	Boys
Immunophenotype	B-ALL	T-ALL (in children)
Age	Child	Adult (or infant <2 years)
Cytogenetics	Normal or hyperdiploidy (>50) TEL rearrangement	Ph+, 11q23 rearrangements
Time to clear blasts from blood	<1 week	>1 week
Time to remission	<4 weeks	>4 weeks
CNS disease at presentation	Absent	Present
Minimal residual disease	Negative at 1–3 months	Still positive at 3–6 months

CNS, central nervous system; Ph+, Philadelphia chromosome positive; WBC, white blood cell count.

patients achieving a long-term cure based on a number of biological variables (Table 12.5). Approximately 90% of children can expect to be cured, whereas in adults this drops significantly to less

than 5% over the age of 70 years. When treatment fails, death usually occurs because of resistant disease or from infections or other complications during treatment.

Table 12.6 Prognosis in acute myeloid leukaemia (AML).*

	Favourable	Unfavourable
Cytogenetics	t(15; 17) t(8; 21) inv(16) NPM mutation	Deletions of chromosome 5 or 7 <i>Fli-3</i> mutation 11q23 t(6; 9) abn(3q) Complex rearrangements
Bone marrow response to remission induction	<5% blasts after first course	>20% blasts after first course
Age	<60 years	>60 years
Onset	Primary	Secondary

* Detection of the relevant molecular abnormality, for example by fluorescence *in situ* hybridization (FISH) analysis, carries the same prognostic significance whether or not the corresponding chromosomal abnormality is present.

Treatment of relapse

The treatment of this is unsatisfactory at present. If it occurs during or soon after initial chemotherapy the outlook is very poor. Novel drugs such as clofarabine may help. Chemotherapy is usually followed, where possible, by allogeneic stem cell transplantation.

Acute myeloid leukaemia

Incidence and pathogenesis

AML occurs in all age groups. It is the common form of acute leukaemia in adults and is increasingly common with age. AML forms only a minor fraction (10–15%) of the leukaemias in childhood. An important distinction is between *primary AML* which appears to arise *de novo* and *secondary AML* which can develop from myelodysplasia and other haematological diseases such as the myeloproliferative diseases or follow previous treatment with chemotherapy. The two types are associated with distinct genetic markers and have different prognoses (Table 12.6). In addition, cytogenetic abnormalities and response to initial treatment have a major influence on prognosis (Table 12.6).

The most common genetic abnormality is the presence of internal tandem repeats of the *FLT-3* gene whose expression is normally tightly regulated in healthy CD34⁺ cells. Some cases of primary AML demonstrate the chromosomal translocations inv(16) and t(8; 21) which generate fusion proteins involving the cord binding factor (CBF) genes

(p. 138). These, together with the t(15; 17) variant (p. 138) exhibit a good prognosis. Many cases of AML with an apparently normal karyotype carry mutations in the nucleophosmin (NPM) gene and these also carry a favourable prognosis. This mutation leads to dysregulated cytoplasmic expression of NPM protein which normally acts as a nucleocytoplasmic shuttling protein and regulates the p53 pathway (Fig. 1.8). On the other hand, other chromosome abnormalities carry an unfavourable prognosis (Table 12.6). Those with normal karyotype may also be subdivided into favourable or poor prognosis on the basis of DNA microarray analysis.

Classification

Classification is usually based on the morphological criteria of the FAB (French–American–British) scheme (modified by the World Health Organization, WHO) in which at least 20% blast cells in the blood or bone marrow are required and which divides AML into eight variants (Table 12.2; Fig. 12.10). These FAB subtypes are associated with characteristic patterns of cytochemical stains (Fig. 12.11), immunophenotype and chromosomal changes (Chapter 10; Table 12.4). The typical ‘myeloid immunophenotype’ is CD13⁺, CD33⁺ and TdT⁺ (Table 12.4; Fig. 12.2) and special antibodies are helpful in the diagnosis of AML M₀ (a rare undifferentiated form), M₆ or M₇ (Table 12.4).

Although the distinct AML subtypes are in fact different genetic diseases, their grouping together is




Fig. 12.10 The French–American–British (FAB) classification of acute myeloid leukaemia. (a) M_1 blast cells show few granules but may show Auer rods, as in this case; (b) M_2 cells show multiple cytoplasmic granules; (c) M_3 blast cells contain prominent granules or multiple Auer rods; (d) M_4 blasts have some monocytoid differentiation; (e) M_{5a} —monoblastic leukaemia in which $>80\%$ of blasts are monoblasts; (f) M_{5b} —monocytic but $<80\%$ of blasts are monoblasts.

(Continued)





Fig. 12.10 (Continued) (g) M_6 showing preponderance of erythroblasts; (h) M_7 —megakaryoblastic leukaemia showing cytoplasmic blebs on blasts.

Fig. 12.11 Cytochemical staining in acute myeloid leukaemia. (a) Sudan black B shows black staining in the cytoplasm. (b) M_4 (myelomonocytic): non-specific esterase/chloracetate staining shows orange-staining monoblast cytoplasm and blue-staining (myeloblast) cytoplasm.



valid as generally their treatment and prognosis are similar. However, differences in treatment according to subtype are being introduced.

Clinical features

Clinical features resemble those of ALL. Anaemia and thrombocytopenia are often profound. A bleeding tendency caused by thrombocytopenia and disseminated intravascular coagulation (DIC) is characteristic of the M_3 variant of AML. Tumour cells can infiltrate a variety of tissues. Gum hypertrophy and infiltration (Fig. 12.12), skin involvement and CNS disease are characteristic of the myelomonocytic (M_4) and monocytic (M_5) types. An isolated mass of leukaemic blasts is usually referred to as a granulocytic sarcoma.

Investigations

The general haematological and biochemical findings are similar to those seen in ALL. Tests for DIC are often positive in patients with the promyelocytic (M_3) variant of AML. As explained earlier, cytogenetic and molecular analysis is critical for determining the prognosis and developing a treatment plan (Table 12.6).

Treatment

Management is both supportive and specific.

1 Supportive treatment is based on the same principles as that for ALL. Problems unique to AML include the haemorrhagic syndrome associated with the AML M_3 variant. The disease may present



Fig. 12.12 Acute myeloid leukaemia FAB type M₅ (monocytic): the gums are swollen and haemorrhagic because of infiltration by leukaemic cells.

with catastrophic haemorrhage or this may develop in the first few days of treatment. It is treated as for DIC with multiple platelet transfusions and replacement of clotting factors with fresh frozen plasma (p. 349). In addition all-*trans* retinoic acid (ATRA) therapy is given in conjunction with chemotherapy for the M₃ disease.

ATRA syndrome is a specific complication that may arise after treatment of AML-M₃ with ATRA. Clinical problems, which are thought to result from the neutrophilia that follows differentiation of promyelocytes from the bone marrow (p. 138), include fever, hypoxia with pulmonary infiltrates and fluid overload. Treatment is with dexamethasone 10 mg intravenously twice daily and ATRA is only discontinued in very severe cases.

2 Specific therapy of AML is primarily with the use of intensive chemotherapy. This is usually given in four blocks each of approximately 1 week and the most commonly used drugs include cytosine arabinoside (in conventional or high doses), daunorubicin, idarubicin, mitoxantrone and etoposide (Fig. 12.13). All the AML subtypes (FAB M₀–M₇) are treated similarly except for the promyelocytic (M₃) variant associated with the t(15; 17) translocation in which ATRA is added to the initial chemotherapy. A typical good response in AML to cytotoxics is shown in Fig. 12.14. The drugs are myelotoxic with limited selectivity between leukaemic and normal marrow cells and so marrow failure resulting from




Fig. 12.13 Acute myeloid leukaemia: flow chart illustrating typical treatment regimen.

the chemotherapy is severe, and prolonged and intensive supportive care is required. Maintenance therapy is of no value except in AML M₃ and CNS prophylaxis is not usually given in AML.

An important concept developing in AML therapy is that of basing the treatment schedule of individual patients on their risk group. Favourable cytogenetics and remission after one course of chemotherapy both predict for a better prognosis. In contrast, monosomy 5 or 7 abnormalities, blast cells




Fig. 12.14 Typical flow chart for the management with chemotherapy of acute myeloid leukaemia.

with *Flt-3* mutations or poorly responsive disease places patients into poor risk groups which may need more intensive treatments (Table 12.6). Monoclonal immunoconjugates targeted against CD33 (e.g. Mylotarg) or CD45 provide an additional therapeutic option for AML therapy.

Stem cell transplantation

Autologous transplantation reduces the rate of relapse but adds further toxicity to the treatment regime and is of no overall benefit. Allogeneic stem cell transplantation (SCT) is used in some

centres in patients under 65 years old with a human leucocyte antigen (HLA) matching sibling donor with standard or poor risk AML in first remission, although other groups save it as an option for relapsed disease. Patients with *t*(8; 21), *t*(15; 17) and *inv*16 who go into remission after the first course do not have SCT unless they subsequently relapse.

Patients over 70 years of age

Results of AML therapy in the elderly are poor because of primary disease resistance and poor tolerability of intensive treatment protocols. Death




Fig. 12.15 Survival in children and adults with acute myeloid leukaemia (Medical Research Council trials). (a) Children 0-14 years; (b) patients 15-59 years; (c) age 60-69 years; (d) 70+ years. (Courtesy of Professor A.K. Burnett)

from haemorrhage, infection or failure of the heart, kidneys or other organs is more frequent than in younger patients. In elderly patients with serious disease of other organs, the decision may be made to use supportive care with or without gentle single-drug chemotherapy. However, in those otherwise well, combination chemotherapy similar to that used in younger patients may produce long-term remissions.

Treatment of relapse

Most patients suffer relapse of their disease and the outlook will depend on age, the duration of the first remission and the cytogenetic risk group. In addition to further chemotherapy, allogeneic SCT is usually performed in those patients who can tolerate the procedure and who have a suitable HLA matching related or unrelated donor. Arsenic trioxide is useful in management of relapsed M_3 variant.

Prognosis

The prognosis for patients with AML has been improving steadily, particularly for younger patients and most recently for the 60-70-year-old group.

Fifty per cent of children and young adults may expect a long-term cure (Fig. 12.15). Cytogenetic abnormalities and initial response to treatment are major predictors of prognosis. Tracking of minimal residual disease using molecular cytogenetic markers or aberrant phenotypes may be helpful in predicting long-term remission or relapse. For the elderly the situation is poor and less than 10% of those over 70 years of age can expect long-term remission.

Bibliography

- Bullinger L., Dohner K., Bair E. *et al.* (2004) Use of gene-expression profiling to identify prognostic subclasses in adult acute myeloid leukemia. *N Engl J Med* 350,1605-16.
- Burnett A.K. (2002) Current controversies: which patients with acute myeloid leukaemia should receive a bone marrow transplantation? An adult treater's view. *Br J Haematol* 118,357-64.
- Döhner K., Schlenk R.F., Habdank M. *et al.* (2005) Mutant nucleophosmin (NPM1) predicts favorable prognosis in younger adults with acute myeloid leukemia and normal cytogenetics: interaction with other gene mutations. *Blood* 106,3740-6.

- Falini B., Mecucci C., Tiacci E. *et al.* (2005) Cytoplasmic nucleophosmin in acute myelogenous leukemia with a normal karyotype. *N Engl J Med* 352,254–66.
- Fröhling S., Scholl C., Gilliland D.G., Levine R.L. (2005) Genetics of myeloid malignancies: pathogenetic and clinical implications. *J Clin Oncol* 23,6285–95.
- Gilham C., Peto J., Simpson J. *et al.* (2005) UKCCS Investigators. Day care in infancy and risk of childhood acute lymphoblastic leukaemia: findings from UK case-control study. *BMJ* 330,1294.
- Gokbuget N., Kneba M., Raff T. *et al.* (2002) Related articles, links risk-adapted treatment according to minimal residual disease in adult ALL. *Best Pract Res Clin Haematol* 15,639–52.
- Ottmann O., Druker B.J., Sawyers C.L. *et al.* (2002) A phase 2 study of imatinib in patients with relapsed or refractory Philadelphia chromosome-positive acute lymphoid leukemias. *Blood* 100,1965–71.
- Pui C.-H. (ed.) (2006) *Childhood Leukemias*, 2nd edn. Cambridge University Press, New York.
- Pui C.-H. and Evans W.E. (2006) Treatment of acute lymphoblastic leukemia. *N Engl J Med* 354,166–78.
- Pui C.-H., Relling M.V., Downing J.R. (2004) Mechanisms of disease: acute lymphoblastic leukemia. *N Engl J Med* 350,1535–48.
- Rowe J.M., Buck G., Burnett A.K. *et al.* (2005) Induction therapy for adults with acute lymphoblastic leukemia: results of more than 1500 patients from the international ALL trial: MRC UKALL XII/ECOG E2993. *Blood* 106,3760–7.
- Tallman M.S., Gilliland D.G., Rowe J.M. (2005) Drug therapy for acute myeloid leukemia. *Blood* 106,1154–63.
- Valk P.J.M., Verhaak R.G.W., Beijnen M.A. *et al.* (2004) Prognostically useful gene-expression profiles in acute myeloid leukemia. *N Engl J Med* 350,1617–28.

Chronic myeloid leukaemia, 174

Variants of chronic myeloid leukaemia, 180

Bibliography, 181

The chronic leukaemias are distinguished from acute leukaemias by their slower progression. Paradoxically, they are also more difficult to cure. Chronic leukaemias can be broadly subdivided into myeloid (Table 13.1) and lymphoid groups (Chapter 15).

Chronic myeloid leukaemia

Chronic myeloid leukaemia (CML) is a clonal disorder of a pluripotent stem cell. The disease accounts for around 15% of leukaemias and may occur at any age. The diagnosis of CML is rarely difficult and is assisted by the characteristic presence of the Ph chromosome (Fig. 13.1d). This results from the t(9;22) (q34;q11) translocation between chromosomes 9 and 22 as a result of which part of the proto-oncogene *c-ABL* is moved to the *BCR* gene on chromosome 22 (Fig. 13.1a) and part of chromosome 22 moves to

chromosome 9. The abnormal chromosome 22 is the Ph chromosome (Fig. 13.1d). In the Ph translocation 5' exons of *BCR* are fused to the 3' exons of *ABL* (Fig. 13.1b,c). The resulting chimeric *BCR-ABL* gene codes for a fusion protein of size 210 kDa (p210). This has tyrosine kinase activity in excess of the normal 145-kDa ABL product. The Ph translocation is also seen in a minority of cases of acute lymphoblastic leukaemia (ALL) and in some of these the breakpoint in *BCR* occurs in the same region as in CML. However, in other cases the breakpoint in *BCR* is further upstream, in the intron between the first and second exons, leaving only the first *BCR* exon intact. This chimeric *BCR-ABL* gene is expressed as a p190 protein which, like p210, has enhanced tyrosine kinase activity. In a minority of patients, the Ph abnormality cannot be seen by microscopic karyotypic analysis but the same molecular rearrangement is detectable by more sensitive techniques; for example, fluorescence *in situ* hybridization (FISH) (Fig. 13.1e) or polymerase chain reaction (PCR) (see p. 90). This Ph-negative *BCR-ABL*-positive CML behaves clinically like Ph-positive CML. As the Ph chromosome is an acquired abnormality of haemopoietic stem cells it is found in cells of both the myeloid (granulocytic, erythroid and megakaryocytic) and lymphoid (B and T cell) lineages.

A great increase in total body myeloid cell mass is responsible for most of the clinical features. In at least 70% of patients who do not respond well to imatinib there is a terminal metamorphosis to acute leukaemia, often preceded by an accelerated phase.

Table 13.1 Classification of chronic myeloid leukaemias (CML).

Chronic myeloid leukaemia, Ph positive (CML, Ph+)
(chronic granulocytic leukaemia, CGL)
Chronic myeloid leukaemia, Ph negative (CML, Ph-)
(atypical)
Juvenile chronic myeloid leukaemia
Chronic neutrophilic leukaemia
Eosinophilic leukaemia
Chronic myelomonocytic leukaemia (CMML) (see
myelodysplasia, p. 186)

Clinical features

This disease occurs in either sex (male : female ratio of 1.4 : 1), most frequently between the ages of 40 and 60 years. However, it may occur in children and neonates, and in the very old. In most cases there are no predisposing factors but the incidence was increased in survivors of the atom bomb exposures in Japan. Its clinical features include the following:

- 1 Symptoms related to hypermetabolism (e.g. weight loss, lassitude, anorexia or night sweats).
- 2 Splenomegaly is nearly always present and is frequently massive. In some patients, splenic enlargement is associated with considerable discomfort, pain or indigestion.
- 3 Features of anaemia may include pallor, dyspnoea and tachycardia.
- 4 Bruising, epistaxis, menorrhagia or haemorrhage from other sites because of abnormal platelet function.
- 5 Gout or renal impairment caused by hyperuricaemia from excessive purine breakdown may be a problem.

caemia from excessive purine breakdown may be a problem.

6 Rare symptoms include visual disturbances and priapism.

7 In up to 50% of cases the diagnosis is made incidentally from a routine blood count.

Laboratory findings

1 Leucocytosis is usually $>50 \times 10^9/L$ and sometimes $>500 \times 10^9/L$ (Fig. 13.2). A complete spectrum of myeloid cells is seen in the peripheral blood. The levels of neutrophils and myelocytes exceed those of blast cells and promyelocytes (Fig. 13.3).

2 Increased circulating basophils.

3 Normochromic, normocytic anaemia is usual.

4 Platelet count may be increased (most frequently), normal or decreased.

5 Neutrophil alkaline phosphatase score is invariably low. It is raised in the myeloproliferative diseases and infections.







Fig. 13.1 The Philadelphia chromosome. (a) There is translocation of part of the long arm of chromosome 22 to the long arm of chromosome 9 and reciprocal translocation of part of the long arms of chromosome 9 to chromosome 22 (the Philadelphia chromosome). This reciprocal translocation brings most of the *ABL* gene into the *BCR* region on chromosome 22 (and part of the *BCR* gene into juxtaposition with the remaining portion of *ABL* on chromosome 9). (b) The breakpoint in *ABL* is between



exons 1 and 2. The breakpoint in *BCR* is at one of the two points in the major breakpoint cluster region (M-BCR) in chronic myeloid leukaemia (CML) or in some cases of Ph+ acute lymphoblastic leukaemia (ALL). (c) This results in a 210-kDa fusion protein product derived from the *BCR-ABL* fusion gene. In other cases of Ph+ ALL, the breakpoint in *BCR* is at a minor breakpoint cluster region (m-BCR) resulting in a smaller *BCR-ABL* fusion gene and a 190-kDa protein. (Continued)



(d)



(i)

(e)



(ii)

Fig. 13.1 (Continued) (d) Karyotype showing the t(9;22)(q34;q11) translocation. The Ph chromosome is arrowed. (e) Visualization of the Philadelphia chromosome on: (i) dividing (metaphase); and (ii) quiescent (interphase) cells by fluorescence *in situ* hybridization (FISH) analysis (ABL probe in red and BCR probe in green) with fusion signals (red/green) on the Ph and der(9) chromosomes. (Courtesy of Dr Ellie Nacheva)



Fig. 13.2 Chronic myeloid leukaemia: peripheral blood showing a vast increase in buffy coat. The white cell count was $532 \times 10^9/L$.

- 6 Bone marrow is hypercellular with granulopoietic predominance.
- 7 Ph chromosome on cytogenetic analysis (conventional or FISH) of blood or bone marrow (Fig. 13.1).
- 8 Serum uric acid is usually raised.

Treatment

Treatment of chronic phase

Tyrosine kinase inhibitors Imatinib (Glivec) was designed as a specific inhibitor of the BCR-ABL fusion protein and blocks tyrosine kinase activity by competing with adenosine triphosphate (ATP) binding (Fig. 13.4). It is the first-line drug in the management of chronic phase disease. At a dose of 400 mg/day it is able to produce a complete haematological response in virtually all patients (Fig. 13.5). Side-effects include skin rash, fluid retention, muscle pains and nausea. Neutropenia and thrombocytopenia may occur and although neutropenia may be managed by granulocyte colony-stimulating factor (G-CSF) in some cases, dose reduction or cessation may be required. Imatinib is highly effective in reducing the number of tumour cells in the bone marrow and this may be monitored by karyotypic analysis of the bone marrow together with PCR analysis for BCR-ABL transcripts in marrow or blood. A complete cytogenetic response is defined as the absence of Ph-positive metaphases on cytogenetic analysis of the bone marrow. The aim is to achieve complete cytogenetic response in all patients as this is associated with improved survival. In the first year after starting treatment, blood should be studied every 3 months and the marrow




Fig. 13.3 Chronic myeloid leukaemia: peripheral blood film showing various stages of granulopoiesis including promyelocytes, myelocytes, metamyelocytes and band and segmented neutrophils.




Fig. 13.4 Mode of action of the tyrosine kinase inhibitor STI-571. It blocks the adenosine triphosphate (ATP) binding site.

at 6 monthly intervals. Patients with suboptimal response can be identified as those who:

- 1 Fail to achieve a complete haematological response after 3 months;
- 2 Fail to achieve any significant cytogenetic response in bone marrow after 6 months; or
- 3 Fail to achieve a major cytogenetic response in bone marrow – <25% Ph⁺ metaphases by cytogenetic analysis after 1 year.

It is unclear how patients who have a suboptimal response should be treated. Potential options are an increase in the dose of imatinib to 600 or 800 mg/day, substitution or addition of second generation tyrosine kinase inhibitors dasatinib or nilotinib which can overcome resistance to imatinib or early allogeneic stem cell transplantation. These questions, as well as combination of imatinib with other drugs (e.g. interferon or cytosine arabinoside), are being assessed in trials. Molecular screening for mutations in the *BCR-ABL* fusion gene can be predictive for




Fig. 13.5 Example of the haematological and cytogenetic response in a patient with chronic myeloid leukaemia who achieves complete remission with imatinib therapy. (a) The white cell count returns to normal within days. (b) Karyotypic examination of the bone marrow reveals a gradual reduction in the number of Philadelphia chromosomes over the first year. (c) Polymerase chain reaction (PCR) analysis of the bone marrow or blood shows a reduction in the number of *bcr-abl* transcripts in comparison with the normal *abl* transcript. *Bcr-abl* transcripts continue to be detected at a very low level but can become negative in some patients. In this case analysis was performed on bone marrow for the first 12 months and on peripheral blood thereafter.

acquired resistance to imatinib in patients in whom there is an increase in bcr-abl transcripts.

It is not yet known if patients may stop imatinib if they become negative for *BRC-ABL* transcripts by PCR analysis. It is possible that imatinib, and similar drugs in development, may prove to be a cure for some patients with CML but this will need much longer clinical follow-up than is available at the present time.

Allopurinol is used in the initial phase of treatment to prevent hyperuricaemia and gout.

Second-line drugs for patients unable to take imatinib or have a poor response to it are discussed below.


Chemotherapy Hydroxyurea treatment can control and maintain the white cell count in the chronic phase but usually needs to be given indefinitely. A typical regimen would be to start with 1.0–2.0 g/day and then to reduce this in weekly increments to a maintenance dosage of 0.5–1.5 g/day. The alkylating agent busulfan is also effective in controlling the disease but has considerable long-term side-effects and is now reserved for patients who are intolerant of hydroxyurea. Imatinib has now largely replaced both drugs.

α -Interferon This was often used after the white cell count had been controlled by hydroxyurea but has now been superseded by imatinib. A typical regi-

men would be from 3 (to 9) megaunits between three to seven times each week given as a subcutaneous injection. The aim is to keep the white cell count low (around $4 \times 10^9/L$). Almost all patients have symptoms of a 'flu-like' illness in the first few days of treatment which responds to paracetamol and gradually wears off. More serious complications include anorexia, depression and cytopenias (see Table 11.1). A minority (approximately 15%) of patients may achieve long-term remission with loss of the Ph chromosome on cytogenetic analysis although the *BCR-ABL* fusion gene can still be detected by PCR. Interferon produces an overall prolongation of the chronic phase with increased life expectancy. Combinations of interferon with pulses of cytosine arabinoside may be more effective than interferon alone.

Stem cell transplantation Allogeneic stem cell transplantation (SCT) is the only established curative treatment for CML but, because of the risk, is usually reserved for imatinib failures. The results are better when it is performed in chronic rather than acute or accelerated phases. Only patients below approximately 65 years of age can tolerate SCT (low intensity in the older group) and only 30% of these will have a matched sibling. The 5-year survival is approximately 50–70% although this is reduced by approximately 10% if transplantation is delayed for

Fig. 13.6 Example of donor leucocyte infusion (DLI) in the treatment of chronic myeloid leukaemia (CML) which relapsed following allogeneic stem cell transplantation (SCT). Polymerase chain reaction (PCR) analysis of the blood for the *BCR-ABL* transcript shows that there was transient loss of the transcript but molecular and cytogenetic relapse occurred at 10 months. One infusion of donor leucocytes led to re-establishment of a durable complete remission.



more than 1 year following diagnosis. Although international bone marrow donor panels are playing an increasingly important part in providing human leucocyte antigen (HLA) matching unrelated donors, allogeneic SCT can only be offered to a minority of patients. Leukaemia relapse post-transplant is a significant problem but donor leucocyte infusions are highly effective in CML (Fig. 13.6 and p. 261), particularly if relapse is diagnosed early by molecular detection of the *BCR-ABL* transcript.

Course and prognosis

CML usually shows an excellent response to imatinib in the chronic phase. Survival is now much prolonged by imatinib. It is possible but not yet established that the best responders to imatinib may never relapse. If death occurs it is usually from terminal acute transformation or from intercurrent haemorrhage or infection. The patients may be divided into prognostic groups according to age, spleen size, platelet count, peripheral blood or bone marrow blast cells on presentation. These are only rough guides to outcome. Ease of response to imatinib is most important.

Accelerated phase disease and blastic transformation

Acute transformation (20% or more blasts in the marrow) may occur rapidly over days or weeks




Fig. 13.7 Chronic myeloid leukemia: acute myeloblastic transformation. Peripheral blood showing frequent myeloblasts.

(Fig. 13.7). More commonly, the patient has an accelerated phase with anaemia, thrombocytopenia and an increase in basophils, eosinophils or blast cells in the blood and marrow. The spleen may be enlarged despite control of the blood count and the marrow may become fibrotic. The patient may be in this phase for several months during which the disease is less easy to control than in the chronic phase. In both the accelerated and acute phases, new chromosome abnormalities are often present. In approximately one-fifth of cases, acute transformation is lymphoblastic and patients may be treated in a similar way to ALL with a number of patients returning to the chronic phase for months or even a year or two. In the majority, transformation is into acute myeloid leukaemia (AML) or mixed types. These are more difficult to treat and survival is rare beyond a few months. Imatinib is valuable in the management of blastic transformation but resistance to treatment usually occurs within a few weeks. New tyrosine kinase inhibitors aimed at overcoming resistance to imatinib are in dasatinib and nilotinib (p. 178) trials. SCT is usually offered to younger patients with an HLA matching donor but the results are less good than in chronic phase.

Variants of chronic myeloid leukaemia

Philadelphia-negative chronic myeloid leukaemia

Less than 5% of patients with features suggestive of CML are negative for the Philadelphia chromosome and *BCR-ABL* translocation. These patients usually have some of the haematological features typical of myelodysplasia. The prognosis appears to be worse than for Ph-positive CML.

Juvenile chronic myeloid leukaemia

This rare condition affects young children and has characteristic clinical features including skin rashes, lymphadenopathy, hepatosplenomegaly and recurrent infections. The blood film shows monocytosis. A high haemoglobin F level is a useful diagnostic feature, the neutrophil alkaline phosphatase score is normal and the Ph chromosome test is negative. The prognosis is poor and SCT is the treatment of choice.

Eosinophilic leukaemia and chronic neutrophilic leukaemia

These are rare conditions in which there is a relatively pure proliferation of mature cells. Some patients with idiopathic hypereosinophilia have an interstitial deletion of chromosome 4 that results in a *FIP1L1-PDGFR*A fusion gene. These patients are often responsive to low doses of imatinib (p. 104). Patients with chronic neutrophilic leukaemia may have mild splenomegaly and in general the prognosis is good.

Chronic myelomonocytic leukaemia

This is classified with the myelodysplastic syndromes and with the myeloproliferative disorders (p. 186).

Bibliography

- Cools J., DeAngelo D.J., Gotlib J. *et al.* (2003) A tyrosine kinase created by fusion of the *PDGFR*A and *FIP1L1* genes as a therapeutic target of imatinib in idiopathic hypereosinophilic syndrome. *N Engl J Med* **348**,1201–14.
- Deininger M., Buchdunger E., Druker B.J. (2005) The development of imatinib as a therapeutic agent for chronic myeloid leukemia. *Blood* **105**,2640–53.
- Druker B.J., Talpaz M., Resta R.N. *et al.* (2001) Efficacy and safety of a specific inhibitor of the BCR-ABL tyrosine kinase in chronic myeloid leukemia. *N Engl J Med* **344**,1031–7.
- Kantarjian H., Giles F., Wunderle L. *et al.* (2006) Nilotinib in imatinib-resistant CML and Philadelphia chromosome positive ALL. *N Engl J Med* **354**,2542–51.
- Mauro M.J. and Deininger M.W.N. (2006) Chronic myeloid leukemia in 2006: a perspective. *Haematologica* **91**,152–9.
- Melo J.V., Deininger M.W. (2004) Biology of chronic myelogenous leukemia — signaling pathways of initiation and transformation. *Hematol Oncol Clin N Am* **18**,545–68.
- Ren R. (2005) Mechanisms of BCR-ABL in the pathogenesis of chronic myelogenous leukaemia. *Nat Rev Cancer* **5**,172–8.
- Talpaz M., Shah N.P., Kantarjian *et al.* (2006) Dasatinib in imatinib-resistant Philadelphia chromosome-positive leukemias. *N Engl J Med* **354**,2531–41.
- Willis S.G., Lange T., Demehri S. *et al.* (2005) High-sensitivity detection of BCR-ABL kinase domain mutations in imatinib-naïve patients: correlation with clonal cytogenetic evolution but not response to therapy. *Blood* **106**,2128–37.

Myelodysplasia (myelodysplastic syndromes), 182
 Myelodysplastic/myeloproliferative diseases, 186

Bibliography, 187

Myelodysplasia (myelodysplastic syndromes)

This is a group of clonal disorders of multipotent haemopoietic stem cells which are characterized by increasing bone marrow failure with quantitative and qualitative abnormalities of all three myeloid cell lines. A hallmark of the disease is ineffective haemopoiesis so that cytopenias often accompany a marrow of normal or increased cellularity. There is a tendency to progress to acute myeloid leukaemia (AML), although death often occurs before this develops. In most cases, the disease is *primary* but in a significant proportion of patients the disease is *secondary* to chemotherapy and/or radiotherapy which has been given for treatment of another malignancy.

Pathogenesis

The pathogenesis of myelodysplastic syndromes (MDS) is unclear but is presumed to start following genetic damage to a multipotent haemopoietic progenitor cell. This leads to increased stem cell proliferation but ineffective differentiation and maturation, leading to the paradox of a hypercellular bone marrow but a pancytopenia in peripheral blood. A high rate of apoptosis is present in bone marrow precursors. The immune system may have a role in suppressing bone marrow function; immunosuppression is sometimes used in treatment (see below).


Clinical features

The disease has an incidence of 4 in 100 000 and a slight male predominance. Over half of patients are over 70 years and fewer than 25% are less than 50 years old. The evolution is often slow and the disease may be found by chance when a patient has a blood count for some unrelated reason. The symptoms, if present, are those of anaemia, infections or of easy bruising or bleeding (Fig. 14.1). In some patients transfusion-dependent anaemia dominates the course, while in others recurring infections or spontaneous bruising and bleeding are the major clinical problems. Because the neutrophils, monocytes and platelets are often functionally impaired, spontaneous infections in some cases, or bruising or bleeding in others, may occur out of proportion to the severity of the cytopenia. The spleen is not usually enlarged.

Dysplastic features in bone marrow may be seen in a wide range of conditions such as excess alcohol intake, megaloblastic anaemia, recovery from cytotoxic chemotherapy and response to growth-factor therapy and these must be ruled out before making a diagnosis of MDS.

Laboratory findings

Peripheral blood Pancytopenia is a frequent finding. The red cells are usually macrocytic or dimorphic but occasionally hypochromic; normoblasts may be present. The reticulocyte count is low. Granulocytes



(a)



(b)

Fig. 14.1 Myelodysplasia. (a) A 78-year-old male patient with refractory anaemia had recurring infections of the face and maxillary sinuses associated with neutropenia (haemoglobin 9.8 g/dL; white cells 1.3×10^9 /L; neutrophils 0.3×10^9 /L; platelets 38×10^9 /L). (b) Purpura in a 58-year-old female with refractory anaemia (haemoglobin 10.5 g/dL; white cells 2.3×10^9 /L; platelets 8×10^9 /L).

are often reduced in number and frequently show lack of granulation (Fig. 14.2). Their chemotactic, phagocytic and adhesive functions are impaired. The Pelger abnormality (single or bilobed nucleus) is often present. The platelets may be unduly large or small and are usually decreased in number but in 10% of cases are elevated. In poor prognosis cases variable numbers of myeloblasts are present in the blood.

Bone marrow The cellularity is usually increased but is decreased in approximately 20% of cases. Multinucleate normoblasts and other dyserythropoietic features are seen (Fig. 14.2). The appearance of ring sideroblasts is caused by iron deposition in the mitochondria of erythroblasts. The granulocyte precursors show defective primary and secondary granulation; cells that are difficult to identify as either agranular myelocytes, monocytes or promonocytes are frequent. Megakaryocytes are abnormal with micronuclear, small binuclear or polynuclear forms. A small number of dysplastic cells may be seen in marrow from normal elderly individuals so at least 10% of the cells in a lineage should be dysplastic in order to consider the diagnosis of MDS. In a minority of cases the marrow is hypocellular and may resemble aplastic anaemia; in others there is fibrosis.

Cytogenetic abnormalities are more frequent in secondary than primary MDS and most commonly constitute partial or total loss of chromosomes 5 or 7 or trisomy 8. The loss of chromosome 5 bands q13 to q33 in elderly females with macrocytic anaemia, normal or raised platelet counts and micromegakaryocytes has been termed the 5q-syndrome and has a good prognosis. *RAS* oncogene (usually *N-RAS*) mutations occur in about 20% of cases and mutations of *FMS* in approximately 15%.

Classification of myelodysplastic syndromes

The myelodysplastic syndromes are classified on the basis of the blood count, their morphological appearance and the number of blast cells in blood or bone marrow (Table 14.1).

Although the classification appears rather complex, the principles are:




Fig. 14.2 Myelodysplasia: appearances of the peripheral blood and bone marrow. (a) Multinucleate polychromatic erythroblasts. (b) Perls' stain showing iron overload in macrophages of a bone marrow fragment. (c) Ring sideroblasts. (d) White cells showing pseudo-Pelger cells, agranular myelocytes and neutrophils. (e) Monocytoid cells and an agranular neutrophil. (f) Mononuclear megakaryocyte.

Table 14.1 The World Health Organization (WHO) classification of myelodysplasia.

Subtype	Blood	Bone marrow
Refractory anaemia (RA)	Anaemia, no blasts	Erythroid dysplasia only, <5% blasts
Refractory anaemia with ringed sideroblasts (RARS)	Anaemia, no blasts	Erythroid dysplasia only and >15% ringed sideroblasts
Refractory cytopenia with multilineage dysplasia (RCMD)	Bi- or pancytopenia or more cell lineages	Dysplasia in >10% of cells in ≥2 myeloid lineages
Refractory cytopenia with multilineage dysplasia (RCMD) and ringed sideroblasts	Bi- or pancytopenia or more cell lineages and	Dysplasia in >10% of cells in ≥2 myeloid lineages >15% ringed sideroblasts
Refractory anaemia with excess blasts-1 (RAEB-1)	Cytopenias; <5% blasts no Auer rods	Uni- or multilineage dysplasia 5–9% blasts
Refractory anaemia with excess blasts-2 (RAEB-2)	Cytopenias or 5–19% blasts or Auer rods	Uni- or multilineage dysplasia 10–19% blasts or Auer rods
Myelodysplasia syndrome unclassified (MDS-U)	Cytopenias no blasts or Auer rods	Myeloid or megakaryocytic dysplasia <5% blasts
MDS associated with isolated del (5q)	Anaemia; normal or increased platelets	Megakaryocytes with hypolobated nuclei <5% blasts

Patients with dysplastic features and over 20% blast cells in the bone marrow are considered to have *acute myeloid leukaemia with multilineage dysplasia* (p. 366).

- Dysplasia may be present solely in red cells (*refractory anaemia*) or present in two or more myeloid lineages (*refractory cytopenia with multilineage dysplasia*, RCMD).
- Both of these subsets can also be associated with *ring sideroblasts* and these subtypes are classified separately.
- If the blast cell count is increased in blood or bone marrow, the diagnosis is made of *refractory anaemia with excess blasts* and these subtypes have a poor prognosis.
- *5q-Syndrome* receives its own classification. It has a particularly good prognosis (Table 14.2).
- Cases with unilineage dysplasia of myeloid or megakaryocytic lineage are termed *unclassified*.

Treatment

This is often extremely difficult because chemotherapy rarely cures the disease and intensive or even low-dose chemotherapy may in some cases make the situation worse rather than better.

Table 14.2 World Health Organization (WHO) classification: clinical and laboratory features.

Subtype	MDS (%)	Cytogenetic abnormalities (%)	Median survival (months)
RA	5–10	25	66
RARS	10	<10	72
RCMD	24	50	33
RCMD-RS	15	50	33
RAEB	40	30–50	10 (RAEB-1)
Isolated del 5q	5	100	116

MDS, myelodysplastic syndromes; RA, refractory anaemia; RAEB, RA with excess blasts; RARS, RA with ringed sideroblasts; RCMD, refractory cytopenia with multilineage dysplasia; RCMD-RS, RCMD with ringed sideroblasts.

Low-risk myelodysplastic syndromes

Patients with less than 5% blasts in the marrow and only one cytopenia and favourable cytogenetics are defined as having low-grade MDS. They are usually

managed conservatively with red cell transfusions, platelet transfusions or antibiotics as required. Attempts may be made to improve marrow function with haemopoietic growth factors, either singly or in combination. Erythropoietin (Epo) benefits the anaemia in many cases of refractory anaemia and may obviate the need for transfusions. Granulocyte colony-stimulating factor (G-CSF) shows synergy with Epo and may increase the response rate. Ciclosporin or antilymphocyte globulin (ALG) occasionally improve patients, particularly those with a hypocellular bone marrow. In the long term, iron overload may be a problem after multiple transfusions; iron chelation therapy (p. 83) should be started after 30–50 units have been transfused and if the anaemia and the need for transfusion continues to be the dominant problem. Other agents (e.g. thalidomide or its derivative Lenalidomide) may be of benefit, especially in the 5q-syndrome. In selected patients, standard or low-intensity allogeneic transplantation offers a permanent cure.

High-risk myelodysplastic syndromes

In these patients, with 5% or more blasts in the marrow and often unfavourable cytogenetics and pancytopenia, a variety of treatments have been attempted to improve the overall prognosis with varying degrees of success. These treatments extend from general support alone to intensive chemotherapy.

General support care only This is most suitable in elderly patients with other major medical problems. Transfusions of red cells and platelets, and therapy with antibiotics and antifungals, are given as needed. Fewer patients in the high-risk group with anaemia are helped by recombinant Epo, even with the addition of G-CSF, than in the low-risk group.

Single-agent chemotherapy Hydroxyurea, etoposide, mercaptopurine or low-dose cytosine arabinoside may be given with some benefit to patients with refractory anaemia with excess blasts (RAEB). The demethylating agents 5' azacytidine or decitabine have been of benefit to approximately 60% of patient trials.

Intensive chemotherapy Chemotherapy as given in AML (p. 169) may be tried in high-risk patients. Although the majority of patients may expect to obtain a remission, relapse is almost inevitable and frequently occurs within a few months. The risks of intensive chemotherapy are great because prolonged pancytopenia may occur in some cases without normal haemopoietic regeneration, presumably because normal stem cells are not present.

Stem cell transplantation SCT offers the prospect of complete cure for MDS and the advent of non-myeloablative conditioning is increasing the age range of patients that may be treated. SCT is usually carried out in MDS without a complete remission being first obtained with chemotherapy, although in high-risk cases initial chemotherapy may be tried to reduce the blast proportion and the risk of recurrence of the MDS. Older age, poor-risk cytogenetics, advanced disease and treatment-related MDS all increase the risk of an unfavourable outcome.

Myelodysplastic/myeloproliferative diseases

These disorders are characterized by the presence of dysplastic features but also increased number of circulating cells in one or more lineages (Table 14.3).

Chronic myelomonocytic leukaemia

This constituted approximately 20% of cases of

Table 14.3 The World Health Organization (WHO) classification of myelodysplastic/myeloproliferative diseases.

Chronic myelomonocytic leukaemia	CMML
Atypical chronic myeloid leukaemia*	aCML
Juvenile myelomonocytic leukaemia	JMML
Myelodysplastic/myeloproliferative disease	Unclassifiable

* This is discussed in Chapter 13.

MDS when it was classified with them. It is defined by a persistent monocytosis of $>1.0 \times 10^9/\text{L}$. The total white cell count is usually raised and may exceed $100 \times 10^9/\text{L}$. Patients may develop skin rashes and around half have splenomegaly. Gum hypertrophy and lymphadenopathy may also be present. Treatment is difficult although hydroxyurea, etoposide or mercaptopurine may be useful. SCT may be tried in younger patients. Median survival is approximately 2 years, with increased marrow blasts a predictor of poor outcome.

Juvenile myelomonocytic leukaemia

This presents in the first 4 years of life and has features of both myelodysplasia and a myeloproliferative disease. There is often a skin rash, hepatosplenomegaly and lymphadenopathy. There is a monocytosis to $>1.0 \times 10^9/\text{L}$, clonal cytogenetic changes but not the *BCR-ABL* fusion gene. The only curative treatment is allogeneic SCT. If untreated, death usually occurs within 4 years, often from acute transformation with leukaemic infiltration (e.g. of the lungs).

Bibliography

- Bennett J.M. (ed.) (2002) *The Myelodysplastic Syndromes: Pathobiology and Clinical Management*. Marcel Dekker, New York.
- Bowen D., Culligan D., Jowitt S. *et al.* (2003) UK MDS Guidelines Group. Guidelines for the diagnosis and therapy of adult myelodysplastic syndromes. *Br J Haematol* 120,187–200.
- Cazzola M. and Malcovati L. (2005) Myelodysplastic syndromes: coping with ineffective hematopoiesis. *N Engl J Med* 352,536–8.
- Deeg, J., Bowen D.T., Gore S.D. *et al.* (eds) (2006) *Hematologic Malignancies*. Springer-Verlag, Berlin.
- Greenberg P., Cox C., LeBeau M.M. *et al.* (1997) International scoring system for evaluating prognosis in myelodysplastic syndromes. *Blood* 89,2079–88.
- Jaffe E.S., Harris N.L., Stein H., Vardiman J.W. (eds) (2001) *World Health Organization Classification of Tumours: Pathology and Genetics of Tumours of Haematopoietic and Lymphoid Tissues*. IARC Press, Lyon.
- List A., Kurtin S., Roe D.J. (2005) Efficacy of lenalidomide in myelodysplastic syndromes. *N Engl J Med* 352,549–57.
- Rowe J.M. (ed.) (2004) Myelodysplastic syndromes: etiology, natural history, current and future therapies. *Clin Haematol* 17,535–661.

CHAPTER 15

15

The chronic lymphoid leukaemias

B-cell diseases, 188

Lymphoma/leukaemia syndromes, 194

T-cell diseases, 195

Bibliography, 195

Several disorders are included in this group characterized by accumulation in the blood of mature lymphocytes of either B- or T-cell type (Table 15.1). There is considerable overlap with the lymphomas. In many cases of non-Hodgkin's lymphoma, lymphoma cells are found in the blood and the distinction between chronic leukaemia and lymphoma is arbitrary, depending on the relative proportion of the disease in soft tissue masses compared to blood and bone marrow. In general, the diseases are incurable but tend to run a chronic and fluctuating course.

Table 15.1 Classification of the chronic lymphoid leukaemias and leukaemia/lymphoma syndromes.*

B-cell	T-cell
<i>Chronic lymphoid leukaemias</i>	
Chronic lymphocytic leukaemia (CLL)	Large granular lymphocytic leukaemia
Prolymphocytic leukaemia (PLL)	T-cell prolymphocytic leukaemia (T-PLL)
Hairy cell leukaemia (HCL)	
Plasma cell leukaemia	
<i>Leukaemia/lymphoma syndromes</i>	
Splenic lymphoma with villous lymphocytes	Sézary syndrome
Follicular lymphoma	Adult T-cell leukaemia/lymphoma
Mantle cell lymphoma	Large cell lymphoma
Lymphoplasmacytic lymphoma	
Large cell lymphoma	

* WHO classification see page 367.

Diagnosis

This group is characterized by a chronic persistent lymphocytosis. Subtypes are distinguished by morphology, immunophenotype and cytogenetics. DNA analysis may be useful in showing a monoclonal rearrangement of either immunoglobulin or T-cell receptor genes.

B-cell diseases

Chronic lymphocytic leukaemia

Chronic lymphocytic leukaemia (CLL) is by far the most common of the chronic lymphoid leukaemias and has a peak incidence between 60 and 80 years of age. The aetiology is unknown but there are geographical variations in incidence. It is the most common of the leukaemias in the Western world but rare in the Far East. In contrast to other forms of leukaemia there is no higher incidence after previous chemotherapy or radiotherapy. There is a seven-fold increased risk of CLL in the close relatives of patients. The tumour cell appears to be a relatively mature B cell with weak surface expression of immunoglobulin (Ig) M or IgD. The cells accumulate in the blood, bone marrow, liver, spleen and lymph nodes as a result of a prolonged lifespan with impaired apoptosis.

Clinical features

1 The disease occurs in older subjects with only 15%



Fig. 15.1 Chronic lymphocytic leukaemia: bilateral cervical lymphadenopathy in a 67-year-old woman. Haemoglobin 12.5 g/dL; white blood count $150 \times 10^9/L$ (lymphocytes $146 \times 10^9/L$); platelets $120 \times 10^9/L$.



Fig. 15.2 Chronic lymphocytic leukaemia: herpes zoster infection in a 68-year-old female.

of cases before 50 years of age. The male to female ratio is 2 : 1.

2 Most cases are diagnosed when a routine blood test is performed. With increasing routine medical check-ups, this proportion is rising.

3 Symmetrical enlargement of cervical, axillary or inguinal lymph nodes is the most frequent clinical sign (Fig. 15.1). The nodes are usually discrete and non-tender. Tonsillar enlargement may be a feature.

4 Features of anaemia may be present. Patients with thrombocytopenia may show bruising or purpura.

5 Splenomegaly and, less commonly, hepatomegaly are common in later stages.

6 Immunosuppression is a significant problem resulting from hypogammaglobulinaemia and cellular immune dysfunction. Early in the disease course bacterial infections predominate but with advanced disease viral and fungal infections such as herpes zoster are also seen (Fig. 15.2).

Laboratory findings

1 Lymphocytosis. The absolute lymphocyte count is $>5 \times 10^9/L$ and may be up to $300 \times 10^9/L$ or more. Between 70 and 99% of white cells in the blood film appear as small lymphocytes. Smudge or smear cells are also present (Fig. 15.3).

2 Immunophenotyping of the lymphocytes shows them to be B cells (surface CD19⁺), weakly expressing surface immunoglobulin (IgM or IgD). This is

shown to be monoclonal because of expression of only one form of light chain (κ or λ ; p. 141). Characteristically, the cells are also surface CD5⁺ and CD23⁺ but CD79b⁻ and FMC7⁻ (Table 15.2). Similar clones are found in some healthy subjects but with a normal total lymphocyte count.

3 Normocytic normochromic anaemia is present in later stages as a result of marrow infiltration or hypersplenism. Autoimmune haemolysis may also occur (see below). Thrombocytopenia occurs in many patients.

4 Bone marrow aspiration shows lymphocytic replacement of normal marrow elements. Lymphocytes comprise 25–95% of all the cells. Trephine biopsy reveals nodular, diffuse or interstitial involvement by lymphocytes.

5 Reduced concentrations of serum immunoglobulins are found and this becomes more marked with advanced disease. Rarely a paraprotein is present.

6 Autoimmunity directed against cells of the haemopoietic system is common. Autoimmune haemolytic anaemia is most frequent but immune thrombocytopenia (p. 283), neutropenia and red cell aplasia are also seen.

Prognostic markers

Cytogenetics The four most common chromosome abnormalities are deletion of 13q14, trisomy 12, deletions at 11q23 and structural abnormalities of




Fig. 15.3 Chronic lymphocytic leukaemia: peripheral blood film showing lymphocytes with thin rims of cytoplasm, coarse condensed nuclear chromatin and rare nucleoli. Typical smudge cells are present.

Table 15.2 Immunophenotype of the chronic B-cell leukaemias/lymphomas (all cases CD19⁺).

	CLL	PLL	HCL	FL	MCL
SIg	Weak	++	++	++	+
CD5	+	-	-	-	+
CD22/FMC7	-	+	+	+	+
CD23	+	-	-	-	-
CD79b	-	++	-/+	++	++
CD103*	-	-	+	-	-

CLL, chronic lymphocytic leukaemia; FL, follicular lymphoma; HCL, hairy cell leukaemia; MCL, mantle cell lymphoma; PLL, prolymphocytic leukaemia.

* CD103 is positive only in HCL.

17p involving the p53 gene. These abnormalities carry prognostic significance (Table 15.3). The 13q14 deletion prevents expression of microRNAs which control expression of proteins relevant to the CLL lifespan (p. 137). Expression of a novel gene CLLU.1 is substantially greater in unmutated than mutated cases.

Somatic hypermutation of the immunoglobulin genes
When B cells recognize antigen in the germinal centre of secondary lymphoid tissues they undergo a process called *somatic hypermutation* in which

random mutations occur in the immunoglobulin heavy-chain gene (p. 113). In CLL, the *IGHV* gene shows evidence of this hypermutation in approximately 50% of cases whereas in the other cases the *VH* genes are unmutated. CLL with unmutated immunoglobulin genes has an unfavourable prognosis (Table 15.3).

Tumour cell phenotype ZAP-70 is a protein tyrosine kinase that is involved in cell signalling following recognition of antigen by antigen receptors on lymphocytes. Its expression is normally restricted to T cells but is also aberrantly expressed in cases of CLL. Its expression is associated with an unfavourable clinical outcome.

Staging

It is useful to stage patients at presentation both for prognosis and for deciding on therapy. The Rai and Binet staging systems are shown in Table 15.4. Typical survival ranges from 12 years for Rai stage 0 to less than 3 years for stage IV.

Treatment

Cures are rare in CLL and so the approach to therapy is conservative, aiming for symptom control rather than a normal blood count. Indeed, chemotherapy given too early in the disease can

Table 15.3 Prognostic factors in chronic lymphocytic leukaemia.

	Good	Bad
Stage	Binet A (Rai 0–I)	Binet B, C (Rai II–IV)
Sex	Female	Male
Lymphocyte doubling time	Slow	Rapid
Bone marrow biopsy appearance	Nodular	Diffuse
Chromosomes	Deletion 13q14	Trisomy 12; deletion 17p; deletion 11q23
VH immunoglobulin genes	Hypermutated	Unmutated
ZAP expression	Low	High
CD38 expression	Negative	Positive
CLLU.1 expression	Low	High
LDH	Normal	Raised

LDH, lactate dehydrogenase.

Table 15.4 Staging of chronic lymphocytic leukaemia (CLL).

(a) Rai classification

Stage	
0	Absolute lymphocytosis $>15 \times 10^9/L^*$
I	As stage 0 + enlarged lymph nodes (adenopathy)
II	As stage 0 + enlarged liver and/or spleen \pm adenopathy
III	As stage 0 + anaemia (Hb <10.0 g/dL)* \pm adenopathy \pm organomegaly
IV	As stage 0 + thrombocytopenia (platelets $<100 \times 10^9/L$)* \pm adenopathy \pm organomegaly

(b) International Working Party classification (From J.L. Binet *et al.* 1981)

Stage	Organ enlargement*	Haemoglobin† (g/dL)	Platelets† ($\times 10^9/L$)
A (50–60%)	0, 1 or 2 areas		
B (30%)	3, 4 or 5 areas	≥ 10	≥ 100
C ($<20\%$)	Not considered	<10	and/or <100

* One area = lymph nodes >1 cm in neck, axillae, groins or spleen, or liver enlargement.

† Secondary causes of anaemia (e.g. iron deficiency) or autoimmune haemolytic anaemia or autoimmune thrombocytopenia must be treated before staging.

shorten rather than prolong life expectancy. Many patients never receive treatment. Treatment is given for troublesome organomegaly, haemolytic episodes and bone marrow suppression. The lymphocyte count alone is not a good guide to treatment. Usually, patients in Binet stage C will need treatment as will some in stage B.

Chemotherapy

Chlorambucil This oral alkylating agent can be used as a daily treatment (e.g. 4–6 mg/day) or in a monthly cycle (e.g. 10 mg/m²/day for 7 days) and is effective in reducing disease bulk in the majority of cases. Typically the drug will need to be given for

several months after which a remission of variable duration will be obtained. Chlorambucil can be reintroduced when required although resistance may develop.

Purine analogues These drugs are effective in the treatment of chronic lymphoid leukaemias and lymphomas. The most effective agent in CLL appears to be fludarabine. Its place in the overall management of CLL is currently under trial. It is the drug of first choice in younger patients as well as having value in patients resistant to chlorambucil. Myelosuppression and prolonged reduction of CD4 (helper) T lymphocytes lead to an increased infection risk and prophylaxis against *Pneumocystis carinii* infection with co-trimoxazole is given until CD4 counts recover. The combination of fludarabine with cyclophosphamide (FC) is more effective than fludarabine alone and fludarabine, cyclophosphamide and rituximab (FCR) increases further the response rate and duration of response.

Monoclonal antibodies Campath-1H (anti-CD52) is a monoclonal antibody that is effective in killing white cells by complement fixation. It is given intravenously or subcutaneously over several weeks and as well as being valuable in resistant and relapsed disease, it may also find a role earlier in treatment. It is highly immunosuppressive and antibacterial and antiviral prophylaxis is advisable.

CD20 expression on CLL cells is not high; rituximab (p. 217), however, appears to be valuable in combination therapy (e.g. with fludarabine). It also has a role in treating autoimmune haemolytic anaemia, thrombocytopenia and red cell aplasia.

Corticosteroids Patients in bone marrow failure should be treated initially with prednisolone alone until there is significant recovery of the platelet, neutrophil and haemoglobin levels. The peripheral lymphocyte count initially rises as infiltrated organs shrink, but later the count falls. Corticosteroids are also indicated in autoimmune haemolytic anaemia or thrombocytopenia.

Other forms of treatment

Radiotherapy This is valuable in reducing the size of

bulky lymph node groups that are unresponsive to chemotherapy. Radiotherapy to the spleen may be valuable in late-stage disease.

Combination chemotherapy Cyclophosphamide, hydroxodaunorubicin, oncovin (vincristine) and prednisolone (CHOP; p. 213) with rituximab is sometimes used in late-stage cases and in patients refractory to chlorambucil.

Ciclosporin Red cell aplasia may respond to ciclosporin.

Splenectomy This is generally reserved for those patients with immune-mediated cytopenias that do not respond to short courses of steroids or those with painful bulky enlargement of the spleen.

Immunoglobulin replacement Immunoglobulin (e.g. 400 mg/kg/month by intravenous infusion) is useful for patients with hypogammaglobulinaemia and recurrent infections.

Stem cell transplantation This is currently an experimental approach in younger patients. Allogeneic stem cell transplantation (SCT) may be curative but has a high mortality rate. Autologous SCT after prior therapy with fludarabine and other drugs is undergoing clinical trials.

Course of disease

Many patients in Binet stage A or Rai stage 0 or I never need therapy and this is particularly likely for those with favourable prognostic markers (Table 15.3). For those who do need treatment a typical pattern is that of a disease that is responsive to several courses of chemotherapy before the gradual onset of extensive bone marrow infiltration, bulky disease and recurrent infection. The disease may transform into a localized high-grade lymphoma (Richter's transformation) or there may be the appearance of an increasing number of prolymphocytes that are resistant to treatment.

Prolymphocytic leukaemia

Although prolymphocytic leukaemia (PLL) may initially appear similar to CLL, the diagnosis is




Fig. 15.4 Prolymphocytic leukaemia: blood film showing prolymphocytes that have prominent central nucleoli and an abundance of pale cytoplasm.

made by the appearance of a majority of prolymphocytes in the blood. The prolymphocyte is around twice the size of a CLL lymphocyte and has a large central nucleolus (Fig. 15.4). B-cell PLL is three times more common than T-cell PLL.

PLL and CLL also differ in their clinical features. PLL typically presents with splenomegaly without lymphadenopathy and with a high and rapidly rising lymphocyte count. Anaemia is a poor prognostic feature. Treatment is difficult in PLL. Splenectomy is usually of benefit and purine nucleoside analogues and rituximab may help.

Hairy cell leukaemia

Hairy cell leukaemia (HCL) is an uncommon B-cell lymphoproliferative disease with a male : female ratio of 4 : 1 and a peak incidence at 40–60 years. Patients typically present with infections, anaemia or splenomegaly. Lymphadenopathy is very uncommon. Pancytopenia is usual at presentation and the lymphocyte count is rarely over $20 \times 10^9/L$. Monocytopenia is a distinctive feature. The blood film reveals a variable number of unusual large lymphocytes with villous cytoplasmic projections (Fig. 15.5). Immunophenotyping shows CD22, FMC7 and CD103 positivity in most cases (Table 15.2). The hairy cells stain for tartrate-resistant acid phosphatase (TRAP). The bone marrow trephine




Fig. 15.5 Hairy cell leukaemia: peripheral blood film showing typical 'hairy' cells with oval nuclei and finely mottled pale grey/blue cytoplasm with an irregular edge.

shows a characteristic appearance of mild fibrosis and a diffuse cellular infiltrate.

There are several effective treatments for HCL and a patient can expect a long-term remission. The

treatment of choice is 2-chlorodeoxyadenosine or deoxycoformycin and both agents achieve responses in over 90% of cases. In two-thirds of cases no relapse occurs, even after 5–10 years. HCL was one of the first diseases in which α -interferon was shown to be effective and it remains an excellent treatment. These drugs have largely replaced the need for splenectomy or combination chemotherapy.

Hairy cell leukaemia variant

Some cases of HCL have clear differences from the typical disease and warrant a separate classification. The white cell count is higher, the hairy cells have a prominent nucleolus and response to purine analogues and interferon is less satisfactory.

Splenic lymphoma with villous lymphocytes

This is now classified within the World Health Organization (WHO) as splenic marginal zone lymphoma (p. 367). It is characterized by massive splenomegaly and circulating monoclonal B cells with a villous appearance. It is a disease of the elderly with a benign clinical course. Although many patients will not need treatment, splenectomy is valuable and purine nucleoside analogues are also effective. The outlook is good.

Plasma cell leukaemia

This rare disease is characterized by a high number of circulating plasma cells. The clinical features tend to be a combination of those found in acute leukaemia (pancytopenia and organomegaly) with features of myeloma (hypercalcaemia, renal involvement and bone disease; Chapter 18). Treatment is with supportive care and systemic chemotherapy; for example, with CHOP, cyclophosphamide-VAD (vincristine, Adriamycin and dexamethasone) or cyclophosphamide, dexamethasone and thalidomide as for multiple myeloma (Chapter 18).

Lymphoma/leukaemia syndromes

Circulating malignant lymphoid cells occur in a variety of syndromes in association with otherwise typical cases of non-Hodgkin's lymphoma. This syndrome is most frequently seen in B-cell tumours of the follicle centre cell type (with circulating indented or cleaved nuclei; Fig. 15.6) and the course is that of the non-Hodgkin's lymphoma (p. 209). Mantle zone lymphoma is often associated with circulating B cells (Fig. 15.6) and, because these are CD5⁺, mantle zone lymphoma may be confused with CLL. Other types of lymphoma may also show




Fig. 15.6 Blood involvement by malignant lymphoma: (a) small cleaved lymphoid cells in follicle centre cell lymphoma; (b) mantle cell lymphoma; (c) large B-cell lymphoma.




Fig. 15.7 (a) Large granular lymphocytes in the peripheral blood. (b) Adult T-cell leukaemia/lymphoma. Typical convoluted lymphoid cells in peripheral blood.

tumour cells in the peripheral blood and bone marrow and in some cases it is difficult to define the disease as either lymphoma (with mainly soft tissue masses) or leukaemia.

T-cell diseases

Large granular lymphocytic leukaemia

Large granular lymphocytic leukaemia (LGL-L) is characterized by the presence of circulating lymphocytes with abundant cytoplasm and large azurophilic granules (Fig. 15.7a). Such cells may be either T or natural killer (NK) cells and show variable expression of CD16, CD56 and CD57. Cytopenia, especially neutropenia, is the main clinical problem although anaemia, splenomegaly and arthropathy with positive serology for rheumatoid arthritis are also common. The mean age is 50 years. Treatment may not be needed but, if required, steroids, cyclophosphamide, ciclosporin or methotrexate may relieve the cytopenia. Granulocyte colony-stimulating factor (G-CSF) or granulocyte-macrophage colony-stimulating factor (GM-CSF) have been used in cases associated with neutropenia.

Adult T-cell leukaemia/lymphoma

Adult T-cell leukaemia/lymphoma (ATLL) was the first malignancy to be associated with a human retrovirus, human T-cell leukaemia/lymphoma virus type 1 (HTLV-1). The virus is endemic in parts

of Japan and the Caribbean but the disease is very rare in people who have not lived in these areas. ATLL lymphocytes have a bizarre morphology with a convoluted 'clover-leaf' nucleus and a consistent CD4⁺ phenotype (Fig. 15.7b).

Most subjects infected with the virus do not develop the disease. The clinical presentation is often acute and dominated by hypercalcaemia, skin lesions, hepatosplenomegaly and lymphadenopathy. Diagnosis is by morphology and serology and although combination chemotherapy may be tried the prognosis is poor. Antiretroviral drugs may prove to have a valuable role.

Sézary syndrome

Patients with Sézary syndrome present with skin disease, usually a pruritic exfoliative erythroderma affecting the palms, soles and face ('red man syndrome'). Biopsy of the skin reveals lymphocytic infiltration and Sézary cells in peripheral blood have a characteristic morphology with deep nuclear clefting, similar to ATLL cells. A variety of treatments are available including chemotherapy, radiotherapy and a photoactivable drug (psoralens) combined with ultraviolet A light (PUVA).

Bibliography

- Binet J.L., Auquier A., Dighiero G. *et al.* (1981) A new prognostic classification of chronic lymphocytic leukemia derived from a multivariate survival analysis. *Cancer* 48,198–206.

- Buhl A.M., Jurlander J., Jørgensen F.S. *et al.* (2006) Identification of a gene on chromosome 12q22 uniquely overexpressed in chronic lymphocytic leukemia. *Blood* 107,2904–11.
- Calin G.A., Ferracin M., Cimmino A. *et al.* (2005) A microRNA signature associated with prognosis and progression in chronic lymphocytic leukemia. *N Engl J Med* 353,1793–801.
- Chiorazzi N., Rai K.R., Ferrarini M. (2005) Chronic lymphocytic leukemia. *N Engl J Med* 352,804–15.
- Crespo M., Bosch F., Villamor N. *et al.* (2003) ZAP-70 expression as a surrogate for immunoglobulin-variable-region mutations in chronic lymphocytic leukemia. *N Engl J Med* 348,1764–75.
- Montserrat E., Moreno C., Esteve J. *et al.* (2006) How I treat refractory CLL. *Blood* 107,1276–83.
- Oscier D., Fegan C., Hillmen P. *et al.* (2004) Guidelines Working Group of the UK CLL Forum. British Committee for Standards in Haematology. Guidelines on the diagnosis and management of chronic lymphocytic leukaemia. *Br J Haematol* 125,294–317.
- Paneesha S. and Milligan D.W. (2005) Stem cell transplantation for chronic lymphocytic leukaemia. *Br J Haematol* 128,145–52.
- Rai K.R., Sawitsky A., Cronkite E.P. *et al.* (1975) Clinical staging of chronic lymphocytic leukemia. *Blood* 46,219–34.
- Rawstron A.C., Yuille M.R., Fuller J. *et al.* (2002) Inherited predisposition to CLL is detectable as subclinical monoclonal B-lymphocyte expansion. *Blood* 100,2289–90.
- Stevenson F.K. and Caligaris-Cappio F. (2004) Chronic lymphocytic leukemia: revelations from the B-cell receptor. *Blood* 103,4389–95.
- Wiestner A., Rosenwald A., Barry T.S. *et al.* (2003) ZAP-70 expression identifies a chronic lymphocytic leukemia subtype with unmutated immunoglobulin genes, inferior clinical outcome, and distinct gene expression profile. *Blood* 101,4944–51.

CHAPTER 16

16

Hodgkin's lymphoma

Pathogenesis, 197

Clinical features, 197

Haematological and biochemical findings, 198

Diagnosis and histological classification, 198

Clinical staging, 200

Treatment, 201

Prognosis, 202

The late effects of Hodgkin's lymphoma and its treatment, 202

Bibliography, 202

Lymphomas are a group of diseases caused by malignant lymphocytes that accumulate in lymph nodes and cause the characteristic clinical features of lymphadenopathy. Occasionally, they may spill over into blood ('leukaemic phase') or infiltrate organs outside the lymphoid tissue.

The major subdivision of lymphomas is into Hodgkin's lymphoma and non-Hodgkin's lymphoma and this is based on the histological presence of Reed–Sternberg (RS) cells in Hodgkin's lymphoma.

Pathogenesis

Hodgkin's disease is a lymphoma in which RS cells are found in the disease tissue. The characteristic RS cells, and the associated abnormal mononuclear cells, are neoplastic whereas the infiltrating inflammatory cells are reactive. Immunoglobulin gene rearrangement studies suggest that the RS cell is of B-lymphoid lineage and that it is often derived from a B cell with a 'crippled' immunoglobulin gene caused by the acquisition of mutations that prevent synthesis of full-length immunoglobulin. The Epstein–Barr virus (EBV) genome has been detected in 50% or more of cases in Hodgkin tissue but its role in the pathogenesis is unclear.

Clinical features

The disease can present at any age but is rare in children and has a peak incidence in young adults. There is an almost 2 : 1 male predominance. The following symptoms are common.

1 Most patients present with painless, non-tender, asymmetrical, firm, discrete and rubbery enlargement of superficial lymph nodes (Fig. 16.1). The cervical nodes are involved in 60–70% of patients, axillary nodes in approximately 10–15% and inguinal nodes in 6–12%. In some cases the size of the nodes decreases and increases spontaneously. They may




Fig. 16.1 Cervical lymphadenopathy in a patient with Hodgkin's lymphoma.

become matted. Typically the disease is localized initially to a single peripheral lymph node region and its subsequent progression is by contiguity within the lymphatic system. Retroperitoneal nodes are also often involved but usually only diagnosed by computed tomography (CT) scan.

2 Clinical splenomegaly occurs during the course of the disease in 50% of patients. The splenic enlargement is seldom massive. The liver may also be enlarged because of liver involvement.

3 Mediastinal involvement is found in 6–11% of patients at presentation. This is a feature of the nodular sclerosing type, particularly in young women. There may be associated pleural effusions or superior vena cava obstruction.

4 Cutaneous Hodgkin's disease occurs as a late complication in approximately 10% of patients. Other organs (e.g. bone marrow, gastrointestinal tract, bone, lung, spinal cord or brain) may also be involved, even at presentation, but this is unusual.

5 Constitutional symptoms are prominent in patients with widespread disease. The following may be seen:

- (a) fever occurs in approximately 30% of patients and is continuous or cyclic;
- (b) pruritus, which is often severe, occurs in approximately 25% of cases;
- (c) alcohol-induced pain in the areas where disease is present occurs in some patients;
- (d) other constitutional symptoms include weight loss, profuse sweating (especially at night), weakness, fatigue, anorexia and cachexia. Haematological and infectious complications are discussed below.

Haematological and biochemical findings

1 Normochromic, normocytic anaemia is most common. Bone marrow involvement is unusual in early disease but if it occurs bone marrow failure may develop with a leucoerythroblastic anaemia.

2 One-third of patients have a neutrophilia; eosinophilia is frequent.

3 Advanced disease is associated with lymphopenia and loss of cell-mediated immunity.

4 The platelet count is normal or increased during early disease, and reduced in later stages.

5 The erythrocyte sedimentation rate (ESR) and C-reactive protein are usually raised and are useful in monitoring disease progress.

6 Serum lactate dehydrogenase (LDH) is raised initially in 30–40% of cases.

Diagnosis and histological classification

The diagnosis is made by histological examination of an excised lymph node. The distinctive multinucleate polyloid RS cell is central to the diagnosis of the four classic types (Figs 16.2 and 16.3) and mononuclear Hodgkin cells are also part of the malignant clone. These cells stain with CD30 and CD15 but are usually negative for B-cell antigen expression. Inflammatory components consist of lymphocytes, neutrophils, eosinophils, plasma cells and variable fibrosis.

Histological classification is into five types (Table 16.1), each of which implies a different prognosis. Nodular sclerosis and mixed cellularity are most frequent. Patients with lymphocyte predominant histology have the most favourable prognosis. Nodular lymphocyte predominant does not show RS cells and has many features of non-Hodgkin's lymphoma and may be treated as such.






Fig. 16.2 Diagrammatic representation of the different cells seen histologically in Hodgkin's lymphoma.



(a)



(b)



(c)

Fig. 16.3 Hodgkin's disease: (a) high power view of a lymph node biopsy showing two typical multinucleate Reed-Sternberg cells, one with a characteristic owl eye appearance, surrounded by lymphocytes, histiocytes and an eosinophil; (b) mixed cellularity; and (c) nodular sclerosing Hodgkin's lymphoma.

Table 16.1 Histology of Hodgkin's lymphoma (WHO classification).

Classic Hodgkin's lymphoma

Nodular sclerosis	Collagen bands extend from the node capsule to encircle nodules of abnormal tissue. A characteristic lacunar cell variant of the Reed-Sternberg cell is often found. The cellular infiltrate may be of the lymphocyte-predominant, mixed cellularity or lymphocyte-depleted type; eosinophilia is frequent
Mixed cellularity	The Reed-Sternberg cells are numerous and lymphocyte numbers are intermediate
Lymphocyte depleted	There is either a reticular pattern with dominance of Reed-Sternberg cells and sparse numbers of lymphocytes or a diffuse fibrosis pattern where the lymph node is replaced by disordered connective tissue containing few lymphocytes. Reed-Sternberg cells may also be infrequent in this latter subtype
Lymphocyte rich	Scanty Reed-Sternberg cells; multiple small lymphocytes with few eosinophils and plasma cells; nodular and diffuse types
Nodular lymphocyte-predominant	Reed-Sternberg cells are absent; abnormal polymorphic B cells

WHO, World Health Organization.

Clinical staging

The selection of appropriate treatment depends on accurate staging of the extent of disease (Table 16.2). Figure 16.4 shows the scheme (Ann Arbor) that is now recommended. Staging is performed by thorough clinical examination together with chest X-ray (Fig. 16.5) and CT scan to detect intrathoracic, intra-abdominal or pelvic disease (Fig. 16.6). It is also used to monitor response to therapy. Magnetic resonance imaging (MRI) scanning may be needed for particular sites (Table 16.2). Bone marrow trephine is carried out in some centres and liver biopsy may also be needed in difficult cases. Positron emission tomography (PET) scanning may also be useful in staging and PET scan combined with CT may be used in staging and to detect small foci of residual disease following treatment (Fig. 17.8).

Table 16.2 Techniques for staging of lymphoma.

Laboratory	Full blood count
	ESR
	Bone marrow aspirate and trephine (not routine)
	Liver function
	LDH
Radiology	C-reactive protein
	Chest radiograph
	CT of thorax, abdomen, chest and pelvis
Special tests	MRI
	Bone scan
	PET or PET/CT

CT, computed tomography; ESR, erythrocyte sedimentation rate; LDH, lactate dehydrogenase; MRI, magnetic resonance imaging; PET, positron emission tomography.




Fig. 16.4 Staging of Hodgkin's lymphoma. Stage I indicates node involvement in one lymph node area. Stage II indicates disease involving two or more lymph nodal areas confined to one side of the diaphragm. Stage III indicates disease involving lymph nodes above and below the diaphragm. Splenic disease is included in stage III but this has special significance (see below). Stage IV indicates involvement outside the lymph node areas and refers to diffuse or disseminated disease in the bone marrow, liver and other extranodal sites. NB. The stage number in all cases is followed by the letter A or B indicating the absence (A) or presence (B) of one or more of the

following: unexplained fever above 38°C; night sweats; or loss of more than 10% of body weight within 6 months. Localized extranodal extension from a mass of nodes does not advance the stage but is indicated by the subscript E. Thus, mediastinal disease with contiguous spread to the lung or spinal theca would be classified as I_E. As involvement of the spleen is often a prelude to widespread haematogenous spread of the disease, patients with lymph node and splenic involvement are staged as III_E. Bulky disease (widening of the mediastinum by more than one-third, or the presence of a nodal mass >10 cm in diameter) is relevant to therapy at any stage.




Fig. 16.5 Chest X-ray in Hodgkin's lymphoma showing widespread enlargement of hilar and mediastinal lymph nodes with associated collapse of the right upper lobe and infiltration or possibly pneumonic changes in the mid zone of the left lung.



Fig. 16.6 Hodgkin's lymphoma (nodular sclerosing type): CT scan of chest showing anterior mediastinal mass of enlarged lymph nodes (arrowed).

Patients are also classified as A or B according to whether or not constitutional features (fever or weight loss) are present (Fig. 16.4).

Treatment

Treatment is with radiotherapy, chemotherapy or a combination of both. The choice depends primarily on the stage and whether clinically A or B (Fig. 16.4) although histological grading is an additional factor. Semen storage, if appropriate, should be carried out before therapy is begun.

Radiotherapy

Patients with stage I and IIA disease may be cured by radiotherapy alone. A dose of 4000 rad (40 Gy) is able to destroy lymph node Hodgkin's tissue in approximately 80% of these patients. Improved high-voltage radiotherapy techniques allow the treatment of all lymph node areas above or below the diaphragm by single 'upper mantle' or 'inverted Y' blocks. Radiotherapy also has a role in the treatment of bulky tumour masses such as mediastinal tumour that remains after chemotherapy or painful skeletal, nodal or soft tissue deposits.

Combined modality therapy

Concerns over late relapse and the long-term effects of radiotherapy have led to the development of regimens in which chemotherapy and radiotherapy are used together. This combination therapy allows short courses of chemotherapy to be combined with reduced levels of radiotherapy and the most effective combinations are being assessed in clinical trials.

Chemotherapy

Cyclical chemotherapy is used for stage III and IV disease and also in stage I and II patients who have bulky disease, type B symptoms or have relapsed following initial radiotherapy. The combination of Adriamycin, bleomycin, vinblastine and dacarbazine (ABVD) is now most widely used. This has replaced the landmark quadruple therapy with mustine, vincristine (Oncovin), procarbazine and prednisolone (MOPP) which was introduced in the

1970s, but is more likely to cause sterility or secondary leukaemia. Variants of this such as ChlVPP (replacing mustine with chlorambucil or cyclophosphamide) are sometimes used. It is usual to give six cycles of chemotherapy or four following achievement of complete remission.

More intensive chemotherapy regimens such as Stanford V or BEACOPP (bleomycin, etoposide, doxorubicin, cyclophosphamide, vincristine, procarbazine, prednisolone) may be useful for poor-risk disease and trials to compare these with ABVD are underway.

Assessment of response to treatment

Clinical examination and imaging (e.g. CT scan) are used to assess response to treatment. Patients with Hodgkin's lymphoma often show residual masses following treatment which may be because of the large degree of fibrosis present within lymph nodes. It can be difficult to assess whether or not such masses represent residual disease and the label of complete response uncertain (CRu) can be applied in these cases. PET scanning combined with CT (see p. 211) is useful in revealing areas of active disease and further biopsy may also be required.

Relapsed cases

The patient is treated with an alternative combination chemotherapy to the initial regimen and, if necessary, with radiotherapy to sites of bulky disease. If the disease remains chemosensitive, high-dose chemotherapy and autologous stem cell transplantation improve the probability of cure and are recommended for most patients below the age of 65 years. Allogeneic transplantation may also be tried in a minority of patients who fail other therapies.

Prognosis

Approximate 5-year survival rates range from 50% to over 90% depending on age, stage and histology.

The late effects of Hodgkin's lymphoma and its treatment

Long-term follow-up of patients has revealed a considerable burden of late disease following treatment. Secondary cancers such as lung cancer and breast cancer appear to be related to radiotherapy whereas myelodysplasia or acute myeloid leukaemia are more associated with the use of alkylating agents. Non-Hodgkin's lymphomas and other cancers also occur with greater frequency than in controls. Non-malignant complications include sterility, intestinal complications, myocardial infarction and other cardiac or pulmonary complications of the mediastinal radiation and chemotherapy. These features are the main reason why less intensive treatment regimens are now being explored for this disease.

Bibliography

- Burton C., Eli P., Linch D. (2004) The role of PET imaging in lymphoma. *Br J Haematol* 126,772–84.
- Cartwright R.A. and Watkins G. (2004) Epidemiology of Hodgkin's disease: a review. *Hematol Oncol* 22,11–26.
- Diehl V., Thomas R.K., Re D. (2004) Part II: Hodgkin's lymphoma – diagnosis and treatment. *Lancet Oncol* 5,19–26.
- Harris N.L., Jaffe E.S., Diebold J. *et al.* (2000) The WHO classification of neoplasms of the hematopoietic and lymphoid tissues. *Hematol J* 1,53–66.
- Juweid M.E. and Cheson B.D. (2006) Positron-emission tomography and assessment of cancer therapy. *N Engl J Med* 354,496–507.
- Re D., Thomas R.K., Behringer K., Diehl V. (2005) From Hodgkin disease to Hodgkin lymphoma: biologic insights and therapeutic potential. *Blood* 105,4553–60.
- Thomas R.K., Re D., Wolf J., Diehl V. (2004) Part I. Hodgkin's lymphoma: molecular biology of Hodgkin and Reed–Sternberg cells. *Lancet Oncol* 5,11–8.
- Zekri J.M., Mouncey P., Hancock B.W. (2004) Trials in advanced Hodgkin's disease: more than 30 years experience of the British National Lymphoma Investigation. *Clin Lymphoma* 5,174–83.

Non-Hodgkin's lymphomas, 203

Clinical features of non-Hodgkin's lymphomas, 203

Specific subtypes of non-Hodgkin's lymphoma, 209

T-cell lymphomas, 214

Bibliography, 215

Non-Hodgkin's lymphomas

These are a large group of lymphoid tumours, most commonly of B-cell origin, whose clinical presentation and natural history are more variable than in Hodgkin's lymphoma. They are characterized by an irregular pattern of spread and a significant proportion of patients develop extranodal disease. Their frequency has increased markedly over the last 50 years and with an incidence of approximately 17 in 100 000 they now represent the fifth most common malignancy in some developed countries. The aetiology of the majority of cases of non-Hodgkin's lymphomas (NHL) is unknown although infectious agents are an important cause in particular subtypes (Chapter 10).

Classification and histopathology

No area of diagnostic histopathology has been associated with greater confusion than the classification of NHL. A number of classifications have been used over the years with no one scheme receiving unanimous support. In 1994, the Revised American European Lymphoma (REAL) classification was published and gained widespread application. The more recent World Health Organization (WHO) classification, based on REAL, includes all lymphoid malignancies as well as lymphomas and is more clinically based than previous classification schemes (Table 17.1). In general terms, there is a

move away from subdividing lymphomas on the basis of subtle histological appearance and more in terms of distinct syndromes each with characteristic morphological, immunophenotypic, genetic and clinical features. It is also useful to think of the likely origin of individual lymphoid malignancies based on their phenotype and immunoglobulin gene rearrangement status (Fig. 17.1). In this chapter we consider each of the common lymphoma subtypes within this classification.

Low- and high-grade non-Hodgkin's lymphomas

The non-Hodgkin's lymphomas are a diverse group of diseases and vary from highly proliferative and rapidly fatal diseases to some of the most indolent and well-tolerated malignancies. For many years clinicians have subdivided lymphomas into low- and high-grade disease, with some falling into an intermediate grade. This approach has been extremely valuable as, in general terms, the low-grade disorders are relatively indolent, respond well to chemotherapy but are very difficult to cure whereas high-grade lymphomas are aggressive and need urgent treatment but are often curable.

Clinical features of non-Hodgkin's lymphomas

1 *Superficial lymphadenopathy* The majority of patients present with asymmetric painless enlargement

Table 17.1 The World Health Organization (WHO) classification of non-Hodgkin's lymphomas. B-cell disorders comprise of 85% of cases. T cell and NK cell together comprise of 15% of cases.

<i>Precursor B-cell neoplasms</i>	<i>Precursor T-cell neoplasms</i>
B-lymphoblastic leukaemia/lymphoma (precursor B-cell ALL, B-ALL/LBL)	T-cell lymphoblastic lymphoma/leukaemia (T-ALL/LBL)
<i>Mature B-cell neoplasms</i>	<i>Mature T-cell and NK cell neoplasms</i>
B-cell chronic lymphocytic leukaemia/small lymphocytic lymphoma	T-cell prolymphocytic leukaemia
B-cell prolymphocytic leukaemia	T-cell granular lymphocytic leukaemia
Lymphoplasmacytic lymphoma	Aggressive NK-cell leukaemia
Splenic marginal zone B-cell lymphoma (\pm villous lymphocytes)	Adult T-cell lymphoma/leukaemia (HTLV-1+)
Hairy cell leukaemia	Extranodal NK/T-cell lymphoma, nasal type
Plasma cell myeloma/plasmacytoma	Enteropathy-type T-cell lymphoma
	Mycosis fungoides/Sézary syndrome
	Anaplastic large cell lymphoma, primary cutaneous type
Extranodal marginal zone B-cell lymphoma of MALT type	
	Peripheral T-cell lymphoma, unspecified
Mantle cell lymphoma	Angioimmunoblastic T-cell lymphoma
Follicular lymphoma	Anaplastic large cell lymphoma, primary systemic type
Nodal marginal zone B-cell lymphoma	
Diffuse large B-cell lymphoma	
Burkitt's lymphoma/Burkitt's cell leukaemia	
Primary effusion lymphoma	
Mediastinal large B-cell lymphoma	

ALL, acute lymphoblastic leukaemia; HTLV, human T-cell leukaemia/lymphoma virus; LBL, lymphoblastic lymphoma; MALT, mucosa-associated lymphoid tissue; NK, natural killer.

of lymph nodes in one or more peripheral lymph node regions.

2 Constitutional symptoms Fever, night sweats and weight loss occur less frequently than in Hodgkin's disease and their presence is usually associated with disseminated disease.

3 Oropharyngeal involvement In 5–10% of patients there is disease of the oropharyngeal lymphoid structures (Waldeyer's ring) which may cause complaints of a 'sore throat' or noisy or obstructed breathing.

4 Presenting features Anaemia, neutropenia with infections or thrombocytopenia with purpura may be presenting features in patients with diffuse bone marrow disease. Cytopenias may also be autoimmune in origin. Infections may occur as a result of neutropenia or reduced cell immunity (e.g. herpes zoster).

5 Abdominal disease The liver and spleen are often

enlarged and involvement of retroperitoneal or mesenteric nodes is frequent. The gastrointestinal tract is the most commonly involved extranodal site after the bone marrow, and patients may present with acute abdominal symptoms.

6 Other organs Skin, brain, testis or thyroid involvement is not infrequent. The skin is also primarily involved in two unusual, closely related T-cell lymphomas: mycosis fungoides and Sézary syndrome.

Investigations

Histology

Lymph node biopsy is the definitive investigation (Figs 17.2 and 17.3) and morphological examination is assisted by immunophenotypic and genetic analysis. For B-cell lymphomas, expression of either κ or λ light chains confirms clonality and distinguishes the disease from a reactive node (Fig. 17.4; Chapter 10).






Fig. 17.1 Proposed cellular origin of B-lymphoid malignancies. Normal B cells migrate from the bone marrow and enter secondary lymphoid tissue. When they encounter antigen a germinal centre is formed and B cells undergo somatic hypermutation of the immunoglobulin genes. Finally, B cells exit the lymph node as memory B cells or plasma cells. The cellular origin of the different lymphoid malignancies can be inferred from


immunoglobulin gene rearrangement status and membrane phenotype. Mantle cell lymphoma and a proportion of B-CLL cases have unmutated immunoglobulin genes whereas marginal zone lymphoma, diffuse large cell lymphoma, follicle cell lymphoma, lymphoplasmacytoid lymphoma and some B-CLL cases have mutated immunoglobulin genes.

Fig. 17.2 Non-Hodgkin's lymphoma: histological sections of lymph nodes showing: (a) a diffuse pattern of involvement in lymphocytic lymphoma with the normal architecture totally replaced by neoplastic lymphocytic cells; (b) a follicular or nodular pattern in follicular lymphoma—the 'follicles' or 'nodules' of neoplastic cells compress surrounding tissue and lack a mantle of small lymphocytes.







(a)



(b)



(c)



(d)

Fig. 17.3 Non-Hodgkin's lymphoma: high power view of lymph node biopsies showing (a) lymphocytic lymphoma showing predominantly small lymphocytes with round nuclei containing densely clumped heterochromatin. (b) Mantle cell lymphoma: showing characteristic deformed pattern of small lymphocytes with angular nuclei ('centrocytes'). (c) Diffuse large B-cell lymphoma: the neoplastic cells are much larger than normal

lymphocytes and have a round nucleus with prominent nucleoli, many of which are adjacent to the nuclear membrane ('centroblasts'). A number of mitotic figures are seen. (d) Diffuse large B-cell lymphoma showing large neoplastic cells with a single prominent nucleolus and abundant dark-staining cytoplasm (previously termed immunoblasts).

The repertoire of CD antigens expressed on the tumour cell surface is also useful in the classification of malignant lymphoma (Table 17.2; see Appendix 1). Trephine biopsy of the marrow and immunoglobulin or T-cell receptor gene rearrangement studies may also be valuable.

Haematological and biochemical findings

1 A normochromic, normocytic anaemia is usual but autoimmune haemolytic anaemia may also occur.

2 In advanced disease with marrow involvement there may be neutropenia, thrombocytopenia (especially if the spleen is enlarged) or leucoerythroblastic features.

3 Lymphoma cells (e.g. mantle zone cells, 'cleaved follicular lymphoma' or 'blast' cells) with variable nuclear abnormalities may be found in the peripheral blood in some patients (Fig. 15.7).

4 Trephine biopsy of marrow is valuable (Fig. 17.6). Paradoxically, bone marrow involvement is found more frequently in low-grade malignant lymphomas.

5 The serum lactate dehydrogenase (LDH) level is




Fig. 17.4 Non-Hodgkin's lymphoma: lymph node stained by immunoperoxidase shows (a) brown ring staining for κ in the malignant lymphoid nodule, and (b) no labelling for λ confirming the monoclonal origin of the lymphoma.

Table 17.2 Characteristic immunophenotype and cytogenetics of B-cell lymphomas.

Lymphoma	Surface immunoglobulin	CD5	CD10	CD20	CD23	CD43	Cyclin D ₁	Typical cytogenetics
CLL/lymphocytic lymphoma	Weak	+	–	+	+	+	–	13q14, 17p or 11q deletions; trisomy 12
Follicular lymphoma	+	–	+	+	+/-	–	–	t(14; 18)
Mantle cell lymphoma	+	+	–	+	–	+	+	t(11; 14)
Diffuse large cell lymphoma	+/-	–	+/-	+	–	+/-	–	various
Burkitt's lymphoma	+	–	+	+	–	–	–	t(8; 14)
MALT lymphoma	+	–	–	+	+/-	+/-	–	t(1; 14)

CLL, chronic lymphocytic leukaemia; MALT, mucosa-associated lymphoid tissue. More detailed cytogenetics are given in Table 10.2.

raised in more rapidly proliferating and extensive disease and may be used as a prognostic marker (Table 17.3). Elevation of serum uric acid may occur. 6 Immunoglobulin electrophoresis may reveal a paraprotein.


Cytogenetics

The various subtypes of NHL are associated with characteristic chromosomal translocations which are of diagnostic and prognostic value (see Tables 10.2; 17.2). Particularly characteristic translocations are


Table 17.3 International prognostic index for high-grade lymphoma.

	Good	Bad
Age	<60 years	>60 years
Performance status	0 or 1	>2
Stage (p. 200)	I or II	III or IV
Number of extranodal sites	0 or 1	>2
Serum LDH	Normal	Raised

LDH, lactate dehydrogenase.



(a)



(b)

Fig. 17.5 (a) DNA microarray of diffuse large B-cell lymphoma (DLCL) which uses expression analysis of 27 genes to divide cases into those with a typical activated B cell (ABC) or germinal centre B cell (GCB) pattern (Wright *et al.*, 2003). (b) Survival of patients with DLCL grouped according to an activated B cell (ABC) or germinal centre B cell (GCB) microarray expression pattern. (After Wright *et al.* 2003)

t(8; 14) in Burkitt's lymphoma, t(14; 18) in follicular lymphoma, t(11; 14) in mantle cell lymphoma and t(2; 5) in anaplastic large cell lymphoma.

In B-cell lymphomas the immunoglobulin genes are clonally rearranged whereas in T-cell lymphomas there is clonal rearrangement of the T-cell receptor genes (Chapter 10). DNA microarray patterns of gene expression have been shown to give valuable diagnostic and prognostic information (Fig. 17.5) but are not yet widely available.

Staging

The staging system is the same as that described for Hodgkin's disease but is less clearly related to prognosis than histological type. Staging procedures usually include chest X-ray, computed tomography (CT) scanning or magnetic resonance imaging (MRI) (Fig. 17.7) and bone marrow aspiration and trephine (Fig. 17.6). Positron-emission tomography (PET) may detect disease not seen on




Fig. 17.6 Iliac crest trephine biopsy in lymphocytic lymphoma. Prominent nodules of lymphoid tissue are seen in the intertrabecular space and paratrabecular areas.

CT scan and is useful for following treatment response (Fig. 17.7). Combined PET-CT is now commonly used in initial staging and for detection of residual disease (Fig. 17.8).

Specific subtypes of non-Hodgkin's lymphoma


Low-grade non-Hodgkin's lymphoma

Follicular lymphoma


This is the most common form of NHL and is associated with the t(14; 18) translocation and constitutive BCL-2 expression in the great majority of cases (Fig. 17.9). Patients are likely to be middle-aged or elderly and their disease is often characterized by a benign course for many years. The median survival from diagnosis is approximately 10 years.

Presentation is usually with painless lymphadenopathy, often widespread, and the majority of patients will have stage III or IV disease. However, sudden transformation may occur to aggressive diffuse tumours which are sometimes associated with a leukaemic phase.

Patients with localized disease may achieve cure with radiotherapy alone. Treatment options for disseminated (stage II–IV) disease range from simple observation through oral chemotherapy to experimental high-dose treatment with stem cell support. When used as initial treatment, chlorambucil or cyclophosphamide, vincristine and prednisolone



(a)



(b)

Fig. 17.7 Non-Hodgkin's lymphoma. (a) Computed tomography (CT) scan of the abdomen showing enlarged mesenteric (M) and retroperitoneal (RP; para-aortic) lymph nodes; B, bowel. (Courtesy of Dr L. Berger) (b) CT scan of the abdomen: enlarged retroperitoneal and mesenteric nodes from a man causing the 'floating aorta' (arrowed) appearance. (Courtesy of Professor A. Dixon and Dr R.E. Marcus)

(Continued)




Fig. 17.7 (Continued) (c) Magnetic resonance imaging (MRI) scan of the chest showing large mediastinal lymph nodes (white and arrowed) adjacent to the great vessels (black). (d) MRI T₂-weighted midline sagittal image of a lumbosacral spine showing compression of the dural sac by an extradural mass. A, spinal cord; B, extradural mass; C, roots of corda equina. (Courtesy of Dr A. Valentine) (e) Positron emission tomography (PET) body scan of a 59-year-old woman with high-grade non-Hodgkin's lymphoma. (i) The first scan showed no evidence of disease prior to allogeneic transplant. Normal

physiological uptake is seen in the brain and bladder. Two months post-transplant the patient relapsed clinically with a mass on the anterior chest wall. (ii) The PET scan showed evidence of widespread relapse in nodal (para-aortic and iliac nodes) and extranodal sites including the lung and bone. The uptake in bone is clearly demonstrated in the left humerus and femur (arrowed). This scan illustrates how well PET can detect both nodal and extranodal disease and allows whole body assessment at a single scanning session. (Courtesy of Dr S.F. Barrington)

(CVP) achieve a response in up to 90% of patients with a median duration of approximately 2 years. Fludarabine alone or in combination with cyclophosphamide or mitoxantrone and dexamethasone (FMD) is another option. Rituximab, a humanized monoclonal antibody to CD20, is also effective and may be used alone or combined with chemotherapy (e.g. R-CVP; Fig. 17.10). Early studies suggest it is also valuable in maintenance.

Disease relapse is initially treated with similar

chemotherapy regimens but the disease becomes increasingly difficult to control and more intensive chemotherapy and radiolabelled antibody therapy may be considered. Autologous stem cell transplantation (SCT) may be valuable in patients with a history of at least one relapse and allogeneic SCT offers the prospect of cure for younger patients.

Lymphocytic lymphomas

Lymphocytic lymphomas are closely related to




Fig. 17.8 Non-Hodgkin's lymphoma. PET/CT scan of male aged 26 years. Red arrow shows the level at which transaxial section is performed. Upper right-hand panel showing FDG uptake in anterior mediastinal mass and right hilar node. Upper left-hand panel shows corresponding CT section, bottom left-hand panel shows

the CT and PET fused image of the corresponding section. Bottom right-hand panel, coronal section showing FDG uptake in mediastinal mass and right hilar node. There is also FDG uptake in the thyroid, heart, bowel, kidneys and bladder. (Courtesy of the Department of Nuclear Medicine, University College, London)

chronic lymphocytic leukaemia (CLL) and this lymphoma appears to be a tissue phase of CLL (Figs 17.2 and 17.3). Many patients with this condition are elderly with slowly progressive disease and may not require treatment for extended periods. Treatment is based on that used for CLL (p. 190).

Lymphoplasmacytoid lymphomas

Lymphoplasmacytoid lymphomas are often associated with the production of monoclonal immunoglobulin M (IgM), in which case they tend to be

termed Waldenström's macroglobulinaemia (p. 223). Complications are anaemia and hyperviscosity syndrome. Treatment is with oral chlorambucil, fludarabine or monoclonal antibodies (e.g. rituximab). Plasma exchanges may be needed to reduce hyperviscosity.

Mantle cell lymphoma

Mantle cell lymphoma is derived from pre-germinal centre cells localized in the primary follicles or in the mantle region of secondary follicles. It has a




Fig. 17.9 Follicular lymphoma: immunohistological detection of BCL-2 protein. (a) Follicular lymphoma positive because the BCL-2 is activated by the (14; 18) translocation; (b) reactive node shows germinal centres unstained for BCL-2 surrounded by positive mantle zone B and T cells. Immunoalkaline phosphatase (APAPP) stain. (Courtesy of Professors K.C. Gatter and D.Y. Mason)




Fig. 17.10 The potential mechanisms of action of rituximab. Rituximab binds to CD20 on the surface of B cells. It can elicit a number of effector mechanisms including: (a) antibody dependent cell-mediated cytotoxicity; (b) complement mediated lysis of tumour cells; and (c) direct apoptosis of the target cell.

characteristic phenotype of CD19⁺ and CD5⁺ (like CLL) but in contrast is CD22⁺, CD23⁻. A specific t(11; 14) (q13; q32) translocation is seen which leads to deregulation of the cyclin D₁ (*BCL-1*) gene by juxtaposing it with the immunoglobulin heavy-chain gene. Clinical presentation is typically with lymphadenopathy and often there is bone marrow

infiltration and tumour cells in the blood. The cells have characteristically angular nuclei in histological sections (Fig. 17.3b) and may circulate in the blood (see Fig. 15.6). Despite its low-grade status, mantle cell lymphoma is characterized by a poor prognosis. Current treatment regimens such as CVP, cyclophosphamide, hydroxodaunorubicin, oncovin (vincristine) and prednisolone (CHOP) or fludarabine, each with rituximab, are used. The role of bortezomid (see p. 223) or autologous stem cell transplantation is being investigated. The prognosis is poor and the median survival is approximately 3 years.

Marginal zone lymphomas

Marginal zone lymphomas are typically extranodal and are usually localized. Mucosa-associated lymphoid tissue (MALT) lymphomas come into this category and usually arise as a consequence of a pre-existing inflammatory or autoimmune disorder at sites such as the stomach or thyroid. Gastric MALT lymphoma is the most common form and is preceded by *Helicobacter (H.) pylori* infection. In the early stages it may respond to antibiotic therapy aimed at eliminating *H. pylori*. Chemotherapy and/or radiotherapy combined with rituximab as for other low-grade lymphoma may be needed.

Splenic marginal zone lymphoma usually presents as splenomegaly and may be associated with circulating 'villous' lymphocytes (p. 194). Splenectomy is useful for symptomatic patients.

High-grade non-Hodgkin's lymphoma

Diffuse large B-cell lymphomas

Diffuse large B-cell lymphomas (DLCL) are a heterogeneous group of disorders representing the classical 'high-grade' lymphomas. As such they typically present with rapidly progressive lymphadenopathy associated with a fast rate of cellular proliferation. Progressive infiltration may affect the bone marrow, gastrointestinal tract, the spinal cord, the kidneys or other organs.

A variety of clinical and laboratory findings are relevant to the outcome of therapy. According to the international prognostic index these include age, performance status, stage, number of extranodal sites and serum LDH (Table 17.3). Bulky disease (major mass >5 cm in diameter) and prior history of low-grade disease or AIDS are also associated with a poorer prognosis. The cell of origin of DLCL has recently been suggested to be of prognostic significance. If this is germinal centre the outlook has been found to be more favourable than if the origin is from an activated peripheral B cell. Cases associated with 3q27 translocation also have a relatively good prognosis. DNA microarray analysis can give prognostic information (Fig. 17.5).

The CHOP regimen has been the mainstay of treatment of DLCL for many years and consists of cyclophosphamide, hydroxycarbamide, vincristine (Oncovin) and prednisolone given in 2-, 3- or 4-weekly cycles, typically for six to eight courses. The addition of anti-CD20 (rituximab) improves the cure rate and so R-CHOP is now the initial treatment. Cycles given at 14-day intervals with G-CSF support are under trial in younger patients. For localized disease, combined radiotherapy and chemotherapy (e.g. three courses of R-CHOP) may be optimal. Prophylactic therapy to prevent central nervous system (CNS) disease such as intrathecal or high-dose systemic methotrexate should be considered for patients with high-risk disease, particularly those with bone marrow involvement.

For patients who relapse and have chemotherapy-sensitive disease, high-dose chemotherapy with ifosfamide, epirubicin and etoposide (IVE) or etoposide, cytosine arabinoside, methylprednisolone and cisplatin (ESHAP), followed by autologous SCT can be effective. For those with primary refractory

or chemoresistant disease the outlook is poor. Bortezomib, a proteasome inhibitor is undergoing trials. Alemtuzumab directed against CD52 may be used in both B- and T-cell NHL. Anti-CD20 radioactively linked to yttrium-90 or iodine-131 is currently reserved for relapsed cases. Overall long-term survival is approximately 65%.

Burkitt's lymphoma

Burkitt's lymphoma is the lymphomatous correlate of L₃ acute lymphoblastic leukaemia (p. 159) and occurs in endemic or sporadic forms. Endemic (African) Burkitt's lymphoma is seen in areas with chronic malaria exposure and is associated with Epstein-Barr virus (EBV) infection. In virtually all cases the C-MYC oncogene is overexpressed because it is translocated to an immunoglobulin gene, usually the heavy-chain locus t(8; 14). Typically, the patient, usually a child, presents with massive lymphadenopathy of the jaw (Fig. 17.11) which is initially very responsive to chemotherapy although long-term cure is uncommon. Sporadic Burkitt's lymphoma may occur anywhere in the world and is not associated with EBV infection. The histological picture of Burkitt's lymphoma is distinctive (Fig. 17.12). The prognosis for such patients was poor until the introduction of chemotherapy



Fig. 17.11 Burkitt's lymphoma: characteristic facial swelling caused by extensive tumour involvement of the mandible and surrounding soft tissues.




Fig. 17.12 Burkitt's lymphoma: histological section of lymph node showing sheets of lymphoblasts and 'starry sky' tingible body macrophages.

regimes which include high-dose methotrexate and cyclophosphamide. These have transformed the outlook and now the majority of patients are cured. B-ALL is treated with the same regimes.

Lymphoblastic lymphomas

Lymphoblastic lymphomas occur mainly in children and young adults and these conditions merge clinically and morphologically with acute lymphoblastic leukaemia (ALL). They are treated in a similar way.

T-cell lymphomas

Peripheral T-cell lymphomas that present with lymphadenopathy rather than extranodal disease are a heterogeneous group of rare tumours and are usually of CD4⁺ phenotype. Several variants of T-cell lymphomas are recognized.

Peripheral T-cell non-Hodgkin's lymphoma, unspecified. These derive from post-thymic T cells at various stages of differentiation. They are treated with combined chemotherapy (e.g. CHOP). The prognosis is poor.

Angioimmunoblastic lymphadenopathy usually occurs in elderly patients with lymphadenopathy, hepatosplenomegaly, skin rashes and a polyclonal increase in serum IgG.



Fig. 17.13 Mycosis fungoides.

Mycosis fungoides is a chronic cutaneous T-cell lymphoma that presents with severe pruritus and psoriasis-like lesions (Fig. 17.13). Ultimately, deeper organs are affected, particularly lymph nodes, spleen, liver and bone marrow.

In **Sézary syndrome** there is dermatitis, erythroderma, generalized lymphadenopathy and circulating T-lymphoma cells. The cells are usually CD4⁺ and have a folded or cerebriform nuclear chromatin. Initial treatment of these conditions is by local irradiation, topical chemotherapy or photochemotherapy with psoralen and ultraviolet light (PUVA). Chemotherapy (e.g. CHOP) may be needed.

Adult T-cell leukaemia/lymphoma is associated with human T-cell leukaemia/lymphoma virus type 1 (HTLV-1) infection and presents with lymphadenopathy, hepatic and splenic enlargement, cutaneous infiltrations and hypercalcaemia.

Angiocentric lymphomas typically involve the nasal sinuses whereas T-cell intestinal lymphoma is associated with gluten-induced enteropathy in many cases.

Anaplastic large cell lymphoma is particularly common in children and is of T-cell or null cell

phenotype. The disease is CD30⁺ and associated with the t(2; 5) (p23; q35) translocation. It has an aggressive course characterized by systemic symptoms and extranodal involvement.

Bibliography

- Abramson J.S. and Shipp M.A. (2005) Advances in the biology and therapy of diffuse large B-cell lymphoma: moving toward a molecularly targeted approach. *Blood* 106,1164–74.
- Barosi G., Carella A., Lazzarino M. *et al.* (2005) Management of nodal indolent (non marginal-zone) non-Hodgkin's lymphomas: practice guidelines from the Italian Society of Hematology, Italian Society of Experimental Hematology and Italian Group for Bone Marrow Transplantation. *Haematologica* 90,1236–57.
- Coiffier B. (2005) Current strategies for the treatment of diffuse large B cell lymphoma. *Curr Opin Hematol* 12,259–65.
- Fisher S.G. and Fisher R.I. (2004) The epidemiology of non-Hodgkin's lymphoma. *Oncogene* 23,6524–34.
- Gandhi M.K. and Marcus R.E. (2005) Follicular lymphoma: time for a re-think? *Blood Rev* 19,165–78.
- Hennessy B.T., Hanrahan E.O., Daly P.A. (2004) Non-Hodgkin lymphoma: an update. *Lancet Oncol* 6,341–53.
- Lossos I.S., Czerwinski D.K., Alizadeh A.A. *et al.* (2004) Prediction of survival in diffuse large-B-cell lymphoma based on the expression of six genes. *N Engl J Med* 350,1828–37.
- Mauch P.M., Armitage J.O., Coiffier B., Dalla-Favera R., Harris N.L. (eds) (2004) *Non-Hodgkin's Lymphomas*. Lippincott Williams and Wilkins, Philadelphia.
- Monti S., Savage K.J., Kutok J.L. *et al.* (2005) Molecular profiling of diffuse large B-cell lymphoma identifies robust subtypes including one characterized by host inflammatory response. *Blood* 105,1851–61.
- Peggs K.S., Mackinnon S., Linch D.C. (2005) The role of allogeneic transplantation in non-Hodgkin's lymphoma. *Br J Haematol* 128,153–68.
- Rastetter W., Molina A., White C.A. (2004) Rituximab: expanding role in therapy for lymphomas and autoimmune diseases. *Annu Rev Med* 55,477–503.
- Reichert J.M., Rosensweig C.J., Faden L.B., Dewitz M.C. (2005) Monoclonal antibody successes in the clinic. *Nat Biotechnol* 23,1073–8.
- Rizvi M.A., Evens A.M., Tallman M.S., Nelson B.P., Rosen S.T. (2006) T-cell non-Hodgkin lymphoma. *Blood* 107,1255–64.
- Seyfarth B., Josting A., Dreyling M., Schmitz N. (2006) Relapse in common lymphoma subtypes: salvage treatment options for follicular lymphoma and Hodgkin's disease. *Br J Haematol* 133,3–18.
- Wright G., Tan B., Rosenwald A. *et al.* (2003) A gene-expression-based method to diagnose clinically distinct subgroups of diffuse large B-cell lymphoma. *Proc Natl Acad Sci USA* 100,9991–6.

CHAPTER 18

18

Multiple myeloma and related disorders

Paraproteinaemia, 216
Multiple myeloma, 216
Other plasma cell tumours, 223
Waldenström's macroglobulinaemia, 223

Monoclonal gammopathy of undetermined significance, 224
Amyloidosis, 225
Hyperviscosity syndrome, 226
Bibliography, 228

Paraproteinaemia

This term refers to the presence of a monoclonal immunoglobulin band in the serum. Normally, serum immunoglobulins are polyclonal and represent the combined output from millions of different plasma cells. A monoclonal band (M-protein), or *paraprotein*, reflects the synthesis of immunoglobulin from a single clone of plasma cells. There are several situations in which this may occur (Table 18.1) and not all require treatment.

Table 18.1 Diseases associated with M-proteins.

<i>Malignant or uncontrolled production</i>
Multiple myeloma
Waldenström's macroglobulinaemia
Malignant lymphoma
Chronic lymphocytic leukaemia
Primary amyloidosis
Plasma cell leukaemia
Heavy chain disease
<i>Benign or stable production</i>
Benign monoclonal gammopathy
Solitary plasmacytoma
Chronic cold haemagglutinin disease
Transient (e.g. with infections)
Acquired immune deficiency syndrome (AIDS)
Gaucher's disease
Rarely with carcinoma and other conditions

Multiple myeloma

Multiple myeloma (myelomatosis) is a neoplastic proliferation characterized by plasma cell accumulation in the bone marrow, the presence of monoclonal protein in the serum and/or urine and related tissue damage. Ninety-eight per cent of cases occur over the age of 40 years with a peak incidence in the seventh decade.

The myeloma cell is a post-germinal centre plasma cell that has undergone immunoglobulin class switching and somatic hypermutation and secretes the paraprotein that is present in serum. Plasma cells naturally home to the bone marrow and this characteristic is retained by the tumour cell. The aetiology of the disease is unknown but it is more common in certain racial groups such as those of Afro-Caribbean origin. Tumour cells have very complex genetic changes but dysregulated or increased expression of cyclin D (p. 9) is believed to be an early unifying event. Approximately half of the tumours have an increased number of chromosomes (*hyperdiploid*) whereas non-hyperdiploid cases have a high incidence of translocations involving the immunoglobulin heavy-chain gene (*IGH*). Monoallelic loss of 13q is frequent in both categories and all these genetic abnormalities are also seen in monoclonal gammopathy of undetermined significance (MGUS; p. 224). The characteristic immunophenotype of malignant plasma cells is CD38^{high},

Table 18.2 Features of benign and malignant paraproteinaemia.

	Benign	Malignant
Bence-Jones proteinuria	Absent	May be present
Serum paraprotein concentration	Usually <20 g/L and stationary	Usually >20 g/L and rising
Serum free light chain ratio	Normal	Abnormal
Immunoparesis (hypogammaglobinaemia)	Absent	Present
Underlying lymphoproliferative disease or myeloma	Absent	Present
Bone lesions	Absent	Present
Plasma cells in marrow	<10%	>10%

CD138^{high} and CD45^{low}. Interleukin 6 is a potent growth factor for myeloma cells and is often active by an autocrine mechanism.

Diagnosis

This depends on three principal findings (Table 18.2):
1 Monoclonal protein in serum and/or urine (Fig. 18.1).




Fig. 18.1 Serum protein electrophoresis in multiple myeloma showing an abnormal paraprotein in the γ -globulin region with reduced levels of background β - and γ -globulins.

2 Increased plasma cells in the bone marrow (Fig. 18.2).

3 Related organ or tissue impairment such as bone disease, renal impairment, anaemia, hypercalcaemia, hyperviscosity, amyloidosis or recurrent infection.

If the bone marrow plasma cell count is >10% but there is **no** evidence of tissue damage the disease is termed *asymptomatic* or *smouldering myeloma*.

Clinical features

1 Bone pain (especially backache) resulting from vertebral collapse and pathological fractures (Fig. 18.3a,b).

2 Features of anaemia: e.g. lethargy, weakness, dyspnoea, pallor, tachycardia.

3 Recurrent infections: related to deficient antibody production, abnormal cell-mediated immunity and neutropenia.

4 Features of renal failure and/or hypercalcaemia: polydipsia, polyuria, anorexia, vomiting, constipation and mental disturbance.

5 Abnormal bleeding tendency: myeloma protein may interfere with platelet function and coagulation factors; thrombocytopenia occurs in advanced disease.

6 Amyloidosis occurs in 5% with features such as macroglossia, carpal tunnel syndrome and diarrhoea.




Fig. 18.2 The bone marrow in multiple myeloma showing large numbers of plasma cells, with many abnormal forms.

7 In approximately 2% of cases there is a hyperviscosity syndrome (Fig. 22.6) with purpura, haemorrhages, visual failure, central nervous system (CNS) symptoms, neuropathies and heart failure.

Laboratory findings include the following:

1 **Presence of a paraprotein** Serum and urine should be screened by immunoglobulin electrophoresis. The paraprotein is immunoglobulin G (IgG) in 60% of cases, IgA in 20% and light chain only in almost all the rest. Less than 1% have IgD or IgE paraprotein. Free light chains are filtered from the serum into the kidney and are therefore often not detectable in serum using electrophoresis. However, the level of free light chain in serum can be measured with a specific antibody. As free light chains are produced by almost all malignant plasma cells the *serum free light chain* assay is useful in diagnosis and monitoring of myeloma and other forms of paraproteinemia. In myeloma the normal $\kappa : \lambda$ free light chain ratio of 2 : 1 is skewed with an excess of κ or λ chains.

Normal serum immunoglobulin levels (IgG, IgA and IgM) are reduced, a feature known as *immune paresis*. The urine contains free light chains, known as *Bence-Jones protein*, in two-thirds of cases. Rare cases of myeloma are non-secretory and therefore not associated with a paraprotein or Bence-Jones proteinuria although some will still show a disturbed free light chain ratio in the serum.


2 There is usually a normochromic normocytic or macrocytic anaemia. Rouleaux formation is marked in most cases (Fig. 18.4). Neutropenia and thrombocytopenia occur in advanced disease. Abnormal plasma cells appear in the blood film in 15% of patients and can be detected by sensitive flow cytometry in over 50%.

3 High erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP).

4 Increased plasma cells in the bone marrow (usually >20%) often with abnormal forms (Fig. 18.2).

5 Radiological investigation of the skeleton reveals bone lesions such as osteolytic areas without evidence of surrounding osteoblastic reaction or sclerosis in 60% of patients (Fig. 18.5) or generalized osteoporosis in 20% (Fig. 18.3). Twenty per cent have no bone lesions. In addition, pathological fractures or vertebral collapse (Fig. 18.3b) are common. The osteolytic lesions are caused by osteoclast activation resulting from high serum levels of RANKL (receptor activator of nuclear factor- κ B (NF- κ B) ligand), produced by plasma cells and bone marrow stroma, which binds to activatory RANK receptors on the osteoclast surface.

6 Serum calcium elevation occurs in 45% of patients. Typically, the serum alkaline phosphatase is normal (except following pathological fractures).



(a)



(b)

Fig. 18.3 (a) Multiple myeloma: X-ray of lumbar spine showing severe demineralization with partial collapse of L₃. (b) Magnetic resonance imaging (MRI) of spine: T₂-weighted study. There is infiltration and destruction of L₃ and L₅ with bulging of the posterior part of the body of L₃ into the spinal canal compressing the corda equina (arrowed). Radiotherapy has caused a marrow signal change in vertebrae C₂–D₄ because of replacement of normal red marrow by fat (bright white signal). (Courtesy of Dr A. Platts)

7 The serum creatinine is raised in 20% of cases. Proteinaceous deposits from heavy Bence-Jones proteinuria, hypercalcaemia, uric acid, amyloid and pyelonephritis may all contribute to renal failure (Fig. 18.6).

8 A low serum albumin occurs with advanced disease.

9 Serum β_2 -microglobulin is often raised and is a useful indicator of prognosis. Levels less than 4 mg/L imply a relatively good prognosis.



Fig. 18.4 The peripheral blood film in multiple myeloma showing rouleaux formations.




Fig. 18.5 Skull X-ray in multiple myeloma showing many 'punched-out' lesions.

Treatment

This may be divided into specific and supportive (Fig. 18.7).

Specific


At the current time the disease remains incurable except for those very few, mostly younger, patients who may be cured by allogeneic stem cell trans-

plantation (SCT). For all other patients the major treatment decision is between the use of *intensive therapy* (mostly for patients aged less than 65–70 years) and *non-intensive therapy* for older patients.


Intensive therapy involves the combination of several courses of chemotherapy to reduce the tumour burden followed by stem cell collection and autologous SCT after high-dose chemotherapy. Repeated intravenous or oral chemotherapy cycles such as cyclophosphamide, dexamethasone and thalidomide (CDT), cyclophosphamide, vincristine, adriamycin (hydroxodaunorubicin) and dexamethasone (CVAD) or idarubicin with dexamethasone (IDEX) may be used. Peripheral blood stem cells are collected after mobilization using a combination of chemotherapy and granulocyte colony-stimulating factor (G-CSF). High-dose melphalan, with or without radiotherapy, is the typical conditioning regime for autologous SCT. Two consecutive SCT procedures are used in some centres.

Although allogeneic transplantation may cure the disease it carries a high procedure-related mortality and patients frequently relapse after the procedure.


Non-intensive therapy In elderly patients monthly courses of the oral alkylating agent melphalan, sometimes in combination with prednisolone, are usually effective in reducing the tumour burden. Many centres now also use thalidomide (see below).




(a)



(b)



(c)



(d)

Fig. 18.6 The kidney in multiple myeloma. (a) Myeloma kidney: the renal tubules are distended with hyaline protein (precipitated light chains or Bence-Jones protein). Giant cells are prominent in the surrounding cellular reaction. (b) Amyloid deposition: both glomeruli and several of the small blood vessels contain an amorphous

pink-staining deposit characteristic of amyloid (Congo red stain). (c) Nephrocalcinosis: calcium deposition (dark 'fractured' material) in the renal parenchyma. (d) Pyelonephritis: destruction of renal parenchyma and infiltration by acute inflammatory cells.




Fig. 18.7 An approach to the management of multiple myeloma.

Typically, paraprotein levels gradually fall, bone lesions show improvement and blood counts may improve. Cyclophosphamide is also effective and simple to use as a single agent. However, after a variable number of courses a 'plateau phase' is reached in which the paraprotein level stops falling. At this point treatment is stopped and the patient is seen at regular intervals in the outpatient clinic. After a variable period of time, often around 18 months, the disease 'escapes' from plateau with rising paraprotein and worsening symptoms. Further chemotherapy may be given although the disease becomes increasingly difficult to control. Trials of

drugs designed to overcome multidrug resistance have not proven successful.

Other treatments

Thalidomide is useful in the management of relapsed disease and is being introduced into first-line therapy (e.g. CDT). Its precise mechanism of action is unknown and it has a number of side-effects such as sedation, constipation, neuropathy and thrombosis. The addition of dexamethasone increases the response rate but venous thrombosis becomes a major concern. Prophylactic anticoagulation with

warfarin or heparin is needed when thalidomide is used in induction regimes. Analogues of thalidomide which have increased immunomodulatory activity are now showing promise.

Bortezomib (Velcade) is a promising drug which inhibits cellular proteasome and NF- κ B activation. Already proven in refractory disease, it is now being assessed in earlier phases of treatment. Its main side-effect is neuropathy.

α -Interferon This may prolong the plateau phase following chemotherapy or transplantation but has little effect on overall survival.

Radiotherapy is highly effective in treating the symptoms of myeloma. It may be used for areas of bone pain or spinal cord compression.

Support care

Renal failure Rehydrate and treat the underlying cause (e.g. hypercalcaemia, hyperuricaemia). Dialysis is generally well tolerated. It is important that all patients with myeloma drink at least 3 L of fluid each day throughout the course of their disease.

Bone disease and hypercalcaemia Bisphosphonates such as pamidronate, clodronate or zoledronic acid are effective in reducing the progression of bone disease and may also improve overall survival. Acute hypercalcaemia is treated with rehydration with isotonic saline, a diuretic and corticosteroids followed by a bisphosphonate.

Compression paraplegia Use decompression laminectomy or irradiation; corticosteroid therapy may help.

Anaemia Transfusion or erythropoietin are used.

Bleeding Bleeding caused by paraprotein interference with coagulation and hyperviscosity syndrome may be treated by repeated plasmapheresis.

Infections Rapid treatment of any infection is essential. Prophylactic infusions of immunoglobulin concentrates together with oral broad-spectrum

antibiotics and antifungal agents may be needed for recurrent infections.

Prognosis

An international prognostic index has been used based on serum β_2 -microglobulin (β_2 M) and albumin levels. Patients with serum β_2 M >5.5 mg/L and an albumin <35 g/L have a poor survival as do those with frequent circulating plasma cells. Overall, the median survival with non-intensive chemotherapy is 3–4 years and this is improved by approximately 1 year with autologous transplantation.

Other plasma cell tumours

Solitary plasmacytoma

These are isolated plasma cell tumours, usually of bone or soft tissue (e.g. the mucosa of the upper respiratory and gastrointestinal tracts or the skin). The associated paraprotein disappears following radiotherapy to the primary lesion.

Plasma cell leukaemia

This occurs either as a late complication of myeloma or as a primary disease characterized by the presence of 20% or more plasma cells in the blood (Chapter 15). The outlook is poor.

Heavy chain disease

In these rare disorders the neoplastic cells secrete only incomplete immunoglobulin heavy chains (γ , α or μ). The most common form is α -heavy chain disease which occurs mainly in the Mediterranean area and starts as a malabsorption syndrome which may progress to lymphoma.

Waldenström's macroglobulinaemia

This is an uncommon condition, seen most frequently in men over 50 years of age, in which there is a lymphoplasmacytoid lymphoma (Fig. 18.8) which produces a monoclonal IgM paraprotein. The cell of origin appears to be a post-germinal centre B cell with the characteristics of an IgM-bearing memory B cell.




Fig 18.8 Lymphoplasmacytoid lymphoma associated with Waldenström's macroglobulinaemia. Bone marrow shows cells with features of lymphocytes and plasma cells.

Clinical features

These are usually of insidious onset, with fatigue and weight loss. Hyperviscosity syndrome (Fig. 22.6) is common. IgM paraprotein increases blood viscosity more than equivalent concentrations of IgG or IgA, and small increases above 30 g/L in concentration lead to large increases in viscosity. Visual upset is frequent and the retina may show a variety of changes such as engorged veins, haemorrhages, exudates and a blurred disc (Fig. 18.9). If the macroglobulin is a cryoglobulin, features of cryoprecipitation, such as Raynaud's phenomenon, may be present. Anaemia, at least partly caused by an increased plasma volume, is usually a significant problem and a bleeding tendency may result from macroglobulin interference with coagulation factors and platelet function. Neurological symptoms, dyspnoea and heart failure may be presenting symptoms. Moderate lymphadenopathy and enlargement of the liver and spleen are frequently seen.

Diagnosis

This is made by the finding of a monoclonal serum IgM together with bone marrow or lymph node infiltration with lymphoplasmacytoid cells. The ESR is raised and there may be a peripheral blood lymphocytosis.

Treatment

Specific

No therapy is needed for patients without symptoms, significant organomegaly or anaemia. Chlorambucil or cyclophosphamide have been the mainstay of therapy but fludarabine or 2-chlorodeoxyadenosine and rituximab are also useful for initial therapy or for relapsed cases. Combination chemotherapy as for non-Hodgkin's lymphoma is needed in late stages. SCT may be considered for advanced disease.

Support

Acute hyperviscosity syndrome is treated with repeated plasmapheresis. As IgM is mainly intravascular, plasmapheresis is more effective than with IgG or IgA paraproteins when much of the protein is extravascular and so rapidly replenishes the plasma compartment. Erythropoietin or regular transfusions may be required for chronic anaemia.

Monoclonal gammopathy of undetermined significance

Transient or persistent paraproteins can occur in many other conditions as well as in multiple myeloma (Table 18.1). A serum paraprotein may be sometimes be detected without any evidence of myeloma or other underlying disease and is termed





Fig. 18.9 Waldenström's macroglobulinaemia: hyperviscosity syndrome. (a) The retina before plasmapheresis shows distension of retinal vessels, particularly the veins which show bulging and constriction (the 'linked sausage' effect) and areas of haemorrhage; (b) following plasmapheresis the vessels have returned to normal and the areas of haemorrhage have cleared.

monoclonal gammopathy of undetermined significance (MGUS). It becomes increasingly common with age, being present in 1% of persons older than 50 years and 3% of those over 70 years. There are no bone lesions, no Bence-Jones proteinuria, and the proportion of plasma cells in the marrow is normal (<4%) or only slightly raised (<10%) (Table 18.2). The concentration of monoclonal immunoglobulin in serum is usually less than 20 g/L and other serum immunoglobulins are not depressed. Serum light chains (either κ or λ) are increased in one-third of patients; the greater the increase, the more the risk of transformation. DNA microarray patterns can distinguish myeloma from MGUS. No treatment is needed but patients with MGUS develop overt myeloma or lymphoma at a rate of approximately 1% each year and so are usually followed up regularly in the outpatient clinic.


Amyloidosis

The amyloidoses are a heterogeneous group of disorders characterized by the extracellular deposition of protein in an abnormal fibrillar form. Amyloidosis may be hereditary or acquired and deposits may be focal, localized or systemic in distribution. The amyloid is made from different amyloid fibril precursor proteins in each type of disease. Except for intracerebral amyloid plaques, all amyloid deposits contain a non-fibrillary glycoprotein amyloid P which is derived from a normal serum precursor structurally related to CRP. The classic diagnostic histological test is red-green birefringence after staining with Congo red and viewing under polarized light (Fig. 18.10).

Amyloidosis is classified in Table 18.3.



(a)



(b)

Fig. 18.10 Amyloidosis:
(a) Congo red staining; and
(b) blue–green birefringence
under polarized light.

Systemic AL amyloidosis

Systemic amyloid disease is caused by deposition of monoclonal light chains produced from a clonal plasma cell proliferation. The level of paraprotein may be very low and is not always detectable in serum or urine. The clinical features are caused by involvement of the heart, tongue (Fig. 18.11), peripheral nerves and kidneys (Fig. 18.12), and the patient may present with heart failure, macroglos-

sia, peripheral neuropathy, carpal tunnel syndrome or renal failure. Treatment is with chemotherapy similar to that used in myeloma, possibly with autologous SCT, which may improve prognosis.

Hyperviscosity syndrome

The most common cause is polycythaemia (p. 233). Hyperviscosity may also occur in patients with myeloma or Waldenström's macroglobulinaemia

Table 18.3 Classification of amyloidosis: types, structure and organ involvement.

Type	Chemical nature	Organs involved
<i>Systemic AL amyloidosis</i>		
Associated with myeloma, Waldenström's macroglobulinaemia or MGUS	Immunoglobulin light chains (AL)	Tongue Skin Heart Nerves Connective tissue Kidneys Liver Spleen
May also occur on its own as primary amyloidosis (associated with an occult plasma cell proliferation)		
May also occur in localized form with local 'immunocyte' proliferation		
<i>Reactive systemic AA amyloidosis</i>		
Rheumatoid arthritis, tuberculosis, bronchiectasis, chronic osteomyelitis, inflammatory bowel disease	Protein A (AA)	Liver Spleen Kidneys Bone marrow
Hodgkin's lymphoma, carcinomas, familial Mediterranean fever		
<i>Familial amyloidosis</i>	e.g. Transthyretin abnormalities	Nerves Heart Eyes
<i>Localized amyloidosis</i>		
Central nervous system	β -amyloid protein	Alzheimer's disease
Endocrine	Peptic hormones	Endocrine tumours
Senile	Various	Heart, brain, joints, prostate, etc.

AA, AL, these are defined by their chemical nature as in the table; MGUS, monoclonal gammopathy of undetermined significance.



Fig. 18.11 Multiple myeloma: the tongue and lips are enlarged because of nodular and waxy deposits of amyloid.

or in patients with chronic or acute leukaemias associated with very high white cell counts. Rarely, haemophiliac patients with circulating inhibitors, being treated with massive doses of cryoprecipitate, have developed hyperviscosity because of the large volumes of fibrinogen infused.

The clinical features of the hyperviscosity syndrome include visual disturbances, lethargy, confusion, muscle weakness, nervous system symptoms and signs, and congestive heart failure. The retina may show a variety of changes: engorged veins, haemorrhages, exudates and a blurred disc (Fig. 18.9).

Emergency treatment varies with the cause: venesection or isovolaemic exchange with a plasma




Fig. 18.12 Serial anterior whole body ^{123}I -labelled serum amyloid P component (SAP) scans of a 52-year-old woman who presented with renal failure resulting from systemic AL amyloidosis. (a) The initial scan demonstrates a large amyloid load with hepatic, splenic, renal and bone marrow deposits. The underlying plasma cell dyscrasia responded to high-dose melphalan followed by autologous stem cell rescue. (b) Follow-up SAP scintigraphy 3 years after chemotherapy showed greatly reduced uptake of tracer indicating substantial regression of her amyloid deposits. (Courtesy of Professor P.N. Hawkins, National Amyloidosis Centre, Royal Free Hospital, London)

substitute for red cells in a polycythaemic patient; plasmapheresis in myeloma, Waldenström's disease or hyperfibrinogenaemia; and leucopheresis or chemotherapy in leukaemias associated with high white counts. The long-term treatment depends on control of the primary disease with specific therapy.

Bibliography

- Anderson K.C., Pazdur R., Farrell A.T. (2005) Development of effective new treatments for multiple myeloma. *J Clin Oncol* 23,7207–11.
- Bergsagel P.L. and Kuehl W.M. (2005) Molecular pathogenesis and a consequent classification of multiple myeloma. *J Clin Oncol* 23,6333–8.
- Bergsagel P.L., Kuehl W.M., Zhan F. *et al.* (2005) Cyclin D dysregulation: an early and unifying pathogenic event in multiple myeloma. *Blood* 106,296–303.
- Child J.A., Morgan G.J., Davies F.E. *et al.* (2003) Medical Research Council Adult Leukaemia Working Party. High-dose chemotherapy with hematopoietic stem-cell rescue for multiple myeloma. *N Engl J Med* 348,1875–83.
- Gahrton G., Durie B.G.M., Samson D.M. (2004) *Multiple Myeloma and Related Disorders*. Arnold, London.
- Greipp P.R., San Miguel J., Durie B.G. *et al.* (2005) International staging system for multiple myeloma. *J Clin Oncol* 23,3412–20.
- International Myeloma Working Group (2003) Criteria for the classification of monoclonal gammopathies, multiple myeloma and related disorders: a report of the International Myeloma Working Group. *Br J Haematol* 121,749–57.

- Johnson S.A., Birchall J., Luckie C., Oscier D.G., Owen R.G. (2006) Guideline on the management of Waldenström macroglobulinaemia. *Br J Haematol* 132,683–97.
- Kyle R.A. and Rajkumar S.V. (2004) Multiple myeloma. *N Engl J Med* 351,1860–73.
- Kyle R.A., Therneau T.M., Rajkumar S.V. (2006) Prevalence of monoclonal gammopathy of undetermined significance. *N Engl J Med* 354,362–9.
- Kyle R.A., Therneau T.M., Rajkumar S.V. *et al.* (2002) A long-term study of prognosis in monoclonal gammopathy of undetermined significance. *N Engl J Med* 346,564–9.
- Lachmann H.J., Gallimore R., Gillmore J.D. *et al.* (2003) Outcome in systemic AL amyloidosis in relation to changes in concentration of circulating free immunoglobulin light chains following chemotherapy. *Br J Haematol* 122,78–84.
- Nowakowski G.S., Witzig T.E., Dingli D. *et al.* (2005) Circulating plasma cells detected by flow cytometry as a predictor of survival in 302 patients with newly diagnosed multiple myeloma. *Blood* 106,2276–9.
- Palumbo A., Bringhen S., Caravita T. *et al.* (2006) Oral melphelan and prednisone chemotherapy plus thalido-
- midide compared with melphelan and prednisone alone in elderly patients with multiple myeloma: randomised controlled trial. *Lancet* 367,825–31.
- Rajkumar S.V., Kyle R.A., Therneau T.M. *et al.* (2005) Serum free light chain ratio is an independent risk factor for progression in monoclonal gammopathy of undetermined significance. *Blood* 106,812–7.
- Richardson P.G., Barlogie B., Berenson J. *et al.* (2003) A phase 2 study of bortezomib in relapsed, refractory myeloma. *N Engl J Med* 348,2609–17.
- Singal S., Mehta J., Desikan R. (1999) Antitumour activity of thalidomide in refractory myeloma. *N Engl J Med* 341,1565–71.
- Smith A., Wisloff F., Samson D., UK Myeloma Forum; Nordic Myeloma Study Group; British Committee for Standards in Haematology (2006) Guidelines on the diagnosis and management of multiple myeloma. *Br J Haematol* 132,410–51.
- Treon S.P., Gertz M.A., Dimopoulos M. *et al.* (2006) Update on treatment recommendations from the Third International Workshop on Waldenström’s macroglobulinemia. *Blood* 107,3442–6.

CHAPTER 19

19

Myeloproliferative disorders

Polycythaemia, 232
 Polycythaemia (rubra) vera, 232
 Primary familial (congenital) polycythaemia, 236
 Secondary polycythaemia, 236
 Apparent polycythaemia, 236

Differential diagnosis of polycythaemia, 236
 Essential thrombocythaemia, 237
 Myelofibrosis, 238
 Bibliography, 240

The term myeloproliferative disorders describes a group of conditions arising from marrow stem cells and characterized by clonal proliferation of one or more haemopoietic components in the bone marrow and, in many cases, the liver and spleen. Three disorders are included in this classification:

- 1 Polycythaemia rubra vera (PRV).
- 2 Essential thrombocythaemia (ET).
- 3 Myelofibrosis.

These disorders are closely related to each other. Indeed, transitional forms occur and in many patients

an evolution from one entity into another occurs during the course of the disease (Fig. 19.1). A single acquired mutation of the cytoplasmic tyrosine kinase Janus-associated kinase 2 (JAK2) (Val617Phe) occurs (heterozygous or homozygous) in the marrow and blood of almost all patients with PRV and in approximately 50% of those with ET and myelofibrosis, showing the common aetiology of these three diseases (Fig. 19.2). The mutation occurs in a highly conserved region of the pseudokinase domain, which is believed to negatively regulate JAK2




Fig. 19.1 Relationship between the three myeloproliferative diseases. They may all arise by somatic mutation in the pluripotent stem and progenitor cells. Many transitional cases occur showing features of two conditions and, in other cases, the disease transforms during its course from one of these diseases to another or to acute myeloid leukaemia. The three diseases, polycythaemia rubra vera, essential thrombocythaemia and myelofibrosis, are characterised by JAK2 mutation in a varying proportion of cases (see text).




Fig. 19.2 The role of JAK2 mutation in the generation of myeloproliferative diseases. (a) (i) Most haemopoietic growth factor receptors do not have intrinsic kinase activity but associate with a protein kinase such as JAK2 in the cytoplasm. (ii) When the receptor binds a growth factor the cytoplasmic domains move closer together and the JAK2 molecules can activate each other by phosphorylation. (iii) The V617F JAK2 mutation allows the JAK protein to become activated even when no growth factor is bound. (b) DNA sequencing shows homozygous G → T mutation in JAK2 in granulocytes but not in T lymphocytes (left-hand panel) and heterozygous mutation in right-hand panel. (After Kralovic R. *et al.*

(2005) *N Engl J Med* 352,1779–90) (c) JAK2 activation leads to cell survival and proliferation through activation of three major pathways; the STAT transcription factors, the PI3K pathway acting through Akt and Ras activation which subsequently activate ERK and MAPK. The net result is production of a diverse range of proteins that promote cell survival and proliferation. (d) A model for the development of myeloproliferative disease following JAK2 mutation. The primary event appears to predispose to an acquired heterozygous mutation of JAK2 (V617F). This leads to a survival advantage. In some patients, a mitotic recombination event leads to a homozygous JAK2 mutation state.

Table 19.1 Radiodilution methods for measuring red cell and plasma volume.

	Normal	Primary or secondary polycythaemia	Relative polycythaemia
Total red cell volume (^{51}Cr)	Men 25–35 mL/kg Women 22–32 mL/kg	Increased	Normal
Total plasma volume (^{125}I -albumin)	40–50 mL/kg	Normal	Decreased

signalling. JAK2 plays a major role in normal myeloid development by transducing signals from diverse cytokines and growth factors including interleukin-3 (IL-3), erythropoietin, granulocyte-macrophage colony-stimulating factor (GM-CSF), granulocyte colony-stimulating factor (G-CSF) and thrombopoietin (Fig. 1.8). Why the same mutation is associated with different myeloproliferative diseases is unclear. The exact cell in which the mutation arises, the genetic background of individual subjects and other factors may be relevant. Chronic myeloid leukaemia, which used to be classified as a myeloproliferative disorder but is now recognized as a separate entity, is discussed in Chapter 13.

Polycythaemia

Polycythaemia (erythrocytosis) is defined as an increase in the haemoglobin concentration above the upper limit of normal for the patient's age and sex.

Classification of polycythaemia

Polycythaemia is classified according to its pathophysiology but the major subdivision is into absolute polycythaemia, in which the red cell mass (volume) is raised, and relative or pseudopolycythaemia in which the red cell volume is normal but the plasma volume is reduced (Table 19.1). Absolute polycythaemia can then be subdivided into primary polycythaemia (PRV) or secondary polycythaemia (Table 19.2).

Polycythaemia (rubra) vera

In PRV, the increase in red cell volume is caused by a clonal malignancy of a marrow stem cell. The disease results from somatic mutation of a single

Table 19.2 Causes of polycythaemia.

Primary

Polycythaemia (rubra) vera
Familial (congenital) polycythaemia

Secondary

Caused by compensatory erythropoietin increase in:

high altitudes
pulmonary disease and alveolar hypoventilation (sleep apnoea)
cardiovascular disease, especially congenital with cyanosis
increased affinity haemoglobin (familial polycythaemia) (Chapter 19)
heavy cigarette smoking

Caused by inappropriate erythropoietin increase in:

renal diseases (e.g. hydronephrosis, vascular impairment, cysts, carcinoma)
tumours such as uterine leiomyoma, hypemephroma, hepatocellular carcinoma, cerebellar haemangioblastoma

Relative

Stress or pseudopolycythaemia
Cigarette smoking
Dehydration: water deprivation, vomiting
Plasma loss: burns, enteropathy

haemopoietic stem cell which gives its progeny a proliferative advantage. The JAK2 mutation is present in haemopoietic cells in almost 100% of patients. Although the increase in red cells is the diagnostic finding, in many patients there is also an overproduction of granulocytes and platelets.

Chromosome abnormalities (e.g. deletions of 9p or 20q) are found in a minority of subjects.

Diagnosis

Making the diagnosis of PRV in a patient who

Table 19.3 Criteria for diagnosis of polycythaemia (rubra) vera.

Category A

Total red cell mass

male >35 mL/kg

female >32 mL/kg

Arterial oxygen saturation >92%

Splenomegaly

JAK2 mutation

Category B

Platelets >400 × 10⁹/L

White cells >12 × 10⁹/L

Increased NAP score

Raised serum vitamin B₁₂ level

NAP, neutrophil alkaline phosphatase.

presents with polycythaemia can be difficult and the diagnostic criteria of the Polycythaemia Vera Study Group are very valuable (Table 19.3).

Clinical features

This is a disease of older subjects with an equal sex incidence. Clinical features are the result of hyperviscosity, hypervolaemia or hypermetabolism.

1 Headaches, dyspnoea, blurred vision and night sweats. Pruritus, characteristically after a hot bath, can be a severe problem.

2 Plethoric appearance: ruddy cyanosis (Fig. 19.3), conjunctival suffusion and retinal venous engorgement.

3 Splenomegaly in 75% of patients (Fig. 19.4a).

4 Haemorrhage (e.g. gastrointestinal, uterine, cerebral) or thrombosis either arterial (e.g. cardiac, cerebral, peripheral) or venous (e.g. deep or superficial leg veins, cerebral, portal or hepatic veins) are frequent.

5 Hypertension in one-third of patients.

6 Gout (as a result of raised uric acid production; Fig. 19.5a).

7 Peptic ulceration occurs in 5–10% of patients.

Laboratory findings

1 The haemoglobin, haematocrit and red cell count are increased. The total red cell volume (TRCV; Table 19.1) is increased.




Fig. 19.3 Polycythaemia vera: facial plethora and conjunctival suffusion in a 63-year-old woman.

Haemoglobin 18 g/dL; total red cell volume 45 mL/kg.

2 A neutrophil leucocytosis is seen in over half of patients, and some have increased circulating basophils.

3 A raised platelet count is present in about half of patients.

4 The *JAK2* mutation is present in the bone marrow and peripheral blood granulocytes in nearly 100% of patients.

5 The neutrophil alkaline phosphatase (NAP) score is usually increased.

6 Increased serum vitamin B₁₂ and vitamin B₁₂-binding capacity because of an increase in haptocorrin.

7 The bone marrow is hypercellular with prominent megakaryocytes, best assessed by a trephine biopsy (Fig. 19.6a).

8 Serum erythropoietin usually low.

9 Expression of PRV-1 (a surface receptor) increased; reduced expression of Mpl (thrombopoietin receptor).

10 Blood viscosity is increased.

11 Plasma urate is often increased; the serum lactate dehydrogenase (LDH) is normal.



(a)



(b)

Fig. 19.4 Splenomegaly: enlarged spleens in male patients with (a) polycythaemia vera and (b) myelofibrosis.

12 Circulating erythroid progenitors (erythroid colony-forming unit, CFU_E , and erythroid burst-forming unit, BFU_E ; p. 2) are increased compared to normal and grow *in vitro* independently of added erythropoietin (endogenous erythroid colonies).

Treatment

Treatment is aimed at maintaining a normal blood count. The haematocrit should be maintained at about 0.45 and the platelet count below $400 \times 10^9/L$.

Venesection

Venesection to reduce the haematocrit to less than 0.45 is particularly useful when a rapid reduction of red cell volume is required (e.g. at the start of therapy). It is especially indicated in younger patients and those with mild disease. The resulting iron deficiency may limit erythropoiesis. Unfortunately, venesection does not control the platelet count.

Cytotoxic myelosuppression

This is considered if there is poor tolerance of venesection, symptomatic or progressive splenomegaly, thrombocytosis, weight loss or night sweats. Daily hydroxycarbamide (hydroxyurea) is valuable in controlling the blood count and may need to be continued for many years (Fig. 19.7). Side-effects of hydroxyurea include myelosuppression, nausea



(a)



(b)

Fig. 19.5 (a) The feet of a 72-year-old man with polycythaemia rubra vera. There is inflammation of the right metatarsophalangeal and other joints caused by uric acid deposits. (b) Gangrene of the left fourth toe in essential thrombocythaemia.

Fig. 19.6 Iliac crest trephine biopsies. (a) Polycythaemia vera: fat spaces are almost completely replaced by hyperplastic haemopoietic tissue. All haemopoietic cell lines are increased with megakaryocytes particularly prominent. (b) Myelofibrosis: normal marrow architecture is lost and haemopoietic cells are surrounded by increased fibrous tissue and intercellular substance.





Fig. 19.7 Haematological response to therapy with hydroxyurea in polycythaemia vera. Hb, haemoglobin; WBC, white blood cells.

and skin toxicity. Busulfan, which can be used intermittently, is sometimes used in older patients. Pipobroman is similar to alkylating agents and is used in Europe but not in the UK. The concern with

cytotoxic drugs, especially busulfan, is that they may be associated with an increased rate of progression to leukaemia. This risk is low and there may be no increased risk with hydroxyurea.

Phosphorus-32 therapy

This is only used for older patients with severe disease. ^{32}P is a β -emitter, with a half-life of 14.3 days. It is concentrated in bone and is a most effective myelosuppressive agent. The usual remission time after a single dose is 2 years. Concern about late development of leukaemia limits its use.

Interferon

α -Interferon subcutaneously suppresses excess proliferation in the marrow and has produced good haematological responses. It is less convenient than the oral agents and side-effects are frequent. It may be particularly valuable in controlling itching. It is the first-line drug for patients less than 40 years old.

Aspirin

Low-dose aspirin reduces thrombotic complications without an increased risk of major haemorrhage.

Course and prognosis

Typically, the prognosis is good with a median survival of 10–16 years. Thrombosis and haemorrhage are the major clinical problems. Increased viscosity, vascular stasis and high platelet levels may all contribute to thrombosis, whereas defective platelet function may promote haemorrhage.

Transition from PRV to myelofibrosis occurs in approximately 30% of patients and approximately 5% of patients progress to acute leukaemia. ^{32}P and busulfan are generally avoided particularly in younger subjects as they may increase this risk.

Primary familial (congenital) polycythaemia

This is a rare condition. In one type there is a mutation in the von Hippel–Lindau protein gene which creates abnormal oxygen sensing with increased erythropoietin production.

Secondary polycythaemia

The causes of secondary polycythaemia are listed in Table 19.2. Hypoxia caused by chronic obstructive airways disease is one of the most frequent, and measurement of arterial oxygen saturation is a valuable

test. Renal and tumour causes of inappropriate erythropoietin secretion are rare. Patients with a high-affinity haemoglobin often have a family history of polycythaemia and present at a young age. Those with cyanotic congenital heart disease should be managed in conjunction with a congenital heart disease unit. Treatment of obesity, hypertension and discontinuation of smoking may lower the haematocrit. These patients are treated by venesection for symptoms such as dizziness, dyspnoea or angina if a raised haematocrit is a contributory factor; also if there has been a previous thrombotic episode. Patients with polycythaemia secondary to hypoxia and pulmonary disease who have symptoms of hyperviscosity or haematocrit of more than 0.56 should have their haematocrit reduced by venesection.

Apparent polycythaemia

Apparent polycythaemia, also known as pseudopolycythaemia, is the result of plasma volume contraction. By definition, the TRCV is normal. The cause is uncertain but it is far more common than PRV. It occurs particularly in young or middle-aged men and may be associated with cardiovascular problems e.g. hypertension (Gaisböck's syndrome) myocardial ischaemia or cerebral transient ischaemic attacks. Diuretic therapy, heavy smoking, obesity and alcohol consumption are frequent associations. Venesection to maintain a haematocrit around 0.45–0.47 is recommended in those with a recent history of thrombosis or with additional risk factors and in all with a haematocrit of more than 0.52.

Differential diagnosis of polycythaemia

A rational approach is needed to evaluate a patient presenting with a high haemoglobin. If PRV is suspected, polymerase chain reaction (PCR) test for the *JAK2* mutation will confirm this in almost all cases. The full blood count, the NAP score, bone marrow aspirate trephine biopsy and ultrasound of the abdomen to assess spleen size and to detect renal abnormalities are the most useful additional tests. Studies with ^{51}Cr -labelled red cells to measure TRCV may be needed if *JAK2* is negative. If absolute

polycythaemia is present look for lung or cardiac disease, check arterial PO_2 and consider a check of the oxygen dissociation curve and haemoglobin (Hb) electrophoresis. Finally, look for erythropoietin secreting tumours by ultrasound, computed tomography (CT) or magnetic resonance imaging (MRI). The serum erythropoietin level is also useful in screening for tumours.

Essential thrombocythaemia

In this condition there is a sustained increase in platelet count, because of megakaryocyte proliferation and overproduction of platelets. The haematocrit is normal and the Philadelphia chromosome or *BCR-ABL* rearrangement are absent. The bone marrow shows no collagen fibrosis. A persisting platelet count of $>400 \times 10^9/L$ is the central diagnostic feature but other causes of a raised platelet count (particularly iron-deficiency, inflammatory or malignant disorder and myelodysplasia) need to be fully excluded before the diagnosis can be made. Half of patients show the *JAK2* (Val617Phe) mutation and these cases tend to resemble more closely polycythaemia vera with higher haemoglobin and white cell counts and splenomegaly than *JAK2* negative cases. There is endogenous megakaryocyte and possibly erythroid colony growth independent of thrombopoietin or erythropoietin. Rare primary familial cases in children have been associated with mutations on the genes for thrombopoietin or its receptor *c-Mpl*.

Clinical and laboratory findings

The dominant clinical features are thrombosis and haemorrhage. Many cases are symptomless and diagnosed on routine blood counts. Thrombosis may occur in the venous or arterial systems (Fig. 19.5b) whereas haemorrhage, as a result of abnormal platelet function, may cause either chronic or acute bleeding. A characteristic symptom is erythromelalgia, a burning sensation felt in the hands or feet and promptly relieved by aspirin. Up to 40% of patients will have palpable splenomegaly whereas in others there may be splenic atrophy because of infarction. Abnormal large platelets and megakaryocyte fragments may be seen on the blood film (Fig. 19.8). The




Fig. 19.8 Peripheral blood film in essential thrombocythaemia showing increased numbers of platelets and a nucleated megakaryocytic fragment.

Table 19.4 Causes of a raised platelet count.

<i>Reactive</i>
Haemorrhage, trauma, postoperative
Chronic iron deficiency
Malignancy
Chronic infections
Connective tissue diseases (e.g. rheumatoid arthritis)
Post-splenectomy
<i>Endogenous</i>
Essential thrombocythaemia (<i>JAK2</i> mutation + or –)
In some cases of polycythaemia vera, myelofibrosis and chronic myeloid leukaemia

bone marrow is similar to that in PRV but an excess of abnormal megakaryocytes is typical. Cytogenetics and molecular analysis for the *BCR-ABL* fusion gene are analysed to exclude chronic myeloid leukaemia. The condition must be distinguished from other causes of a raised platelet count (Table 19.4). Platelet function tests (p. 277) are consistently abnormal, failure of aggregation with adrenaline being particularly characteristic.

Treatment

The principle is to control the platelet count so as to reduce the risk of thrombosis which is the major clinical problem. The patients most at risk are those over 60 years old, with a platelet count

$>1000 \times 10^9/L$ and with previous episodes of thrombosis or haemorrhage. The thrombotic risk depends on other risk factors such as smoking history and hypertension, and the treatment should take account of these risks. In those with a high risk the aim is to keep the platelet count below $600 \times 10^9/L$. Hydroxyurea is the most widely used treatment. α -Interferon is also valuable in younger patients but has more side-effects. Anagrelide is effective in reducing the platelet count but also has more side-effects, particularly on the cardiovascular system, than hydroxyurea. A possible increased risk of myelofibrosis is also of concern. Busulfan and ^{32}P were used but are not now favoured because of possible long-term complications. Platelet pheresis may be helpful in short-term management. Aspirin is commonly used to reduce thrombotic risk and in patients younger than 60 years with no previous thrombosis or haemorrhage, and platelets $<1000 \times 10^9/L$, it may be the treatment of choice.

Course

Often the disease is stationary for 10–20 years or more. The disease may transform after a number of years to myelofibrosis; the risk of transformation to acute leukaemia is relatively low ($<5\%$). Patients have been stratified as low risk if less than 40 years old, with no previous thrombosis, hypertension or diabetes and platelets $<1000\text{--}1500 \times 10^9/L$ and are treated with aspirin alone; intermediate risk if less than 60 years with no prior thrombosis, hypertension or diabetes and platelets $<1000\text{--}1500 \times 10^9/L$; or high risk if over 60 years or with prior thrombosis or haemorrhage, hypertension or diabetes, platelet count $>1000 \times 10^9/L$. High-risk patients receive hydroxyurea and aspirin. For intermediate risk, trials of aspirin versus hydroxyurea and aspirin are in progress.

Myelofibrosis

Confusingly, the condition has many names: idiopathic myelofibrosis; myelosclerosis; agnogenic myeloid metaplasia; or myelofibrosis with myeloid metaplasia (MMM).

The predominant feature of myelofibrosis is a progressive generalized reactive fibrosis of the bone

marrow in association with the development of haemopoiesis in the spleen and liver (known as myeloid metaplasia). Clinically this leads to anaemia and massive hepatosplenomegaly. In some patients there is osteosclerosis. Myelofibrosis is a clonal stem cell disease. The fibrosis of the bone marrow is secondary to hyperplasia of abnormal megakaryocytes. It is thought that fibroblasts are stimulated by platelet-derived growth factor and other cytokines secreted by megakaryocytes and platelets. The *JAK2* mutation occurs in approximately 50% of patients. The absence of the mutation in many patients shows genetic heterogeneity in myelofibrosis as in ET. Non-specific cytogenetic abnormalities may be found in approximately half the patients.

One-third or more of the patients have a previous history of PRV and some patients present with clinical and laboratory features of both disorders.

Clinical features

- 1 An insidious onset in older people is usual with symptoms of anaemia.
- 2 Symptoms resulting from massive splenomegaly (e.g. abdominal discomfort, pain or indigestion) are frequent; splenomegaly is the main physical finding (Fig. 19.4b).
- 3 Hypermetabolic symptoms such as loss of weight, anorexia, fever and night sweats are common.
- 4 Bleeding problems, bone pain or gout occur in a minority of patients.

Myelofibrosis and chronic myeloid leukaemia are responsible for most cases of massive (>20 cm) splenic enlargement in the UK and North America (see Table 9.1).

Laboratory findings

- 1 Anaemia is usual but a normal or increased haemoglobin level may be found in some patients.
- 2 The white cell and platelet counts are frequently high at the time of presentation. Later in the disease leucopenia and thrombocytopenia are common.
- 3 A leucoerythroblastic blood film is found. The red cells show characteristic 'tear-drop' poikilocytes (Fig. 19.9).
- 4 Bone marrow is usually unobtainable by aspiration. Trephine biopsy (Fig. 19.6b) shows a fibrotic




Fig. 19.9 Peripheral blood film in myelofibrosis. Leucoerythroblastic change with 'tear-drop' cells and an erythroblast.

hypercellular marrow. Increased megakaryocytes are frequently seen. In 10% of cases there is increased bone formation with increased bone density on X-ray.

5 JAK2 kinase is mutated in approximately 50% of cases.

6 NAP score is usually increased.

7 High serum urate and LDH levels reflect the increased but largely ineffective turnover of haemopoietic cells.

8 Transformation to acute myeloid leukaemia occurs in 10–20% of patients.

Treatment

This is usually palliative and aimed at reducing the effects of anaemia and splenomegaly. Blood transfusions and regular folic acid therapy are used in severely anaemic patients. Hydroxyurea may help to reduce splenomegaly and hypermetabolic symptoms. Trials of thalidomide, cyclophosphamide or etanercept (anti-TNF α) are in progress. Danazol, an androgen derivative may improve anaemia in approximately 30% of patients. Erythropoietin may also be tried. Splenectomy is considered for patients with severe symptomatic splenomegaly—mechanical discomfort, thrombocytopenia, portal hypertension, excessive transfusion requirements or hypermetabolic symptoms. Splenic irradiation is an alternative but usually provides relief only for

3–6 months. Allopurinol is indicated in virtually all patients to prevent gout and urate nephropathy from hyperuricaemia. Allogeneic stem cell transplantation may be curative for young patients.

The median survival is approximately 3.5 years and causes of death include heart failure, infection and leukaemic transformation. A haemoglobin level of less than 10 g/dL, a white cell count of less than 4 or greater than $30 \times 10^9/L$ and the presence of abnormal chromosomes are associated with a worse prognosis.

Systemic mastocytosis

Mast cells are derived from haemopoietic stem cells. Mature cells survive for months or years in vascular tissues and most organs. Systemic mastocytosis is a clonal myeloproliferative disorder involving usually the bone marrow, heart, spleen, lymph nodes and skin.

The somatic *c-kit* mutation Asp816Val is detected in the majority of patients and may be partly responsible for autonomous growth and enhanced survival of the neoplastic mast cells. In many patients this mutation is also detected in other haemopoietic cells.

Symptoms are related to histamine and prostaglandin release and include flushing, pruritus, abdominal pain and bronchospasm. The skin usually shows urticaria pigmentosa.

In many patients the course is indolent. In other cases, the disease is associated with acute leukaemia or other dysplastic conditions.

Bibliography

- Baxter E.J., Scott L.M., *et al.* (2005) A mutation of the tyrosine kinase JAK2 in myeloproliferative disorders. *Lancet* 366,122.
- Belláné-Chantelot C., *et al.* (2006) Genetic and clinical features of JAK2 mutation in myeloproliferative diseases. *Blood* 108,341–346.
- Cross N.C. and Reiter J.E. (2005) JAK2 mutations in chronic myeloproliferative disorders. *Blood* 106,1207–12.
- Finazzi G., Caruso V., *et al.* (2005) JAK2 mutation in polycythemia vera: patients enrolled in a large international study. *Blood* 105,2664–70.
- Harrison C.N., Campbell P.H., *et al.* (2005) Hydroxyurea compared with no therapy: a risk essential thrombocythemia. *Blood* 106,45–50.
- Harrison C.N., Gale R.J., *et al.* (2005) A large proportion of patients with essential thrombocythemia do not have a JAK2 mutation and may be at lower risk of thrombotic events. *Blood* 106,93,417–24.

CHAPTER 20

20

Aplastic anaemia and bone marrow failure

Pancytopenia, 241
Aplastic anaemia, 241
Red cell aplasia, 246

Schwachman–Diamond syndrome, 246
Congenital dyserythropoietic anaemia, 246
Bibliography, 247

Pancytopenia

Pancytopenia describes a reduction in the blood count of all the major cell lines—red cells, white cells and platelets. It has several causes (Table 20.1) which can be broadly divided into decreased bone marrow production or increased peripheral destruction.

Aplastic anaemia

Aplastic (hypoplastic) anaemia is defined as pancytopenia resulting from aplasia of the bone marrow. It is classified into primary (congenital or acquired) or secondary types (Table 20.2).

Pathogenesis

The underlying defect in all cases appears to be a

Table 20.1 Causes of pancytopenia.

Decreased bone marrow function

Aplasia

Acute leukaemia, myelodysplasia, myeloma

Infiltration with lymphoma, solid tumours, tuberculosis

Megaloblastic anaemia

Paroxysmal nocturnal haemoglobinuria

Myelofibrosis

Haemophagocytic syndrome

Increased peripheral destruction

Splenomegaly

substantial reduction in the number of haemopoietic pluripotential stem cells, and a fault in the remaining stem cells or an immune reaction against them, which makes them unable to divide and differentiate sufficiently to populate the bone marrow (Fig. 20.1). A primary fault in the marrow micro-environment has also been suggested but the success of stem cell transplantation (SCT) shows this can only be a rare cause because normal donor stem cells are usually able to thrive in the recipient's marrow cavity.

Congenital

The Fanconi type has an autosomal recessive pattern of inheritance and is often associated with growth retardation and congenital defects of the skeleton (e.g. microcephaly, absent radii or thumbs), of the renal tract (e.g. pelvic or horseshoe kidney) (Fig. 20.2) or skin (areas of hyper- and hypopigmentation); sometimes there is mental retardation. The syndrome is genetically heterogeneous with eight different complementation groups termed A, B, C, D1, D2, E, F and G. The *FAND1* gene is identical to *BRCA2*, the breast cancer susceptibility gene. The encoded proteins cooperate in a common cellular pathway involved in DNA repair. Cells from Fanconi's anaemia (FA) patients show an abnormally high frequency of spontaneous chromosomal breakage and the diagnostic test is elevated breakage after incubation of peripheral blood lymphocytes

Table 20.2 Causes of aplastic anaemia.

Primary	Secondary
Congenital (Fanconi and non-Fanconi types)	Ionizing radiation: accidental exposure (radiotherapy, radioactive isotopes, nuclear power stations)
Idiopathic acquired	Chemicals: benzene, organophosphates and other organic solvents, DDT and other pesticides, organochlorines, recreational drugs (ecstasy)
	Drugs
	Those that regularly cause marrow depression (e.g. busulfan, cyclophosphamide, anthracyclines, nitrosoureas)
	Those that occasionally or rarely cause marrow depression (e.g. chloramphenicol, sulphonamides, gold, anti-inflammatory, antithyroid, psychotropic, anticonvulsant/antidepressant drugs)
	Viruses: viral hepatitis (non-A, non-B, non-C, non-G in most cases), EBV

EBV, Epstein-Barr virus.





Fig. 20.1 Aplastic anaemia: low power views of bone marrow show severe reduction of haemopoietic cells with an increase in fat spaces. (a) Aspirated fragment. (b) Trephine biopsy.

with the DNA cross-linking agent diepoxybutane (DEB test). Dyskeratosis congenita is a rare sex-linked disorder with nail and skin atrophy and is associated with mutations in the *DKC1* (dyskerin)

or *TERC* (telomerase reverse transcriptase RNA template) genes which are both involved in the maintenance of telomere length.

The usual age of presentation of FA is 5–10 years.



(a)



(b)

Fig. 20.2 (a) X-rays showing absent thumbs in a patient with Fanconi's anaemia (FA). (b) Intravenous pyelogram in a patient with FA showing a normal right kidney but a left kidney abnormally placed in the pelvis.

Approximately 10% of cases develop acute myeloid leukaemia. Treatment is usually with androgens and/or SCT. The blood count usually improves with androgens but side-effects, especially in children, are distressing (virilization and liver abnormalities); remission rarely lasts more than 2 years. SCT may cure the patient. Because of the sensitivity of the patient's cells to DNA damage, conditioning regimes are mild.

Idiopathic acquired

This is the most common type of aplastic anaemia, accounting for at least two-thirds of acquired cases. In most cases haemopoietic tissue is the target of an immune process dominated by oligoclonal expression of cytotoxic T cells which secrete γ -interferon and tumour necrosis factor. In approximately one-third of cases short telomeres are found in leucocytes, especially in those with a prolonged clinical course. Mutations in the telomere repair complex have been described but their relevance is unclear. The favourable responses to antilymphocyte globulin (ALG) and ciclosporin support the concept that autoimmune T-cell mediated damage, possibly against functionally and structurally altered stem cells, is important.

Secondary

This is often caused by direct damage to the haemopoietic marrow by radiation or cytotoxic drugs. The antimetabolite drugs (e.g. methotrexate) and mitotic inhibitors (e.g. daunorubicin) cause only temporary aplasia but the alkylating agents, particularly busulfan, may cause chronic aplasia closely resembling the chronic idiopathic disease. Some individuals develop aplastic anaemia as a rare idiosyncratic side-effect of drugs such as chloramphenicol or gold which are not known to be cytotoxic (Table 20.2). They may also develop the disease during or within a few months of viral hepatitis (rarely hepatitis A, B or C, but more frequently non-A, non-B, non-C). Because the incidence of marrow toxicity is particularly high for chloramphenicol, this drug should be reserved for treatment of those infections that are life-threatening and for which it is the optimum antibiotic (e.g. typhoid).

Chemicals such as benzene may be implicated and rarely aplastic anaemia may be the presenting feature of acute lymphoblastic or myeloid leukaemia, especially in childhood. Myelodysplasia (Chapter 14) may also present with a hypoplastic marrow.

Clinical features

The onset is at any age with a peak incidence around 30 years and a slight male predominance; it can be insidious or acute with symptoms and signs resulting from anaemia, neutropenia or thrombocytopenia. Infections, particularly of the mouth and throat, are common and generalized infections are frequently life-threatening; bruising, bleeding gums, epistaxes and menorrhagia are the most frequent haemorrhagic manifestations and the usual presenting features, often with symptoms of anaemia. The lymph nodes, liver and spleen are not enlarged.

Laboratory findings

- 1 Anaemia is normochromic, normocytic or macrocytic (mean cell volume (MCV) often 95–110 fL). The reticulocyte count is usually extremely low in relation to the degree of anaemia.
- 2 Leucopenia. There is a selective fall in granulocytes, usually but not always to below $1.5 \times 10^9/L$. In severe cases, the lymphocyte count is also low. The neutrophils appear normal.
- 3 Thrombocytopenia is always present and, in severe cases, is less than $20 \times 10^9/L$.
- 4 There are no abnormal cells in the peripheral blood.
- 5 The bone marrow shows hypoplasia, with loss of haemopoietic tissue and replacement by fat which comprises over 75% of the marrow. Trepine biopsy is essential and may show patchy cellular areas in a hypocellular background (Fig. 20.1b). The main cells present are lymphocytes and plasma cells; megakaryocytes in particular are severely reduced or absent.

Severe cases show neutrophils $<0.5 \times 10^9/L$ (very severe $<0.2 \times 10^9/L$), platelets $<20 \times 10^9/L$, reticulocytes $<20 \times 10^9/L$ and marrow cellularity $<25\%$.

Diagnosis

The disease must be distinguished from other causes of pancytopenia (Table 20.1) and this is not

usually difficult provided an adequate bone marrow sample is obtained. Cytogenetic analysis should be performed. Paroxysmal nocturnal haemoglobinuria (PNH) must be excluded by flow-cytometry testing of red cells for CD55 and CD59. In older patients, hypoplastic myelodysplasia may show similar appearances. Qualitative abnormalities of the cells and clonal cytogenetic changes suggest myelodysplasia rather than aplastic anaemia. Some patients diagnosed as having aplastic anaemia develop PNH, myelodysplasia or acute myeloid leukaemia in subsequent years. This may occur even in patients who have responded well to immunosuppressive therapy.

Treatment

General

The cause, if known, is removed (e.g. radiation or drug therapy is discontinued). Initial management consists largely of supportive care with blood transfusions, platelet concentrates, and treatment and prevention of infection. All blood products should be filtered to reduce the risk of alloimmunization and irradiated to prevent grafting of live donor lymphocytes. In severely thrombocytopenic (platelet count $<10 \times 10^9/L$) and neutropenic (neutrophils $<0.5 \times 10^9/L$) patients, management is similar to the supportive care of patients receiving intensive chemotherapy with reverse barrier isolation. An antifibrinolytic agent (e.g. tranexamic acid) may be used in patients with severe prolonged thrombocytopenia. Oral antifungal agents and oral antibiotics are used prophylactically in some units to reduce the incidence of infection.

Specific

This must be tailored to the severity of the illness as well as the age of the patient and potential sibling stem cell donors. Severity is assessed by the reticulocyte, neutrophil and platelet counts and degree of marrow hypoplasia. Severe cases have a high mortality in the first 6–12 months unless they respond to specific therapy. Less severe cases may have an acute transient course or a chronic course with ultimate recovery, although the platelet count often remains subnormal for many years. Relapses, sometimes severe and occasionally fatal,

may also occur and rarely the disease transforms into myelodysplasia, acute leukaemia or PNH (Chapter 5).

The following 'specific' treatments are used with varying success.

1 Antilymphocyte or antithymocyte globulin (ALG or ATG) This is prepared in animals (e.g. horse or rabbit) and is of benefit in approximately 50–60% of acquired cases. It is usually given with corticosteroids which also reduce the side-effects of ALG including the serum sickness of fever, rash and joint pains which may occur approximately 7 days after administration. Corticosteroids should not be used alone as they increase the risk of infection. Typically, if there is no response to ALG after 4 months a second course may be tried, prepared from another species. Overall, up to 80% of patients respond to combined ALG, steroids and ciclosporin.

2 Ciclosporin This is an effective agent which appears particularly valuable in combination with ALG and steroids.

3 Androgens These are beneficial in some patients with FA and acquired aplastic anaemia although an overall improved survival in acquired aplastic anaemia has not been proven. Oxymetholone is usually tried but side-effects are marked including virilization, salt retention and liver damage with cholestatic jaundice or rarely hepatocellular carcinoma. If there is no response in 4–6 months, androgens should be stopped. If there is a response the drug should be withdrawn gradually.

4 Stem cell transplantation Allogeneic transplantation offers the chance of permanent cure. For aplastic anaemia conditioning is with cyclophosphamide without irradiation but with ciclosporin which reduces the risks of graft failure and (with methotrexate) of graft-versus-host disease. The relative role of SCT versus immunosuppressive therapy in individual patients with aplastic anaemia is under constant review. In general terms, SCT is favoured in younger patients with severe aplastic anaemia and a human leucocyte antigen (HLA) matching sibling donor. Cure rates of up to 80% are obtained. However, non-myeloablative transplants (p. 262) are used in selected patients over the age of 40 years. SCT using unrelated volunteer donors and mismatched family members is achieving a survival ratio over 50% in patients with severe aplastic

anaemia. In older subjects and those with less severe disease, immunosuppression is usually tried first.

5 Haemopoietic growth factors Granulocyte colony-stimulating factor (G-CSF) may produce minor responses but does not lead to sustained improvement. Other growth factors have not proved helpful.

Red cell aplasia

Chronic form

This is a rare syndrome characterized by anaemia with normal leucocytes and platelets and grossly reduced or absent erythroblasts in the marrow (Fig. 20.3). The congenital form is known as Diamond–Blackfan syndrome (Table 20.3) and is inherited as a recessive condition. It is associated with a varying number of somatic disorders (e.g. of the face or heart). Mutation of a gene on chromosome 19 that encodes a ribosomal protein underlies some cases. Corticosteroids are the first line of treatment and SCT may be curative. Androgens may also produce improvement but side-effects on growth can be severe.

The acquired chronic form can occur without any obvious associated disease or precipitating factor (idiopathic), or may be seen with autoimmune diseases (especially systemic lupus erythematosus), with a thymoma, lymphoma or chronic lymphocytic leukaemia. Red cell aplasia from anti-erythropoietin antibodies has been rarely described in patients with chronic renal failure receiving recombinant erythropoietin. In some cases, immunosuppression

with corticosteroids, ciclosporin, azathioprine or ALG is helpful. Monoclonal antibodies (e.g. Campath (anti-CD52) or rituximab (anti-CD20)) are being increasingly used in treatment of refractory acquired red cell aplasia and other autoimmune cytopenias.

If regular blood transfusions are needed, iron chelation therapy will also be necessary. SCT has been carried out in some severe cases.

Transient form

Parvovirus B19 infects red cell precursors via the P antigen and causes a transient (5–10 days) red cell aplasia with the rapid onset of severe anaemia in patients with pre-existing shortened red cell survival (e.g. sickle cell disease or hereditary spherocytosis; Fig. 20.4). Transient red cell aplasia with anaemia may also occur in association with drug therapy (Table 20.3), and in normal infants or children, often with a history of a viral infection in the preceding 3 months.

Schwachman–Diamond syndrome (rare autosomal recessive syndrome)

This is a rare autosomal recessive syndrome characterized by varying degrees of cytopenia, especially neutropenia with a propensity to transform to myelodysplasia or acute myeloid leukaemia. Exocrine pancreatic dysfunction is an invariable feature while skeletal abnormalities, hepatic impairment and short stature are frequent.

Congenital dyserythropoietic anaemia

Congenital dyserythropoietic anaemias (CDAs) are a group of hereditary refractory anaemias characterized by ineffective erythropoiesis and erythroblast multinuclearity. The patient may be jaundiced with bone marrow expansion. The white cell and platelet counts are normal. The reticulocyte count is low for the degree of anaemia, despite increased marrow cellularity. The anaemia is of variable severity and is usually first noted in infancy or childhood. Iron overload may develop and splenomegaly is common. The CDAs have been classified into four types based on the degree to which megaloblastic changes, giant erythroblasts and dyserythropoietic changes




Fig. 20.3 The bone marrow in primary red cell aplasia. There is selective loss of erythropoiesis.




Fig. 20.4 Parvovirus infection: flow chart showing transient fall in haemoglobin and reticulocytes in a patient with hereditary spherocytosis.

Table 20.3 Classification of pure red cell aplasia.

Acute, transient	Chronic Congenital	Acquired
Parvovirus infection Infancy and childhood Drugs (e.g. azathioprine, co-trimoxazole)	Diamond–Blackfan syndrome	Idiopathic Associated with thymoma, systemic lupus erythematosus, rheumatoid arthritis, lymphoma, chronic lymphocytic leukaemia, T-large granular lymphocytosis, myelodysplasia, viral infection, drugs

are present. Type II is known as HEMPAS (hereditary erythroblast multinuclearity with a positive acidified serum lysis test) with some sera but not the patient's serum. The basic lesion in HEMPAS is a genetic defect in one of two enzymes, N-acetylglucosaminyltransferase or α -mannosidase, which are involved in glycosylation of several red cell membrane proteins. α -Interferon has induced remission in some cases.

Bibliography

- Dokal I. (2003) Inherited aplastic anaemia. *Hematol J* 4,3–9.
- Dror Y. and Freedman M.H. (2002) Shwachman–Diamond syndrome. *Br J Haematol* 118,701–13.
- Frickhofen N., Heimpel H., Kaltwasser J.P., Schrezenmeier H. for the German Aplastic Anemia Study Group. (2003) Antithymocyte globulin with or without cyclosporin A: 11-year follow-up of a randomized trial comparing treatments of aplastic anemia. *Blood* 101,1236–42.
- Issaragrisil S., Kaufman D.W., Anderson T. *et al.* (2006) The epidemiology of aplastic anemia in Thailand. *Blood* 107,1299–307.
- Joenje H. and Patel K.J. (2001) The emerging genetic and molecular basis of Fanconi anaemia. *Nat Rev Genet* 2,446–57.
- Locasciulli A., Bruno B., Rambaldi A. *et al.* GITMO Prospective Randomized Study Group (2004) Treatment of severe aplastic anemia with antilymphocyte globulin, cyclosporine and two different granulocyte colony-stimulating factor regimens: a GITMO prospective randomized study. *Haematologica* 89,1054–61.
- Marsh J.C. (2005) Management of acquired aplastic anaemia. *Blood Rev* 19,143–51.
- Marsh J.C., Ball S.E., Gordon-Smith E.C. *et al.* for the British

- Committee for Standards in Haematology (2003) Guidelines for the diagnosis and management of acquired aplastic anaemia. *Br J Haematol* 123,782–801.
- Muir K.R., Chilvers C.E., Harriss C. *et al.* (2003) The role of occupational and environmental exposures in the aetiology of acquired severe aplastic anaemia: a case-control investigation. *Br J Haematol* 123,906–14.
- Robak T. (2004) Monoclonal antibodies in the treatment of autoimmune cytopenias. *Eur J Haematol* 72,79–88.
- Taniguchi T. and D'Andrea A.D. (2006) Molecular pathogenesis of Fanconi anemia: recent progress. *Blood* 107,4223–33.
- Vulliamy T., Marrone A., Dokal I., Mason P.J. (2002) Association between aplastic anaemia and mutations in telomerase RNA. *Lancet* 359,2168–70.
- Yamaguchi H., Calado R.T., Ly H. *et al.* (2005) Mutations in TERT, the gene for telomerase reverse transcriptase, in aplastic anemia. *N Engl J Med* 352,1413–24.
- Young, N.S. and Brown K.E. (2004) Parvovirus B19. *N Engl J Med* 350,586–97.

Principles of stem cell transplantation, 249

Autologous stem cell transplantation, 253

Allogeneic stem cell transplantation, 254

Bibliography, 262

Principles of stem cell transplantation

Stem cell transplantation (SCT) involves eliminating a patient's haemopoietic and immune system by chemotherapy and/or radiotherapy and replacing it with stem cells either from another individual or with a previously harvested portion of the patient's own haemopoietic stem cells (Fig. 21.1). The term encompasses both *bone marrow transplantation* (BMT), in which stem cells are collected from bone marrow, and *peripheral blood stem cell* (PBSC) *transplantation* in which stem cells are collected from peripheral blood.

SCT may be *syngeneic* (from an identical twin), *allogeneic* (from another person) or *autologous* (from the patient's own stem cells) (Table 21.1).

The principal diseases for which SCT is performed are listed in Table 21.2. However, the exact role of SCT in the management of each disease is complex and depends on factors such as disease severity and subtype, remission status, age and, for allogeneic transplantation, availability of donors.

Collection of stem cells

Stem cells can be collected from bone marrow or from peripheral blood.

Bone marrow collection

The donor is given a general anaesthetic and 500–

1200 mL of marrow is harvested from the pelvis. The marrow is heparinized and a mononuclear cell count is taken to assess the yield, which should be approximately $2\text{--}4 \times 10^8$ nucleated cells/kg body weight of the recipient.

Peripheral blood stem cell collection

PBSCs are taken using a cell-separator machine connected to the patient or donor via peripheral cannulae (Fig. 21.2). Blood is taken through one cannula and pumped around the machine where mononuclear cells are collected by centrifugation before the red cells are returned to the patient. This continuous process may take a few hours before enough mononuclear cells are collected.

Peripheral blood normally contains too few haemopoietic stem cells to allow collection of sufficient numbers for transplantation. Chemotherapy and growth factors can each increase the number by around 100 times. In future it is possible that stem cell populations may also be expanded *in vitro*.

Chemotherapy is used in patients undergoing autologous stem cell collection but not in healthy donors. PBSCs are usually collected during the recovery phase from a cycle of chemotherapy such as 1.5 g/m^2 of cyclophosphamide.

Granulocyte colony-stimulating factor (G-CSF) is given to patients or donors as a course of injections (typically $10 \mu\text{g/kg/day}$ for 4–6 days) until the white cell count starts to rise. PBSC collections are taken and, depending on the efficiency of stem cell





Fig. 21.1 Procedures for (a) allogeneic, and (b) autologous stem cell transplantation. G-CSF, granulocyte colony-stimulating factor; GVHD, graft-versus-host disease.

Table 21.1 Stem cell transplantation: potential donors.

HLA-matching sibling	}	Allogeneic
Unrelated HLA-matching volunteer		
Umbilical cord blood		
Identical twin	}	Syngeneic
Self		Autologous

HLA, human leucocyte antigen.

mobilization, repeated collections may be needed for up to 3 days. The adequacy of the collection may be assessed by:

1 CD34⁺ cell count. Generally $>2.5 \times 10^6/\text{kg}$ are needed for autologous transplantation.

2 *In vitro* colony assays, particularly granulocyte-macrophage colony-forming unit (CFU-GM) (p. 2), of which $1-5 \times 10^5/\text{kg}$ would be considered adequate for transplantation.

Table 21.2 Stem cell transplantation: indications.

Allogeneic (or syngeneic)	Autologous
Acute lymphoblastic or myeloid leukaemia	Hodgkin's lymphoma and non-Hodgkin's lymphoma
Chronic myeloid leukaemia	Multiple myeloma
Other malignant disorders of the marrow (e.g. myelodysplasia, multiple myeloma, lymphoma, chronic lymphocytic leukaemia)	Acute and chronic leukaemias
Severe aplastic anaemia including Fanconi's anaemia	Severe autoimmune disease
Inherited disorders: thalassaemia major, sickle cell anaemia, immune deficiencies, inborn errors of metabolism in the haemopoietic and mesenchymal system (e.g. osteopetrosis)	Amyloidosis
Other acquired severe marrow diseases (e.g. paroxysmal nocturnal haemoglobinuria, red cell aplasia, myelofibrosis)	For 'gene therapy' of genetic disease (e.g. adenosine deaminase deficiency)

Fig. 21.2 Peripheral blood stem cell (PBSC) collection: a donor undergoing collection of PBSCs on a cell separator.






Fig. 21.3 Peripheral blood stem cell collection: enriched CD34⁺ cells stained by May-Grünwald-Giemsa stain. The cells have the appearance of small and medium-sized lymphocytes.

Stem cell processing

After collection the stem cell harvest is processed with removal of red cells and concentration of the mononuclear cells. Autologous collections may be 'purged' by chemotherapy or antibodies in an attempt to remove residual malignant cells. Allogeneic collections may be treated with antibodies to remove T cells to reduce graft-versus-host disease (GVHD). CD34⁺ stem cells may be selected from both types of harvest (Fig. 21.3).

Conditioning

Prior to infusion of haemopoietic stem cells patients receive chemotherapy, sometimes in combination with total body irradiation (TBI; Fig. 21.1) in a procedure called *conditioning*. This is designed to eradicate the patient's haemopoietic and immune system and, if present, malignancy. In addition, in the setting of allogeneic SCT it helps to suppress the host immune system and thereby prevent rejection of the 'foreign' stem cells. An important development that has occurred in SCT is a major shift from *myeloablative* conditioning regimes to *non-myeloablative* conditioning. Non-myeloablative conditioning does not completely ablate the patient's bone marrow and is discussed more fully later (p. 262).

TBI is usually used in patients with malignant disease and is administered as a single dose or in smaller doses over several days (fractionated). The most commonly used drug is cyclophosphamide but busulfan, melphalan, cytosine arabinoside, etoposide or nitrosoureas are given in some protocols. At least 36 h are allowed for the elimination of the drugs from the circulation following the last dose of chemotherapy before donor stem cells are infused. Conditioning therapy is often complicated by mucositis and patients sometimes need parenteral nutrition. Trials are taking place in which monoclonal antibodies directed against specific antigens such as CD45 are attached to toxins or radioactive isotopes in an attempt to selectively target white cells as an aid to conditioning.

Post-transplant engraftment and immunity

After a period of typically 1–3 weeks of severe pancytopenia, the first signs of successful engraftment are monocytes and neutrophils in the blood with a subsequent increase in platelet count (Fig. 21.4). A reticulocytosis also begins and natural killer (NK) cells are among the earliest donor-derived lymphocytes to appear. G-CSF may be used to reduce the period of neutropenia. Engraftment is usually quicker following PBSC transplantation compared with BMT.




Fig. 21.4 Typical haematological chart of a patient undergoing allogeneic marrow transplantation for aplastic anaemia. WBC, white blood cells.

The marrow cellularity gradually returns to normal but the marrow reserve remains impaired for 1–2 years. There is profound immunodeficiency for 3–12 months with a low level of CD4 helper cells and a raised CD8 : CD4 ratio for 6 months or more. Immune recovery is quicker after autologous and syngeneic SCT than following allogeneic SCT. The patient's blood group changes to that of the donor and antigen-specific immunity becomes that of the donor after approximately 60 days.

Autologous stem cell transplantation

This allows the delivery of a high dose of chemo-

therapy, with or without radiotherapy, which otherwise would result in prolonged bone marrow aplasia. Stem cells are harvested and stored before the treatment is given and are then reinfused to 'rescue' the patient from the myeloablative effects of the treatment (Fig. 21.1). A limitation of the procedure is that tumour cells contaminating the stem cell harvest may be reintroduced into the patient. Nevertheless, autografting has a major role in the treatment of haematological diseases such as lymphoma and myeloma and is under investigation in other diseases such as acute leukaemia and severe autoimmune diseases. The major problem associated with autografting is recurrence of the original disease.

GVHD is not an issue. Procedure-related mortality is generally well below 5%. Trials of autologous bone marrow cell transfer to treat myocardial infarct and other non-haemopoietic diseases are in progress.

Allogeneic stem cell transplantation

In this procedure, stem cells harvested from another person are infused into the patient. The procedure has a significant morbidity and mortality and one of the major reasons is the immunological incompatibility between donor and patient despite matching of the human leucocyte antigens (HLA). This may manifest as immunodeficiency, GVHD or graft failure. Paradoxically, there is also a graft-versus-leukaemia (GVL) effect which probably underlies much of the success of the procedure.

The human leucocyte antigen system

Allografting would be impossible without the ability to perform HLA typing. The short arm of chromosome 6 contains a cluster of genes known as the major histocompatibility complex (MHC) or the HLA region (Fig. 21.5a). Genes in this region encode the HLA antigens and several other molecules including complement components, tumour necrosis factor (TNF) and proteins associated with antigen processing. HLA proteins are divided into two

types: class I and II (Table 21.3). Their role is to bind intracellular peptides and 'present' these to T lymphocytes for antigen recognition. Class I molecules (HLA-A, -B and -C) present antigen to CD8⁺ T cells and class II molecules (HLA-DR, -DQ and -DP) present to CD4⁺ T cells (Fig. 21.5b).

Class I antigens are present on most nucleated cells and on the cell surface they are associated with β_2 -microglobulin. Class II antigens have a more restricted tissue distribution and comprise α and β chains, both encoded by genes in the MHC locus. The inheritance of the four loci (HLA-A, -B, -C and -DR) is closely linked, one set of loci is inherited from each parent so that there is approximately a one in four chance of two siblings having identical HLA antigens (Fig. 21.6a). Crossing-over of genes during meiosis accounts for occasional unexpected disparities. The inheritance is independent of sex or blood group.

Human leucocyte antigen and transplantation

The natural role of HLA molecules is in directing T-lymphocyte responses and the greater the HLA mismatch the more severe is the immune response between transplanted cells. HLA typing is critical in donor selection for allogeneic SCT.

Minor histocompatibility antigens are peptides that are presented by HLA molecules and are able to act as antigens in SCT either because they are




Fig. 21.5 (a) The human leucocyte antigen (HLA) complex. (b) HLA class I and II molecules showing protein domains and bound peptide.

Table 21.3 The human leucocyte antigens.

	Class I	Class II
Antigens	HLA-A, -B, -C	HLA-DR, -DP, -DQ
Distribution	All nucleated cells, platelets	B lymphocytes Monocytes Macrophages Activated T cells
Structure	Large polypeptide chain (MHC coded) and a β_2 -microglobulin	Two polypeptide chains (α and β) both MHC coded
Interacts with	CD8 lymphocytes	CD4 lymphocytes

HLA, human leucocyte antigen; MHC, major histocompatibility complex.

polymorphic in the population or because they are encoded on the Y chromosome and therefore represent novel antigens when a female immune system engrafts in a male. They are likely to be important antigens in GVHD and the GVL reaction (see below).

HLA typing may be carried out by serological or molecular techniques. Serological testing involves the use of antibodies that are specific for individual HLA alleles or small families of alleles. Positivity may be detected by direct binding of a labelled antibody or by the use of complement to kill target cells that bind antibody (the two-stage lymphocytotoxicity test).

Molecular testing is performed on DNA. The most widely used methods use PCR sequence-specific primers or PCR sequence-specific oligonucleotide probes. Both methods use DNA primers or probes that react with polymorphic sequence motifs present within the nucleotide sequence of the HLA allele.

The nomenclature for HLA alleles is now standardized. A single antigenic specificity defined by serological typing (e.g. HLA-A) can be divided into different alleles by DNA sequencing. Each allele is given in numerical designation. The gene name is followed by an asterisk. The first two subsequent digits indicate the allele group. The third and fourth digits list subtypes. Subsequent digits indicate minor differences in non-coding regions. As an example, alleles at the HLA-A loci are written as HLA-A*0101

to HLA-A*8001, where the first two digits after the asterisk indicate the type of the allele and the last two digits indicate the subtype. The type often corresponds to the serological antigen carried by the alleles (e.g. HLA-A2 for the HLA-A*0201 to HLA-A*0230 alleles). The nomenclature for the class II genes is similar but complicated by the fact that there may be more than one HLA-DRB gene on each chromosome (Fig. 21.6b).

Further cellular tests of histocompatibility that are sometimes employed, particularly in unrelated donor SCT, are limiting dilution analyses of cytotoxic T-lymphocyte or helper T-lymphocyte precursors (CTLp and HTLp). A higher value indicates greater mismatch and appears to correlate with increased GVHD.

When searching for an unrelated donor the aim is to match HLA-A, -B and -DR between recipient and donor and this is then called a 6/6 match. Occasionally a single mismatch (5/6) can be tolerated but it is rare to accept a donor with a greater mismatch than this. An exception is when a donor with only a single HLA haplotype match, usually a parent or sibling, is used in a *haploidentical SCT*. Such transplants usually require a stem cell graft that is heavily depleted of T cells in order to limit the development of GVHD. There are over 4 million volunteer donors on international registries and the chance of identifying a matched unrelated donor for a patient lacking an HLA identical sibling (depending on the ethnic group) is usually greater than 50%.





Fig. 21.6 (a) An example of the possible pattern of inheritance of the A, B and DR (*DRB1*) series alleles of the human leucocyte antigen (HLA) complex. (b) Molecular genetics of the HLA class II gene complex. There are four major haplotypes of MHC class II genes in the population and each individual may have up to two (one on each chromosome). The *DRA1* gene codes for the DRα protein and the *DRB1*, *DRB3*, *DRB4* and *DRB5* genes encode DRβ

chains. Expression from the *DRB1* gene is higher than from the other genes. The number of alleles at each gene is shown underneath the gene in italics. Alleles at each locus have a standard nomenclature (e.g. the alleles at the *DRB1* gene are termed *DRB1*0101* to *DRB1*1608*). It is now known that the DR51, DR52 and DR53 antigens, which are defined by serological testing, are encoded from the *DRB5*, *DRB4* and *DRB3* genes, respectively.

Complications (Table 21.4)

Graft-versus-host disease

This is caused by donor-derived immune cells, particularly T lymphocytes, reacting against recipient

tissues. Its incidence is increased with increasing age of donor and recipient and if there is any degree of HLA mismatch between them. GVHD prophylaxis is usually given as ciclosporin (intravenously or orally for 6–12 months) and methotrexate (three

Table 21.4 Complications of stem cell transplantation.

Early (usually <100 days)	Late (usually >100 days)
Infections, especially bacterial, fungal, herpes simplex virus, CMV	Infections, especially varicella-zoster, capsulate bacteria
Haemorrhage	Chronic GVHD (arthritis, malabsorption, hepatitis, scleroderma, sicca syndrome, lichen planus, pulmonary disease, serous effusions)
Acute GVHD (skin, liver, gut)	Chronic pulmonary disease
Graft failure	Autoimmune disorders
Haemorrhagic cystitis	Cataract
Interstitial pneumonitis	Infertility
Others: veno-occlusive disease, cardiac failure	Second malignancies

CMV, cytomegalovirus; GVHD, graft-versus-host disease.

Table 21.5 Acute graft-versus-host disease: clinical staging (Seattle system).

Stage	Skin	Liver (bilirubin, $\mu\text{mol/L}$)	Gut (diarrhoea, L/day)
I	Rash <25%	20–35	0.5–1.0
II	Rash 25–50%	35–80	1.0–1.5
III	Erythroderma	80–150	1.5–2.5
IV	Bullae, desquamation	>150	>2.5; severe pain, ileus

Fig. 21.7 Widespread erythematous skin rash in acute graft-versus-host disease following bone marrow transplantation.



or four injections). An alternative is to remove T cells from the donor stem cell infusion. In addition, anti-T-cell antibodies may be given to the patient.

In *acute GVHD*, occurring in the first 100 days, the skin, gastrointestinal tract or liver are affected (Table 21.5). The skin rash typically affects the face, palms, soles and ears but may in severe cases affect the whole body (Fig. 21.7). The diagnosis is usually

confirmed by skin biopsy which shows initially single cell necrosis in the basal layer of the epidermis; lymphocyte infiltration may be scanty. Diarrhoea may lead to fluid and electrolyte depletion. Typically, bilirubin and alkaline phosphatase are raised but the other hepatic enzymes are relatively normal. Acute GVHD is usually treated by high doses of corticosteroids which are effective in the majority of cases.




Fig. 21.8 Time sequence for development of different types of infection following allogeneic bone marrow transplantation. CMV, cytomegalovirus; Gr+, Gr-, Gram-positive or -negative; GVHD, graft-versus-host disease; HSV, herpes simplex virus.


In *chronic GVHD*, occurring after 100 days and usually evolving from acute GVHD, these tissues are involved but also the joints and other serosal surfaces, the oral mucosa and lacrimal glands. Features of scleroderma, Sjögren's syndrome and lichen planus may develop. The immune system is impaired (including hyposplenism) with risk of infection. Malabsorption and pulmonary abnormalities are frequent. Drugs such as ciclosporin, azathioprine, mycophenolate mofetil, thalidomide or corticosteroids are used although response may be poor.

Infections


In the early post-transplant period, bacterial or fungal infections are frequent (Fig. 21.8). These may be reduced by reverse barrier nursing with laminar or positive pressure air flow and the use of skin and mouth antiseptics. Prophylactic therapy with aciclovir, antifungal agents and oral antibiotics is often

added. If a fever or other evidence of an infection occurs, broad-spectrum intravenous antibiotics are commenced immediately after blood cultures and other appropriate microbiological specimens have been taken. Failure of response to antibacterial agents is usually an indication to commence systemic antifungal therapy with amphotericin B, caspofungin or voriconazole (p. 151). Fungal infections, especially *Candida* and *Aspergillus* species (Fig. 21.9) are a particular problem because of the prolonged neutropenia. Fluconazole is effective in reducing the risk of *Candida* infection and itraconazole may provide some prophylaxis against both organisms. The standard formulation of amphotericin is nephrotoxic and the newer preparations such as liposomal amphotericin are better tolerated.

Viral infections, particularly with the herpes group of viruses, are frequent with herpes simplex, cytomegalovirus (CMV) and varicella zoster virus (VZV) occurring at different peak intervals (Fig. 21.8).



(a)



(b)

Fig. 21.9 (a) Chest radiograph showing an aspergilloma in a patient following stem cell transplantation. (b) Cytology of sputum illustrates the branching septate hyphae of *Aspergillus* (methenamine silver stain).

CMV presents a particular threat and is associated with a potentially fatal interstitial pneumonitis as well as with hepatitis and falling blood counts. The infection may be caused by reactivation of CMV in the recipient or a new infection transmitted by the

donor. In CMV-seronegative patients with CMV-seronegative donors, CMV-negative blood products or filtered blood must be given. Aciclovir may be useful in prophylaxis. Most centres screen patients regularly for evidence of CMV reactivation

following allogeneic transplantation using polymerase chain reaction (PCR) or antibody-based tests. If these tests become positive, ganciclovir may suppress the virus before disease occurs. Ganciclovir, foscarnet and CMV immunoglobulin may be tried for established CMV infection.


Pneumocystis carinii is another cause of pneumonitis that may be prevented by prophylactic co-trimoxazole. VZV infection is also frequent post-SCT but occurs later with a median onset at 4–5 months. Rarely, disseminated VZV infection occurs. Intravenous aciclovir is indicated. Epstein–Barr virus (EBV) infections and EBV-associated lymphoproliferative disease are less frequent after SCT than after solid organ transplants.

Interstitial pneumonitis

This is one of the most frequent causes of death post-SCT (Fig. 21.10). CMV is a frequent agent but other herpes viruses and *P. carinii* account for other cases; in many, no cause other than the previous radiation and chemotherapy can be implicated. Bronchoalveolar lavage or open lung biopsy may be needed to establish the diagnosis.


Blood product support

Platelet concentrates are given to maintain a count of $10 \times 10^9/L$ or more. Platelets and blood transfusions given in the post-transplant period must be irradiated prior to administration in order to kill any lymphocytes that might cause GVHD.



(a)

Fig. 21.10 (a) Chest radiograph showing interstitial pneumonitis following bone marrow transplantation. Widespread diffuse mottling can be seen. The patient had received total body irradiation and had grade III graft-versus-host disease. No infective cause of the pneumonitis was identified. Possible causes include pneumocystis,



(b)

(c)

cytomegalovirus, herpes zoster, fungal infection or a combination of these. (b) Sputum cytology: intranuclear CMV inclusion body in a pulmonary cell. Papanicolaou stain. (c) *Pneumocystis carinii* in bronchial washings, Gram–Weigert stain.

Other complications of allogeneic transplantation

Graft failure

The risk of graft failure is increased if the patient has aplastic anaemia or if T-cell depletion of donor marrow is used as GVHD prophylaxis. This suggests that donor T cells are needed to overcome host resistance to engraftment of stem cells.

Haemorrhagic cystitis

This is usually caused by the cyclophosphamide metabolite acrolein. Mesna is given in an attempt to prevent this. Certain viruses (e.g. adenovirus or polyomavirus) may also cause this complication.

Other complications

These include veno-occlusive disease of the liver (manifest as jaundice, hepatomegaly and ascites or weight gain) and cardiac failure as a result of the conditioning regime (especially high doses of cyclophosphamide) and previous chemotherapy on the heart. Haemolysis because of ABO incompatibility between donor and recipient may cause problems in the first weeks. Microangiopathic haemolytic anaemia may also occur.

Late complications

Relapse of the original disease (e.g. acute or chronic leukaemia) may occur. Bacterial infections are frequent, especially with Gram-negative or encapsulated organisms affecting the respiratory tract. Oral penicillin is given prophylactically to reduce this risk. VZV and fungal infections are also frequent. The use of prophylactic co-trimoxazole and oral aciclovir for 3–6 months reduces the risk of *Pneumocystis* and herpes infections, respectively.

Delayed pulmonary complications include restrictive pneumonitis and bronchiolitis obliterans. Endocrine complications include hypothyroidism, growth failure with low growth hormone levels in children, impaired sexual development and infertility. These endocrine problems are more marked if TBI has been used. Clinically apparent autoimmune disorders are infrequent and include myasthenia, rheumatoid arthritis, anaemia, thrombocytopenia or neutropenia. Autoantibodies are frequently detected in the absence of symptoms. Second malignancies

(especially non-Hodgkin's lymphoma) occur with a six- or sevenfold incidence compared with controls. CNS complications include neuropathies and eye problems caused by chronic GVHD (sicca syndrome) or cataracts.

Umbilical cord transplants

Fetal blood contains a large number of haemopoietic stem cells and umbilical cord blood has been used successfully as a source of stem cells. The main problem is the small amount of blood and therefore limited number of stem cells that can be collected from each sample and this limits the application of umbilical blood transplants to children or small adults. *Ex vivo* expansion of progenitor cells or administration of two or more cord units may be of value in the future. The immunological properties of cord blood cells are under investigation.

Graft-versus-leukaemia effect and donor leucocyte infusions

After allogeneic transplantation the donor immune system helps to eradicate the patient's leukaemia, a phenomenon known as the *graft-versus-leukaemia* (GVL) effect. Evidence includes the decreased relapse rate in patients with severe GVHD, the increased relapse rate in identical twins and, most convincingly, the ability of *donor leucocyte infusions* (DLI) to cure relapsed leukaemia in some patients. Graft-versus-lymphoma and -myeloma effects also exist. The principle of DLI is that peripheral blood mononuclear cells are collected from the original allograft donor and directly infused into the patient at the time of leukaemia relapse (Fig. 13.6).

There is a large difference in the outcome of different diseases treated by DLI. Chronic myeloid leukaemia (CML) is most sensitive whereas acute lymphoblastic leukaemia rarely responds. In CML the response to DLI is better in cases of early relapse. PCR is used to monitor serial blood samples for evidence of recurrence of the *BCR-ABL* transcript before karyotypic or clinical relapse occurs (Fig. 13.6). DLI can then be used in cases of molecular relapse. The response to DLI may take several weeks but usually results in a permanent cure. The mechanism is unclear but a T-cell-mediated

Table 21.6 Vectors used in gene therapy.

Vector	Properties
<i>Viral vectors</i>	
Retroviral	Integrate into DNA; infect variety of cell types. Only infect dividing cells
Adenovirus	Do not integrate, therefore transient expression. Infect dividing and non-dividing cells. Often provoke immune response
Adeno-associated virus	Limited integration sites
Herpes virus	Can carry large genes; no integration. Can infect resting cells
<i>Non-viral vectors</i>	
Liposomal	Relatively simple and cheap to make
Naked DNA	Low efficiency of entry into cells
Ballistic (gene gun)	

alloreactive immune response is likely to be a major component.

Non-myeloablative transplants

In order to reduce the morbidity and mortality of allogeneic transplantation, a number of low-intensity *non-myeloablative conditioning regimens* have been introduced which do not completely destroy the host bone marrow. These can include agents such as fludarabine, low-dose irradiation, antilymphocyte globulin and low doses of busulfan or cyclophosphamide. The aim in these 'mini-transplants' is to use enough immunosuppression to allow donor stem cells to engraft without completely eradicating host marrow stem cells. DLIs are commonly used at a late stage in order to encourage complete donor engraftment. Such regimes rely heavily on the ability of the GVL effect to cure the underlying malignant disease (Fig. 21.11) and extend the age range and increase the treatment indications for allogeneic transplantation.

Gene therapy and SCT

The ability to introduce novel genes into cells using appropriate vectors offers the opportunity to manipulate stem cells before reinfusion. Gene therapy vectors may be viral or non-viral and their relative properties are shown in Table 21.6. Many

different genes are being investigated for possible expression in cells such as haemopoietic stem cells and specific differentiated populations such as T lymphocytes. These include genes to correct specific inborn errors of metabolism, genes such as neomycin to 'mark' specific populations for subsequent analysis, 'suicide' genes such as thymidine kinase that render cells susceptible to ganciclovir and resistance genes to protect normal stem cells from high-dose chemotherapy. Unfortunately, haemopoietic stem cells are difficult to transfect because they are relatively rare and few are in cell cycle at any particular time. Therefore the sustained correction of metabolic disorders following the introduction of transduced autologous haemopoietic progenitors remains a difficult problem.

Bibliography

- Appelbaum F.R. (2003) The current status of hematopoietic cell transplantation. *Annu Rev Med* 54,491–512.
- Appelbaum F.R., Blume K.G., Forman S.J., Donnelly T.E. (2003) *Thomas' Hematopoietic Cell Transplantation*. Blackwell Science, Oxford.
- Copelan E.A. (2006) Hematopoietic stem-cell transplantation. *N Engl J Med* 354,1813–26.
- Diaconescu R., Flowers C.R., Storer B. *et al.* (2004) Morbidity and mortality with nonmyeloablative compared with myeloablative conditioning before hematopoietic cell transplantation from HLA-matched related disorders. *Blood* 104,1550–8.

- Hebart H. and Einsele H. (2004) Clinical aspects of CMV infection after stem cell transplantation. *Hum Immunol* 65,432–6.
- Jacobsohn D.A., Duerst R., Tse W., Kletzel M. (2004) Reduced intensity haemopoietic stem-cell transplantation for treatment of non-malignant diseases in children. *Lancet* 364,156–62.
- Little A-M., Marsh S.G.E., Madrigal J.A. (2005) Histocompatibility. In: Hoffbrand A.V., Tuddenham E.G.D., Catovsky D. (eds) *Postgraduate Haematology*, 5th edn. Blackwell Publishing, Oxford: pp. 395–418.
- Mathur A. and Martin J.F. (2004) Stem cells and repair of the heart. *Lancet* 364,183–92.
- Peggs K.S. and Mackinnon S. (2004) Immune reconstitution following haematopoietic stem cell transplantation. *Br J Haematol* 124,407–20.
- Peggs K.S., Mackinnon S., Linch D.C. (2005) The role of allogeneic transplantation in non-Hodgkin's lymphoma. *Br J Haematol* 128,153–68.
- Wollert K.C., Meyer G.P., Lote J. *et al.* (2004) Intracoronary autologous bone-marrow cell transfer after myocardial infarction: the BOOST randomised controlled clinical trial. *Lancet* 364,141–8.

CHAPTER 22

22

Platelets, blood coagulation and haemostasis

Components of haemostatic response, 265

Blood coagulation, 270

Fibrinolysis, 274

Haemostatic response, 275

Tests of haemostatic function, 276

Bibliography, 277

The normal haemostatic response to vascular damage depends on closely linked interaction between the blood vessel wall, circulating platelets and blood coagulation factors (Fig. 22.1).

An efficient and rapid mechanism for stopping bleeding from sites of blood vessel injury is clearly

essential for survival. Nevertheless, such a response needs to be tightly controlled to prevent extensive clots developing and to break down such clots once damage is repaired. The haemostatic system thus represents a delicate balance between procoagulant and anticoagulant mechanisms allied to a process for




Fig. 22.1 The involvement of blood vessels, platelets and blood coagulation in haemostasis. ADP, adenosine diphosphate.

fibrinolysis. The five major components involved are platelets, coagulation factors, coagulation inhibitors, fibrinolysis and blood vessels. These are described later in the haemostatic response section on p. 275.

Components of haemostatic response

Platelets

Platelet production

Platelets are produced in the bone marrow by fragmentation of the cytoplasm of megakaryocytes, one of the largest cells in the body. The precursor of the megakaryocyte—the megakaryoblast—arises by a process of differentiation from the haemopoietic stem cell (Fig. 22.2). The megakaryocyte matures by endomitotic synchronous replication (i.e. DNA replication in the absence of nuclear or cytoplasmic division) enlarging the cytoplasmic volume as the number of nuclear lobes increase in multiples of two. A picture of mature polyploid megakaryocytes is shown in Fig. 22.3. Very early on invaginations of plasma membrane are seen, called the demarcation membrane, which evolves through the development of the megakaryocyte into a highly branched network. At a variable stage in development, most commonly at the eight nucleus stage, the cytoplasm becomes granular. Mature megakaryocytes are extremely large, with an eccentric placed single lobu-

lated nucleus and a low nuclear to cytoplasmic ratio. Platelets form by fragmentation of megakaryocyte cytoplasm, approximately each megakaryocyte giving rise to 1000–5000 platelets. The time interval from differentiation of the human stem cell to the production of platelets averages approximately 10 days.

Thrombopoietin is the major regulator of platelet production and is constitutively produced by the liver and kidneys. Thrombopoietin increases the number and rate of maturation of megakaryocytes via c-Mpl receptor. Platelet levels start to rise 6 days after the start of therapy and remain high for 7–10 days. Unfortunately, thrombopoietin is not available for routine clinical practice. Platelets also have c-Mpl receptors for thrombopoietin and remove it from the circulation. Therefore, levels are high in thrombocytopenia as a result of marrow aplasia and vice versa.

The normal platelet count is approximately $250 \times 10^9/\text{L}$ (range $150\text{--}400 \times 10^9/\text{L}$) and the normal platelet lifespan is 7–10 days. Up to one-third of the marrow output of platelets may be trapped at any one time in the normal spleen but this rises to 90% in cases of massive splenomegaly (p. 286).

Platelet structure

Platelets are extremely small and discoid, $3.0 \times 0.5 \mu\text{m}$ in diameter, with a mean volume 7–11 fL. The ultrastructure of platelets is represented in




Fig. 22.2 Simplified diagram to illustrate platelet production from megakaryocytes.




Fig. 22.3 Megakaryocytes: (a) immature form with basophilic cytoplasm; (b) mature form with many nuclear lobes and pronounced granulation of the cytoplasm.




Fig. 22.4 The ultrastructure of platelets. ADP, adenosine diphosphate; PF, platelet factor; VWF, von Willebrand factor.

Fig. 22.4. The glycoproteins of the surface coat are particularly important in the platelet reactions of adhesion and aggregation which are the initial events leading to platelet plug formation during haemostasis. Adhesion to collagen is facilitated by glycoprotein Ia (GPIa). Glycoproteins Ib (defective in Bernard-Soulier syndrome) and IIb/IIIa (defective in thrombasthenia) are important in the attachment of platelets to von Willebrand factor (VWF) and hence to vascular subendothelium (Fig. 22.5) where metabolic interactions occur (Fig. 22.6). The binding site for IIb/IIIa is also the receptor for

fibrinogen which is important in platelet-platelet aggregation.

The plasma membrane invaginates into the platelet interior to form an open membrane (canalicular) system which provides a large reactive surface to which the plasma coagulation proteins may be selectively absorbed. The membrane phospholipids (previously known as platelet factor 3) are of particular importance in the conversion of coagulation factor X to Xa and prothrombin (factor II) to thrombin (factor IIa) (Fig. 22.7).

The platelet contains three types of storage





Fig. 22.5 Platelet adhesion. The binding of glycoprotein (GP) Ib (which consists of four proteins: GPIb α , GPIb β , GPIX, GPV) to von Willebrand factor leads to adhesion to the subendothelium and also exposes the GPIIb/IIIa ($\alpha_{IIb}\beta_3$ integrin) binding sites to fibrinogen and von Willebrand factor leading to platelet aggregation. The GPIa site permits direct adhesion to collagen.

Fig. 22.6 The synthesis of prostacyclin and thromboxane. The opposing effects of these agents are mediated by changes in the concentration of cyclic adenosine monophosphate (cAMP) in platelets via stimulation or inhibition of the enzyme adenylate cyclase. cAMP controls the concentration of free calcium ions in the platelet which are important in the processes which cause adhesion and aggregation. High levels of cAMP lead to low free calcium ion concentrations and prevent aggregation and adhesion. ATP, adenosine triphosphate; Ca, calcium; PG, prostaglandin (G₂ and H₂).



granules: dense, α and lysosomes. The more frequent specific α granules contain a heparin antagonist (PF4), platelet-derived growth factor (PDGF), β -thromboglobulin, fibrinogen, VWF and other clotting factors. Dense granules are less common and

contain adenosine diphosphate (ADP), adenosine triphosphate (ATP), 5-hydroxytryptamine (5-HT) and calcium. Lysosomes contain hydrolytic enzymes and peroxisomes contain catalase. During the release reaction described below, the contents of




Fig. 22.7 The pathway of blood coagulation initiated by tissue factor (TF) on the cell surface. When plasma comes into contact with TF, factor VII binds to TF. The complex of TF and activated VII (VIIa) activates X and IX. TF pathway inhibitor (TFPI) is an important inhibitor of TF/VIIa. VIIa-IXa complex greatly amplifies Xa production from X. The generation of thrombin from prothrombin by the action of Xa-Va complex leads to fibrin formation. Thrombin also activates XI (dashed line), V and XIII. Thrombin cleaves VIII from its carrier von Willebrand factor (VWF), greatly increasing the formation of VIIa-IXa and hence of Xa-Va. Pale green, serine proteases; yellow, cofactors.

the granules are discharged into the open canalicular system.

Platelets are also rich in signalling and cytoskeletal proteins which support the rapid switch from quiescent to activation that follows vessel damage.

Platelet antigens

Several platelet surface proteins have been found to be important antigens in platelet-specific autoimmunity and they have been termed human platelet antigens (HPA). In most cases, two different alleles exist, termed a or b alleles (e.g. HPA-1a). Platelets also express ABO and human leucocyte antigen (HLA) class I but not class II antigens.

Platelet function

The main function of platelets is the formation of mechanical plugs during the normal haemostatic response to vascular injury. In the absence of platelets, spontaneous leakage of blood through small vessels may occur. The immobilization of platelets at the sites of vascular injury requires specific platelet-vessel wall (adhesion) and platelet-platelet (aggregation) interactions. The blood flow conditions determine the specific receptor ligand interactions.

Platelet adhesion and activation

Following blood vessel injury, platelets adhere to exposed subendothelial matrix proteins via specific adhesive glycoproteins (GP). Under condition high shear, e.g. arterioles, the exposed subendothelial matrix is initially coated with VWF multimer. The platelets then make contact with VWF via the GPIIb-IIIa-V complex on platelets. This initiates platelet rolling in the direction of blood flow over the exposed VWF with activation of GPIIb/IIIa receptor. Firm adhesion is established by the stronger interaction of other glycoproteins including activated GPIIb/IIIa with VWF and GPVI and integrin $\alpha 1/\beta 2$ with collagen and other components of the subendothelial matrix. Under static or low shear conditions, platelets adhere predominantly to collagen of the subendothelium. Collagen binds initially to GPIa/IIa, cross-links many of these integrin molecules, and in this way activates platelets.

This ligand receptor binding results in a complex cascade of signals which result in platelet activation. The events that follow are shape change and spreading, activation of GPIIb/IIIa and granule secretion. Platelets become more spherical and extrude long pseudopodia which enhance platelet vessel wall

and platelet–platelet interaction. The end result of spreading is a flattened spread out platelet with granules and organelles in the centre, resulting in a characteristic fried egg appearance. These changes are brought about by the actin cytoskeleton. The granules are secreted from the centre of the cell.

Von Willebrand factor VWF is involved in platelet adhesion to the vessel wall and to other platelets (aggregation) (Fig. 22.5). It also carries factor VIII (see below) and used to be referred to as factor VIII-related antigen (VIII-Rag). It is a large cysteine-rich glycoprotein, with multimers made up on average of 2–50 subunits, with a molecular weight (MW) of $0.8\text{--}20 \times 10^6$. VWF is encoded by a gene on chromosome 12 and is synthesized both in endothelial cells and megakaryocytes, and stored in Weibel–Palade bodies and platelet α granules respectively. Plasma VWF is almost entirely derived from endothelial cells, with two distinct pathways of secretion. The majority is continuously secreted and a minority is stored in Weibel–Palade bodies. The stored VWF can rise the plasma levels and it can be released under the influence of several secretagogues, like stress, exercise, adrenaline and infusion of desmopressin (1-deamino-8-D-arginine vasopressin, DDAVP). The VWF released from Weibel–Palade bodies is in the form of large and ultra large multimers, the most adhesive and reactive form of VWF. They are in turn cleaved in plasma to monomeric VWF and smaller multimers by the specific plasma metalloprotease, ADAMTS-13.

Platelet aggregation

It is characterized by cross-linking of platelets through active GPIIb/IIIa receptors with fibrinogen bridges. A resting platelet has about 50–80 000 GPIIb/IIIa receptors, which do not bind fibrinogen, VWF or other ligands. Stimulation of a platelet leads to an increase in GPIIb/IIIa molecules, due to binding of α -granule membrane (rich in receptors) with the plasma membrane, activation of surface-exposed GPIIb/IIIa, enabling platelet cross-linking with fibrinogen bridges. Binding brings about molecular conformational changes resulting in a firm connection and further activation of the platelet.

Platelet release reaction and amplification

Primary activation by various agonists induces

intracellular signalling, leading to the release of α and δ -granules. α -Granule contents play an important role in platelet aggregate formation and stabilization and, in addition, the ADP released from dense granules plays a major positive feedback role in promoting platelet activation. TXA₂ is the second of the two major platelet positive feedback loops important in secondary amplification of platelet activation to form a stable platelet aggregate. It is formed *de novo* upon activation of cytosolic phospholipase A₂ (PL_{A2}) which is the rate limiting step. This liberates arachidonic acid from the membrane phospholipids, and is metabolized by cyclooxygenase to TXA₂. It is a labile substance and lowers platelet cyclic adenosine monophosphate (cAMP) levels and initiates the release reaction (Fig. 22.6). Thromboxane A₂ not only potentiates platelet aggregation, but also has powerful vasoconstrictive activity. The release reaction is inhibited by substances that increase the level of platelet cAMP. One such substance is the prostaglandin prostacyclin (PGI₂) which is synthesized by vascular endothelial cells. It is a potent inhibitor of platelet aggregation and prevents their deposition on normal vascular endothelium.

Clot formation and retraction

The highly localized enhancement of ongoing platelet activation by ADP and TXA₂ results in a platelet plug large enough to plug the area of endothelial injury. In this platelet plug the platelets are completely degranulated and adherent to each other. This is followed by clot retraction which is mediated by GPIIb/IIIa receptors which link the cytoplasmic actin filaments to the surface bound fibrin polymers.

Platelet procoagulant activity

After platelet aggregation and release, the exposed membrane phospholipid (platelet factor 3) is available for two reactions in the coagulation cascade. Both phospholipid-mediated reactions are calcium-ion dependent. The first (tenase) involves factors IXa, VIIIa and X in the formation of factor Xa (Fig. 22.7). The second (prothrombinase) results in the formation of thrombin from the interaction of factors Xa, Va and prothrombin (II). The phospholipid surface forms an ideal template for the crucial concentration and orientation of these proteins.




Fig. 22.8 The endothelial cell forms a barrier between platelets and plasma clotting factors and the subendothelial connected tissues. Endothelial cells produce substances that can initiate coagulation, cause vasodilatation, inhibit platelet aggregation or haemostasis, or activate fibrinolysis.

Growth factor

PDGF found in the specific granules of platelets stimulates vascular smooth muscle cells to multiply and this may hasten vascular healing following injury.

Natural inhibitors of platelet function

Nitric oxide (NO) is constitutively released from endothelial cells and also from macrophages and platelets. It has a short half-life of 3–5 s. It inhibits platelet activation and promotes vasodilatation. Prostacyclin synthesized by endothelial cells also inhibits platelet function (Fig. 22.8) and causes vasodilatation by raising cyclic guanosine monophosphate (GMP) levels. The transmembrane protein PECAM-1 is expressed also on endothelial cells. It is its own ligand and inhibits platelet activation by collagen.

Blood coagulation

The coagulation cascade

Blood coagulation involves a biological amplification system in which relatively few initiation substances sequentially activate by proteolysis a cascade of circulating precursor proteins (the coagulation factor enzymes) which culminates in

the generation of thrombin; this, in turn, converts soluble plasma fibrinogen into fibrin (Fig. 22.7). Fibrin enmeshes the platelet aggregates at the sites of vascular injury and converts the unstable primary platelet plugs to firm, definitive and stable haemostatic plugs. A list of the coagulation factors appears in Table 22.1. They are assembled from a small number of molecules or domains (Fig. 22.9). The operation of this enzyme cascade requires local concentration of circulating coagulation factors at the site of injury.

Surface-mediated reactions occur on exposed collagen, platelet phospholipid and tissue factor. With the exception of fibrinogen, which is the fibrin clot subunit, the coagulation factors are either enzyme precursors or cofactors (Table 22.1). All the enzymes, except factor XIII, are serine proteases (i.e. their ability to hydrolyse peptide bonds depends upon the amino acid serine at their active centre; Fig. 22.10). The scale of amplification achieved in this system is dramatic, (e.g. 1 mol of activated factor XI through sequential activation of factors IX, X and prothrombin may generate up to 2×10^8 mol of fibrin).

Coagulation *in vivo*

The generation of thrombin *in vivo* is a complex network of amplification and negative feedback loops

Table 22.1 The coagulation factors.

Factor number	Descriptive name	Active form
I	Fibrinogen	Fibrin subunit
II	Prothrombin	Serine protease
III	Tissue factor	Receptor/cofactor*
V	Labile factor	Cofactor
VII	Proconvertin	Serine protease
VIII	Antihaemophilic factor	Cofactor
IX	Christmas factor	Serine protease
X	Stuart–Prower factor	Serine protease
XI	Plasma thromboplastin antecedent	Serine protease
XII	Hageman (contact) factor	Serine protease
XIII	Fibrin-stabilizing factor	Transglutaminase
	Prekallikrein (Fletcher factor)	Serine protease
	HMWK (Fitzgerald factor)	Cofactor*

HMWK, high molecular weight kininogen.

* Active without proteolytic modification.

to ensure a localized and limited production. The generation of thrombin is dependant on three enzyme complexes, each consisting of protease, cofactor, phospholipids (PL) and calcium. They are extrinsic Xase (VIIa, TF, PL, Ca^{2+}) and intrinsic Xase (IXa, VIIIa, PL, Ca^{2+}) generating FXa, prothrombinase complex (Xa, Va, PL, Ca^{2+}) generating thrombin. The generation of thrombin following vascular injury occurs in two waves of very different magnitude with different functions. During the initiation phase small amounts are generated (picomolar concentrations) which prepares the coagulation cascade for the second larger thrombin burst in the amplification when micromolar concentrations are produced, i.e. a million-fold higher concentration than produced during the initiation phase.

Initiation

Coagulation is initiated by the interaction of the membrane bound TF exposed by vascular injury, with plasma factor VIIa. One to two per cent of the total factor VII circulates in the activated form, but does not express proteolytic activity unless bound to TF. The factor VIIa-tissue factor (extrinsic factor Xase) complex activates both factor IX and factor X. The factor Xa, in the absence of its cofactor, forms small amounts of thrombin from prothrombin. This is insufficient to initiate significant fibrin polymer-

ization, it activates the coenzymes, factor V and factor VIII, platelets and factor XI.

Amplification

The initiation pathway or extrinsic Xase is rapidly inactivated by TFPI which forms a quaternary complex of VIIa, TF, Xa and TFPI. The thrombin generation is now dependant on the traditional intrinsic pathway which has been primed by the small amounts of thrombin generated during initiation. In the amplification phase the intrinsic Xase formed by IXa and VIIIa on phospholipid surface in the presence of Ca^{2+} activates sufficient Xa which then in combination with Va, PL and Ca^{2+} forms the prothrombinase complex and results in the explosive generation of thrombin which acts on fibrinogen to form the fibrin clot.

In the 'classic' pathway formulated to explain *in vitro* coagulation testing, initiation of the pathway required contact reactions between factor XII, kallikrein and high molecular weight kininogen (HMWK) leading to the activation of factor XI. However, the lack of abnormal bleeding in individuals with hereditary deficiencies of these contact factors suggests that these reactions are not required for physiological coagulation *in vivo*.

Factor XI does not seem to have a role in the physiological initiation of coagulation. It has a




Fig. 22.9 Domains of the enzymes, receptors and cofactors involved in blood coagulation and regulation. The components of blood coagulation are proenzymes, procofactors and regulator proteins. The proenzymes, including protein C, contain a catalytic domain, an activation region and a signal peptide. The vitamin K-dependent proteins include a propeptide and a γ -carboxyglutamic acid (Gla) domain. Other important domains include the epidermal growth factor-like (EGF) domain, the kringle domain and the repeat-sequence

domain. Tissue factor is an integral membrane protein unrelated to other known proteins. Factors V and VIII have marked similarities in structure. Sites of intracellular peptide bonds cleaved during synthesis are indicated by thin arrows, and sites of peptide bonds cleaved during protein activation are indicated by thick arrows. The transmembrane domain of tissue factor is shown within the phospholipid bilayer. (From B. Furie and B.C. Furie 1992, courtesy of the *New England Journal of Medicine* with permission.)




Fig. 22.10 Serine (Se) protease activity. This example shows the activation of factor X by factor IX.

supplementary role in the activation of factor IX (see above) and may be important at major sites of trauma or at operations and potentially causes excess bleeding in factor XI deficient individuals.

Thrombin hydrolyses fibrinogen, releasing fibrinopeptides A and B to form fibrin monomers (Fig. 22.11). Fibrin monomers link spontaneously by hydrogen bonds to form a loose insoluble fibrin

Table 22.2 The coagulation factors.

Factor	Plasma half-life (h)	Plasma concentration (mg/L)	Comments
II	65	100	Prothrombin group: vitamin K needed for synthesis; require Ca^{2+} for activation Thrombin interacts with them; increase in inflammation, pregnancy, oral contraceptives
VII	5	0.5	
IX	25	5	
X	40	10	
I	90	3000	
V	15	10	
VIII	10	0.1	
XI	45	5	
XII	50	30	




Fig. 22.11 The formation and stabilization of fibrin.

polymer. Factor XIII is also activated by thrombin together with calcium. Activated factor XIII stabilizes the fibrin polymers with the formation of covalent bond cross-links.

Fibrinogen has a MW 340 000 and consists of two identical subunits, each containing three dissimilar polypeptide chains ($\text{A}\alpha$, $\text{B}\beta$ and γ) which are linked by disulphide bonds. After cleavage by thrombin of fibrinopeptides A and B, fibrin monomer consists of three paired α , β and γ chains.

Some of the properties of the coagulation factors are listed in Table 22.2. The activity of factors II, VII, IX and X is dependent upon vitamin K which is responsible for carboxylation of a number of terminal glutamic acid residues on each of these molecules (see Fig. 24.7).

The serine protease coagulation factors along with those of the fibrinolytic system (see below)

have a high degree of homology and contain characteristic structural domains (Fig. 22.9), such as the kringles which are concerned with substrate binding and the carboxylated glutamic acid (Gla) residues which bind to phospholipid. There are also regions of homology with fibronectin (finger regions) and with epidermal growth factor. Although factor VIII and V cofactors are not protease enzymes, they circulate in a precursor form that requires limited cleavage by thrombin for expression of full cofactor activity.

Physiological limitation of blood coagulation

Unchecked blood coagulation would lead to dangerous occlusion of blood vessels (thrombosis) if the protective mechanisms of coagulation factor inhibitors, blood flow and fibrinolysis were not in operation.

Coagulation factor inhibitors

It is important that the effect of thrombin is limited to the site of injury. The first inhibitor to act is tissue factor pathway inhibitor (TFPI) which is synthesized in endothelial cells and is present in plasma and platelets and accumulates at the site of injury caused by local platelet activation. This inhibits Xa and VIIa and tissue factor to limit the main *in vivo* pathway by forming the quaternary complex. There is direct inactivation of thrombin and other serine protease factors by other circulating inhibitors of which antithrombin is the most potent. It inactivates serine proteases by combining with them by peptide bonding to form high molecular weight stable complexes




Fig. 22.12 Activation and action of protein C by thrombin which has bound to thrombomodulin on the endothelial cell surface. Protein S is a cofactor that facilitates binding of activated protein C to the platelet surface. The inactivation of factors Va and VIIIa results in the inhibition of blood coagulation. The inactivation of tissue plasminogen activator inhibitor (tPAI) enhances fibrinolysis. EPCR, endothelial protein C receptor.

(Fig. 25.6). Heparin potentiates its action markedly. Another protein, heparin cofactor II, also inhibits thrombin. α_2 -Macroglobulins, α_2 -antiplasmin, C_1 esterase inhibitor and α_1 -antitrypsin also exert inhibitory effects on circulating serine proteases.

Protein C and protein S

These are inhibitors of coagulation cofactors V and VIII. Thrombin binds to an endothelial cell surface receptor, thrombomodulin. The resulting complex activates the vitamin K-dependent serine protease protein C which is able to destroy activated factors V and VIII, thus preventing further thrombin generation. The action of protein C is enhanced by another vitamin K-dependent protein, S, which binds protein C to the platelet surface (Fig. 22.12). An endothelial protein C receptor localizes protein C to the endothelial surface, promoting protein C activation by the thrombin–thrombomodulin complex (Fig. 22.12). In addition, activated protein C enhances fibrinolysis (Fig. 22.12).

As with other serine proteases, activated protein C is subject to inactivation by serum protease inactivators (serpins), e.g. antithrombin.

Blood flow

At the periphery of a damaged area of tissue, blood flow rapidly achieves a dilution and dispersal of activated factors before fibrin formation has occurred. Activated factors are destroyed by liver parenchy-

mal cells and particulate matter is removed by liver Kupffer cells and other reticuloendothelial cells.

Fibrinolysis

Fibrinolysis (like coagulation) is a normal haemostatic response to vascular injury. Plasminogen, a β -globulin proenzyme in blood and tissue fluid, is converted to the serine protease plasmin by activators either from the vessel wall (intrinsic activation) or from the tissues (extrinsic activation) (Fig. 22.13). The most important route follows the release of tissue plasminogen activator (tPA) from endothelial cells. tPA is a serine protease that binds to fibrin. This enhances its capacity to convert thrombus-bound plasminogen into plasmin. This fibrin dependence of tPA action strongly localizes plasmin generation by tPA to the fibrin clot. Release of tPA




Fig. 22.13 The fibrinolytic system. tPA, tissue plasminogen activator.

occurs after such stimuli as trauma, exercise or emotional stress. Activated protein C stimulates fibrinolysis by destroying plasma inhibitors of tPA (Fig. 22.12). On the other hand, thrombin inhibits fibrinolysis by activating thrombin-activated fibrinolysis inhibitor (TAFI).

Plasmin generation at the site of injury limits the extent of the evolving thrombus. The split products of fibrinolysis are also competitive inhibitors of thrombin and fibrin polymerization. Normally, α_2 -antiplasmin inhibits any local free plasmin.

Fibrinolytic agents are widely used in clinical practice (p. 317). Therapeutic recombinant tPA has been synthesized using recombinant DNA technology. The bacterial agent streptokinase is a peptide produced by haemolytic streptococci and forms a complex with plasminogen, which converts other plasminogen molecules to plasmin. Urokinase is a tPA initially isolated from human urine.

Plasmin is capable of digesting fibrinogen, fibrin, factors V and VIII and many other proteins. Cleavage of peptide bonds in fibrin and fibrinogen produces a variety of split (degradation) products (Fig. 22.13). Large amounts of the smallest fragments D and E can be detected in the plasma of patients with disseminated intravascular coagulation (p. 298).

Inactivation of plasmin

Tissue plasminogen activator is inactivated by plasminogen activator inhibitor (PAI). Circulating plasmin is inactivated by potent inhibitors α_2 -antiplasmin and α_2 -macroglobulin.

Endothelial cells

The endothelial cell has an active role in the maintenance of vascular integrity. This cell provides the basement membrane that normally separates collagen, elastin and fibronectin of the subendothelial connective tissue from the circulating blood (Fig. 22.8). Loss or damage to the endothelium results in both haemorrhage and activation of the haemostatic mechanism. The endothelial cell also has a potent inhibitory influence on the haemostatic response, largely through the synthesis of PGI_2 and NO, which have vasodilatory properties and inhibit platelet aggregation. In contrast, endothelins are a family of

vasoactive peptides that can activate fibrinolysis via the release of tPA. Synthesis of tissue factor which initiates haemostasis only occurs in endothelial cells following activation and its natural inhibitor, tFPI, is also synthesized. Synthesis of prostacyclin, VWF, plasminogen activator, antithrombin and thrombomodulin, the surface protein responsible for activation of protein C, provides agents that are vital to both platelet reactions and blood coagulation (Fig. 22.8).

Haemostatic response

Vasoconstriction

An immediate vasoconstriction of the injured vessel and reflex constriction of adjacent small arteries and arterioles is responsible for an initial slowing of blood flow to the area of injury. When there is widespread damage this vascular reaction prevents exsanguination. The reduced blood flow allows contact activation of platelets and coagulation factors. The vasoactive amines and thromboxane A_2 liberated from platelets, and the fibrinopeptides liberated during fibrin formation, also have vasoconstrictive activity (Fig. 22.1).

Platelet reactions and primary haemostatic plug formation

Following a break in the endothelial lining, there is an initial adherence of platelets to exposed connective tissue, potentiated by VWF. Collagen exposure and thrombin produced at the site of injury cause the adherent platelets to release their granule contents and also activate platelet prostaglandin synthesis leading to the formation of thromboxane A_2 . Released ADP causes platelets to swell and aggregate. Additional platelets from the circulating blood are drawn to the area of injury. This continuing platelet aggregation promotes the growth of the haemostatic plug which soon covers the exposed connective tissue. The unstable primary haemostatic plug produced by these platelet reactions in the first minute or so following injury is usually sufficient to provide temporary control of bleeding. It seems likely that prostacyclin, produced by endothelial and smooth muscle cells in the vessel wall adjacent to the area of damage, is important in limiting the extent of the initial platelet plug.

Stabilization of the platelet plug by fibrin

Definitive haemostasis is achieved when fibrin formed by blood coagulation is added to the platelet mass and by platelet-induced clot retraction/compaction.

Following vascular injury, the formation of extrinsic Xase (VIIa, TF, PL and Ca^{2+}) initiates the coagulation cascade. Platelet aggregation and release reactions accelerate the coagulation process by providing abundant membrane phospholipid. Thrombin generated at the injury site converts soluble plasma fibrinogen into fibrin, potentiates platelet aggregation and secretion and also activates factor XI and XIII and cofactors V and VIII. The fibrin component of the haemostatic plug increases as the fused platelets autolyse and after a few hours the entire haemostatic plug is transformed into a solid mass of cross-linked fibrin. Nevertheless, because of incorporation of plasminogen and tPA (p. 273), this plug begins to autodigest during the same time frame.

Tests of haemostatic function

Defective haemostasis with abnormal bleeding may result from:

- 1 A vascular disorder;
- 2 Thrombocytopenia or a disorder of platelet function; or
- 3 Defective blood coagulation.

A number of simple tests are employed to assess the platelet, vessel wall and coagulation components of haemostasis.

Blood count and blood film examination

As thrombocytopenia is a common cause of abnormal bleeding, patients with suspected bleeding disorders should initially have a blood count including platelet count and blood film examination. In addition to establishing the presence of thrombocytopenia, the cause may be obvious (e.g. acute leukaemia).

Screening tests of blood coagulation

Screening tests provide an assessment of the 'extrinsic' and 'intrinsic' systems of blood coagulation and also the central conversion of fibrinogen to fibrin (Table 22.3).

The prothrombin time (PT) measures factors VII, X, V, prothrombin and fibrinogen. Tissue thromboplastin (a brain extract) and calcium are added to citrated plasma. The normal time for clotting is 10–14 s. It may be expressed as the international normalized ratio (INR) (p. 314).

The activated partial thromboplastin time (APTT) measures factors VIII, IX, XI and XII in addition to factors X, V, prothrombin and fibrinogen. Three substances—phospholipid, a surface activator (e.g. kaolin) and calcium—are added to citrated plasma. The normal time for clotting is approximately 30–40 s.

Prolonged clotting times in the PT and APTT because of factor deficiency are corrected by the addition of normal plasma to the test plasma (50 : 50 mix). If there is no correction or incomplete

Table 22.3 Screening tests used in the diagnosis of coagulation disorders.

Screening tests	Abnormalities indicated by prolongation	Most common cause of disorder
Thrombin time (TT)	Deficiency or abnormality of fibrinogen or inhibition of thrombin by heparin or FDPs	Disseminated intravascular coagulation Heparin therapy
Prothrombin time (PT)	Deficiency or inhibition of one or more of the following coagulation factors: VII, X, V, II, fibrinogen	Liver disease Warfarin therapy DIC
Activated partial thromboplastin time (APTT or PTTK)	Deficiency or inhibition of one or more of the following coagulation factors: XII, XI, IX (Christmas disease), VIII (haemophilia), X, V, II, fibrinogen	Haemophilia, Christmas disease (+ conditions above)
Fibrinogen quantitation	Fibrinogen deficiency	Disseminated intravascular coagulation, liver disease

FDPs, fibrin degradation products.

NB. Platelet count and the tests of platelet function are also used in screening patients with a bleeding disorder.

correction with normal plasma, the presence of an inhibitor of coagulation is suspected.

The thrombin (clotting) time (TT) is sensitive to a deficiency of fibrinogen or inhibition of thrombin. Diluted bovine thrombin is added to citrated plasma at a concentration giving a clotting time of 14–16 s with normal subjects.

Specific assays of coagulation factors

Most factor assays are based on an APTT or PT in which all factors except the one to be measured are present in the substrate plasma. This usually requires a supply of plasma from patients with hereditary deficiency of the factor in question or artificially produced factor-deficient plasma. The corrective effect of the unknown plasma on the prolonged clotting time of the deficient substrate plasma is then compared with the corrective effect of normal plasma. Results are expressed as a percentage of normal activity.

A number of chemical, chromogenic and immunological methods are available for quantification of other proteins such as fibrinogen, VWF and factor VIII. Factor XIII activity can be assessed by testing for clot solubility in urea.

Bleeding time

The bleeding time is a useful test for abnormal platelet function including the diagnosis of VWF deficiency. It has largely been replaced by the platelet function analysis-100 (PFA-100) test (see below). It will be prolonged in thrombocytopenia but is normal in vascular causes of abnormal bleeding. The test involves the application of pressure to the upper arm with a blood pressure cuff, after which small incisions are made in the flexor surface forearm skin. Bleeding stops normally in 3–8 min.

Tests of platelet function

The most valuable investigation is platelet aggregometry which measures the fall in light absorbance in platelet-rich plasma as platelets aggregate. Initial (primary) aggregation is caused by an external agent, the secondary response to aggregating agents released from the platelets themselves. The five external aggregating agents most commonly used

are ADP, collagen, ristocetin, arachidonic acid and adrenaline. The pattern of response to each agent helps to make the diagnosis (see Fig. 23.1). Flow cytometry is now increasingly used in routine practice to identify platelet glycoprotein defects.

In the PFA-100 test, citrated blood is aspirated through a capillary tube onto a membrane coated with collagen/ADP or collagen/adrenaline. Blood flow is maintained. Platelets begin to adhere and aggregate, primarily via VWF interactions with GPIb and GPIIb/IIIa, resulting in occlusion of the aperture.

The PFA-100 analysis may give false negative results with relatively common platelet defects. Full platelet function tests and VWF screening may be required to exclude abnormal platelet function, even if the PFA-100 test is normal.

Test of fibrinolysis

Increased levels of circulating plasminogen activator may be detected by demonstrating shortened euglobulin clot lysis times. A number of immunological methods are available to detect fibrinogen or fibrin degradation products (including D-dimers) in serum. In patients with enhanced fibrinolysis, low levels of circulating plasminogen may be detected.

Bibliography

- Aird W.C. (2003) Endothelial cell heterogeneity. *Crit Care Med* 31, S221–30.
- Dahlback B. and Villoutreic B.O. (2003) Molecular recognition in the protein C anticoagulant pathway. *J Thromb Haemost* 1, 1525–34.
- Harrison P. (2005) The role of PFA-100 testing in the investigation and management of haemostatic defects in children and adults. *Br J Haematol* 130, 3–10.
- Italiano J.E. Jr. and Shivdasani R.A. (2003) Megakaryocytes and beyond: the birth of platelets. *J Thromb Haemost* 1, 1174–82.
- Kemball-Cook, G., Tuddenham E.G.D., McVey J.H. (2005) Normal haemostasis. In: Hoffbrand A.V., Catovsky D., Tuddenham E. (eds) *Postgraduate Haematology*, 5th edn. Blackwell Publishing, Oxford: pp. 783–801.
- Michelson A.D. (ed.) (2002) *Platelets*. Academic Press, San Diego, CA.
- Watson, S.P. and Harrison P. (2005) The vascular function of platelets. In: Hoffbrand A.V., Catovsky D., Tuddenham E. (eds) *Postgraduate Haematology*, 5th edn. Blackwell Publishing, Oxford: pp. 808–824.

CHAPTER 23

23

Bleeding disorders caused by vascular and platelet abnormalities

Vascular bleeding disorders, 278
Thrombocytopenia, 280
Disorders of platelet function, 286

Diagnosis of platelet disorders, 287
Platelet transfusions, 288
Bibliography, 289

Abnormal bleeding may result from:

- 1 Vascular disorders;
- 2 Thrombocytopenia;
- 3 Defective platelet function; or
- 4 Defective coagulation.

The pattern of bleeding is relatively predictable depending on the aetiology. Vascular and platelet disorders tend to be associated with bleeding from mucous membranes and into the skin whereas in coagulation disorders the bleeding is often into joints or soft tissue (Table 23.1).

The first three categories are discussed in this chapter and the disorders of blood coagulation follow in Chapter 24.

Vascular bleeding disorders

The vascular disorders are a heterogeneous group of conditions characterized by easy bruising and spontaneous bleeding from the small vessels. The underlying abnormality is either in the vessels themselves or in the perivascular connective tissues. Most cases of bleeding caused by vascular defects alone are not severe. Frequently, the bleeding is mainly in the skin causing petechiae, ecchymoses or both (Fig. 23.1). In some disorders there is also bleeding from mucous membranes. In these conditions the standard screening tests are normal. The bleeding time is normal and the other tests of haemostasis are also normal. Vascular defects may be inherited or acquired.

Inherited vascular disorders

Hereditary haemorrhagic telangiectasia

In this uncommon disorder, which is transmitted as an autosomal dominant trait. There are dilated microvascular swellings which appear during childhood and become more numerous in adult life. These telangiectasia develop in the skin, mucous membranes (Fig. 23.1a) and internal organs. Pulmonary and cerebral arteriovenous malformations are seen in a minority of cases. Recurrent epistaxes are frequent and recurrent gastrointestinal tract haemorrhage

Table 23.1 Clinical differences between diseases of platelets/vessel walls or coagulation factors.

Finding	Platelet/vessel walls	Coagulation
Mucosal bleeding	Common	Rare
Petechiae	Common	Rare
Deep haematomas	Rare	Characteristic
Bleeding from skin cuts	Persistent	Minimal
Sex of patient	Equal	>80% male



Fig. 23.1 (a) Hereditary haemorrhagic telangiectasia: the characteristic small vascular lesions are obvious on the lips and tongue. (b) Senile purpura. (c) Characteristic perifollicular petechiae in vitamin C deficiency (scurvy).

may cause chronic iron deficiency anaemia. Treatment is with embolization, laser treatment, oestrogens, tranexamic acid and iron supplementation.

Connective tissue disorders

In the Ehlers–Danlos syndrome there are hereditary collagen abnormalities with purpura resulting from defective platelet aggregation, hyperextensibility of joints and hyperelastic friable skin. Pseudoxanthoma elasticum is associated with arterial haemorrhage and thrombosis. Mild cases may present with superficial bruising and purpura following minor trauma.

Giant cavernous haemangioma

These congenital malformations occasionally cause chronic activation of coagulation leading to laboratory features of disseminated intravascular coagulation (DIC) and in some cases thrombocytopenia.

Acquired vascular defects

1 Simple easy bruising is a common benign disorder

which occurs in otherwise healthy women, especially those of child-bearing age.

2 Senile purpura caused by atrophy of the supporting tissues of cutaneous blood vessels is seen mainly on dorsal aspects of the forearms and hands (Fig. 23.1b).

3 Purpura associated with infections. Many bacterial, viral or rickettsial infections may cause purpura from vascular damage by the organism or as a result of immune complex formation (e.g. measles, dengue fever or meningococcal septicaemia).

4 The Henoch–Schönlein syndrome is usually seen in children and often follows an acute upper respiratory tract infection. It is an immunoglobulin A (IgA)-mediated vasculitis. The characteristic purpuric rash accompanied by localized oedema and itching is usually most prominent on the buttocks and extensor surfaces of the lower legs and elbows (Fig. 23.2). Painful joint swelling, haematuria and abdominal pain may also occur. It is usually a self-limiting condition but occasional patients develop renal failure.



Fig. 23.2 Henoch-Schönlein purpura: (a) unusually severe purpura on legs with bullous formation in a 6-year-old child; and (b) early urticarial lesions.

5 Scurvy. In vitamin C deficiency, defective collagen may cause perifollicular petechiae, bruising and mucosal haemorrhage (Fig. 23.1c).

6 Steroid purpura. The purpura, which is associated with long-term steroid therapy or Cushing's syndrome, is caused by defective vascular supportive tissue.

Tranexamic acid and aminocaproic acid are useful antifibrinolytic drugs that may reduce bleeding resulting from vascular disorders or thrombocy-

topenia but are contraindicated in the presence of haematuria because they might lead to clots obstructing the renal tract.

Thrombocytopenia

Abnormal bleeding associated with thrombocytopenia or abnormal platelet function is characterized by spontaneous skin purpura (Fig. 23.3) and mucosal haemorrhage and prolonged bleeding after




Fig. 23.3 (a) Typical purpura; and (b) massive subcutaneous haemorrhage in a patient with drug-induced thrombocytopenia.

Table 23.2 Causes of thrombocytopenia.**Failure of platelet production****Selective megakaryocyte depression**

- rare congenital defects
- drugs, chemicals, viral infections

Part of general bone marrow failure

- cytotoxic drugs
- radiotherapy
- aplastic anaemia
- leukaemia
- myelodysplastic syndromes
- myelofibrosis
- marrow infiltration (e.g. carcinoma, lymphoma)
- multiple myeloma
- megaloblastic anaemia
- HIV infection

Increased consumption of platelets**Immune**

- autoimmune
- idiopathic
 - associated with systemic lupus erythematosus, chronic lymphocytic leukaemia or lymphoma
- infections: HIV, other viruses, malaria
- drug-induced
- heparin
- post-transfusional purpura
- feto-maternal alloimmune thrombocytopenia

Disseminated intravascular coagulation**Thrombotic thrombocytopenic purpura****Abnormal distribution of platelets****Splenomegaly****Dilutional loss****Massive transfusion of stored blood to bleeding patients**

HIV, human immunodeficiency virus.

trauma (Table 23.1). The main causes of thrombocytopenia are listed in Tables 23.2 and 23.3.

Failure of platelet production

This is the most common cause of thrombocytopenia and is usually part of a generalized bone marrow failure (Table 23.2). Selective megakaryocyte depression may result from drug toxicity or viral infection. Rarely, it is congenital as a result of mutation of the c-MPL thrombopoietin receptor, in association with absent radii, or in May-Hegglin anomaly, with large inclusions in granulocytes, or

Table 23.3 Thrombocytopenia as a result of drugs or toxins.**Bone marrow suppression****Predictable (dose-related)**

- ionizing radiation, cytotoxic drugs, ethanol

Occasional

- chloramphenicol, co-trimoxazole, idoxuridine, penicillamine, organic arsenicals, benzene, etc.

Immune mechanisms (proven or probable)**Analgesics, anti-inflammatory drugs,**

- gold salts

Antimicrobials

- penicillins, sulphonamides, trimethoprim, para-aminosalicylate

Sedatives, anticonvulsants

- diazepam, sodium valproate, carbamazepine

Diuretics

- acetazolamide, chlorothiazides, frusemide

Antidiabetics

- chlorpropamide, tolbutamide

Others

- digitoxin, heparin, methyldopa, oxyprenolol, quinine, quinidine

Platelet aggregation**Ristocetin, heparin (p. 313)**

Wiskott-Aldrich syndrome (WAS) with eczema and immune deficiency. Diagnosis of these causes of thrombocytopenia is made from the clinical history, peripheral blood count, the blood film and bone marrow examination.

Increased destruction of platelets**Autoimmune (idiopathic) thrombocytopenic purpura**

Autoimmune (idiopathic) thrombocytopenic purpura (ITP) may be divided into chronic and acute forms.

Chronic idiopathic thrombocytopenic purpura

This is a relatively common disorder. The highest incidence has been considered to be in women aged 15–50 years although some reports suggest an increasing incidence with age. It is the most common cause of thrombocytopenia without anaemia or neutropenia. It is usually idiopathic but may




Fig. 23.4 The pathogenesis of thrombocytopenia in autoimmune thrombocytopenic purpura.

be seen in association with other diseases such as systemic lupus erythematosus (SLE), human immunodeficiency virus (HIV) infection, chronic lymphocytic leukaemia (CLL), Hodgkin's disease or autoimmune haemolytic anaemia (Table 23.2).

Pathogenesis

Platelet autoantibodies (usually IgG) result in the premature removal specific to platelets from the circulation by macrophages of the reticuloendothelial system, especially the spleen (Fig. 23.4). In many cases, the antibody is directed against antigen sites on the glycoprotein (GP) IIb-IIIa or Ib complex. The normal lifespan of a platelet is 7–10 days but in ITP this is reduced to a few hours. Total megakaryocyte mass and platelet turnover are increased in parallel to approximately five times normal.

Clinical features

The onset is often insidious with petechial haemorrhage, easy bruising and, in women, menorrhagia. Mucosal bleeding (e.g. epistaxes or gum bleeding) occurs in severe cases but fortunately intracranial haemorrhage is rare. The severity of bleeding in ITP is usually less than that seen in patients with comparable degrees of thrombocytopenia from bone

marrow failure; this is attributed to the circulation of predominantly young, functionally superior platelets in ITP. Chronic ITP tends to relapse and remit spontaneously so the course may be difficult to predict. Many asymptomatic cases are discovered by a routine blood count.

The spleen is not palpable unless there is an associated disease causing splenomegaly.

Diagnosis

- 1 The platelet count is usually $10\text{--}50 \times 10^9/\text{L}$. The haemoglobin concentration and white cell count are typically normal unless there is iron deficiency anaemia because of blood loss.
- 2 The blood film shows reduced numbers of platelets, those present often being large.
- 3 The bone marrow shows normal or increased numbers of megakaryocytes.
- 4 Sensitive tests are able to demonstrate specific antiglycoprotein GPIIb/IIIa or GPIb antibodies on the platelet surface or in the serum in most patients. Platelet-associated IgG assays are less specific.

Treatment

As this is a chronic disease the aim of treatment should be to maintain a platelet count above the level at which spontaneous bruising or bleeding




Fig. 23.5 Response to prednisolone in chronic immune thrombocytopenic purpura with subsequent relapse and response to splenectomy.

occurs with the minimum of intervention. In general, a platelet count above $50 \times 10^9/L$ does not require treatment.

1 Corticosteroids Eighty per cent of patients remit on high-dose corticosteroid therapy. Prednisolone 1 mg/kg/day is the usual initial therapy in adults and the dosage is gradually reduced after 10–14 days. In poor responders the dosage is reduced more slowly but splenectomy or alternative immunosuppression is considered.

2 Splenectomy (Fig. 23.5) This operation is recommended in patients who have symptoms and still have platelets $<30 \times 10^9/L$ after 3 months of steroid therapy or who require unacceptably high doses of steroids to maintain a platelet count above $30 \times 10^9/L$. Good results occur in most of the patients, but in patients with ITP refractory to steroids or immunoglobulin there may be little benefit. Splenunculi must be removed otherwise subsequent relapse of ITP can occur.

3 High-dose intravenous immunoglobulin therapy is able to produce a rapid rise in platelet count in the majority of patients. A regimen of 400 mg/kg/day for 5 days or 1 g/kg/day for 2 days is recommended. It is particularly useful in patients with life-threatening haemorrhage, in steroid-refractory ITP, during pregnancy or prior to surgery. The mechanism of action may be blockage of Fc recep-

tors on macrophages or modification of autoantibody production.

4 Immunosuppressive drugs (e.g. vincristine, cyclophosphamide, azathioprine, mycophenolate mofetil or ciclosporin alone or in combination) are usually reserved for those patients who do not respond sufficiently to steroids and splenectomy.

5 Monoclonal antibody Rituximab (anti-CD20) produces responses in approximately 50%, which are often durable.

6 Other treatments that may elicit a remission include danazol (an androgen which may virilize in women) and intravenous anti-D immunoglobulin. It is often necessary to combine two drugs (e.g. danazol and an immunosuppressive agent).

7 Platelet transfusions Platelet concentrates are beneficial in patients with acute life-threatening bleeding. Their benefit will only last a few hours.

8 Stem cell transplantation has cured some severe cases.

Acute idiopathic thrombocytopenic purpura

This is most common in children. In approximately 75% of patients the episode follows vaccination or an infection such as chickenpox or infectious mononucleosis. Most cases are caused by non-specific immune complex attachments. Spontaneous remissions are usual but in 5–10% of cases the disease becomes chronic (lasting >6 months). Fortunately, morbidity and mortality in acute ITP is very low.

The diagnosis is one of exclusion and there is debate as to the need for bone marrow aspiration. If the platelet count is over $30 \times 10^9/L$ no treatment is necessary unless the bleeding is severe. Those with counts below $20 \times 10^9/L$ may be treated with steroids and/or intravenous immunoglobulin, especially if there is significant bleeding.

Infections

It seems likely that the thrombocytopenia associated with many viral and protozoal infections is immune-mediated. In HIV infection, reduced platelet production is also involved (p. 328).

Post-transfusion purpura

Thrombocytopenia occurring approximately 10 days after a blood transfusion has been attributed to antibodies in the recipient developing against the human




Fig. 23.6 Usual type of platelet damage caused by drugs in which an antibody–drug–protein complex is deposited on the platelet surface. If complement is attached and the sequence goes to completion, the platelet may be lysed directly. Otherwise it is removed by reticuloendothelial cells because of opsonization with immunoglobulin and/or the C3 component of complement.

platelet antigen-1a (HPA-1a) (absent from the patient's own platelets) on transfused platelets. The reason why the patient's own platelets are then destroyed is unknown. Treatment is with intravenous immunoglobulin, plasma exchange or corticosteroids.

Drug-induced immune thrombocytopenia

An immunological mechanism has been demonstrated as the cause of many drug-induced thrombocytopenias (Fig. 23.6). Quinine (including that in tonic water), quinidine and heparin are particularly common causes (Table 23.3).

The platelet count is often less than $10 \times 10^9/L$, and the bone marrow shows normal or increased numbers of megakaryocytes. Drug-dependent antibodies against platelets may be demonstrated in the sera of some patients. The immediate treatment is to stop all suspected drugs but platelet concentrates should be given to patients with dangerous bleeding.

Thrombotic thrombocytopenic purpura and haemolytic uraemic syndrome

Thrombotic thrombocytopenic purpura (TTP) occurs in familial or acquired forms. There is deficiency of a ADAMTS13 metalloprotease which breaks down ultra large von Willebrand factor multimers (ULVWF) (Fig. 23.7). In the familial forms more than 50 ADAMTS13 mutations have been reported whereas in acquired forms it follows the development of an inhibitory IgG auto-antibody, the presence of which may be stimu-

lated by infection, autoimmune/connective tissue disease, certain drugs, stem cell transplantation or cardiac surgery. ULVWF multimeric strings secreted from Weibel–Palade bodies are anchored to the endothelial cells, and passing platelets adhere via their GPIb α receptors. Increasing platelet aggregation onto the ULVWF multimeric strings has the potential to form large, occlusive, platelet thrombi. These strings are capable of embolising to microvessels downstream contributing to organ ischaemia (Fig. 23.8). In the closely related haemolytic uraemic syndrome (HUS) ADAMTS13 levels are normal.

TTP has traditionally been described as pentad of thrombocytopenia, microangiopathic haemolytic anaemia, neurologic abnormalities, renal failure and fever. The microvascular thrombosis causes variable degrees of tissue ischaemia and infarction and is responsible for the microangiopathic haemolytic anaemia and thrombocytopenia. In current clinical practice, thrombocytopenia, schistocytosis, and an impressively elevated serum lactate dehydrogenase (LDH) value are sufficient to suggest the diagnosis. The serum LDH is derived both from ischaemic or necrotic tissue cells and lysed red cells. Coagulation tests are normal in contrast to the findings in DIC (Fig. 24.8). The serum LDH is raised. ADAMTS13 is absent or severely reduced in plasma. Treatment is with plasma exchange, using fresh frozen plasma (FFP) or cryosupernatant. This removes the large molecular weight VWF multimers and the antibody and provides ADAMTS13.




Fig. 23.7 Proposed pathogenesis of thrombotic thrombocytopenic purpura (TTP). Von Willebrand factor (VWF) consists of a series of VWF multimers each of molecular weight (MW) 250 kDa which are covalently linked. **(a)** Under physiological circumstances a metalloprotease ADAMTS13 cleaves high molecular weight multimers at a Tyr-842–Met-843 bond and the resulting VWF has an MW of 500–20 000 kDa. **(b)** In non-familial TTP, an antibody develops to the metalloprotease and so blocks cleavage of VWF multimers. **(c)** In congenital forms of TTP, the protease appears to be absent. In both cases, the resultant ultra large VWF multimers can bind platelets under high shear stress conditions and lead to platelet aggregation.

The platelet count and serum LDH are useful for monitoring the response to treatment. More recently, rituximab (anti-CD20) has been found effective in therapy, used in conjunction with plasma infusions or with plasma exchange. In refractory cases and chronic relapsing cases, high-dose corticosteroids, vincristine, intravenous immunoglobulin, rituximab and immunosuppressive therapy with azathioprine or cyclophosphamide have been used. In untreated cases mortality may approach 90%. Relapses are frequent.

HUS in children has many common features but organ damage is limited to the kidneys. There is also usually diarrhoea. Fits are frequent. Many cases are associated with *Escherichia coli* infection with vero-


toxin 0157 or with other organisms, especially *Shigella*. Supportive renal dialysis and control of hypertension and fits are the mainstays of treatment. Platelet transfusions are contraindicated in HUS and TTP.

Disseminated intravascular coagulation


Thrombocytopenia may result from an increased rate of platelet destruction through consumption of platelets because of their participation in DIC (p. 298).

Increased splenic pooling

The major factor responsible for thrombocytopenia in splenomegaly is platelet 'pooling' by the spleen. In splenomegaly, up to 90% of platelets may be



(a)



(b)

Fig. 23.8 Thrombotic thrombocytopenic purpura. (a) Platelet thrombus in a small cardiac vessel with minor endothelial and inflammatory reaction. (Courtesy of Dr J.E. McLaughlin) (b) Peripheral blood film showing red cell fragmentation.

sequestered in the spleen whereas normally this accounts for approximately one-third of the total platelet mass (Fig. 23.9). Platelet lifespan is normal and in the absence of additional haemostatic defects, the thrombocytopenia of splenomegaly is not usually associated with bleeding.

Massive transfusion syndrome

Platelets are unstable in blood stored at 4°C and the platelet count rapidly falls in blood stored for more than 24 h. Patients transfused with massive amounts of stored blood (more than 10 units over a 24-h period) frequently show abnormal clotting and thrombocytopenia. These should be corrected by the use of platelet transfusions and FFP.

Disorders of platelet function

Disorders of platelet function are suspected in patients who show skin and mucosal haemorrhage and in whom the bleeding time is prolonged despite a normal platelet count. These disorders may be hereditary or acquired.

Hereditary disorders

Rare inherited disorders may produce defects at each of the different phases of the platelet reactions leading to the formation of the haemostatic platelet plug.




Fig. 23.9 The platelet distribution between the circulation and spleen in normal individuals (left), and in patients with moderate or massive splenomegaly (right).

Thrombasthenia (Glanzmann's disease)

This autosomal recessive disorder leads to failure of primary platelet aggregation because of a deficiency of membrane GPIIb (gene on chromosome 17). It usually presents in the neonatal period and, characteristically, platelets fail to aggregate *in vitro* to any agonist except ristocetin.

Bernard-Soulier syndrome

In this disease the platelets are larger than normal and there is a deficiency of GPIb (chromosome 23). There is defective binding to VWF, defective adherence to exposed subendothelial connective tissues and platelets do not aggregate with ristocetin. There is a variable degree of thrombocytopenia.

Storage pool diseases

In the rare grey platelet syndrome, the platelets are larger than normal and there is a virtual absence of α granules with deficiency of their proteins. In the more common β -storage pool disease there is a deficiency of dense granules.

Platelet function is abnormal in von Willebrand disease because of an inherited defect in VWF (p. 295).

Acquired disorders

Antiplatelet drugs

Aspirin therapy is the most common cause of defective platelet function. It produces an abnormal bleeding time and, although purpura may not be obvious, the defect may contribute to the associated gastrointestinal haemorrhage. The cause of the aspirin defect is inhibition of cyclo-oxygenase with impaired thromboxane A_2 synthesis (see Fig. 25.8). There is consequent impairment of the release reaction and aggregation with adrenaline and adenosine diphosphate (ADP). After a single dose the defect lasts 7–10 days, i.e. the life of the platelet. Aspirin is contraindicated in patients with gastrointestinal or genitourinary bleeding, retinal bleeding, peptic ulcer, haemophilia or severe hypertension. Dipyridamole inhibits platelet aggregation by blocking reuptake of adenosine and is usually used as an adjunct to oral anticoagulants. Clopidogrel inhibits binding of ADP to its platelet receptor and is mainly used for prevention of thrombotic events

(e.g. after coronary stenting or angioplasty) in patients with a history of symptomatic atherosclerotic disease. Intravenous agents abciximab, eptifibatide and tirofiban are inhibitors of GPIIb/IIIa receptor sites and may be used in patients undergoing percutaneous coronary intervention with unstable angina and acute coronary syndromes. There is a risk of transient thrombocytopenia with these agents, especially with abciximab, and platelet transfusions may be needed.

Hyperglobulinaemia

Hyperglobulinaemia associated with multiple myeloma or Waldenström's disease may cause interference with platelet adherence, release and aggregation.

Myeloproliferative and myelodysplastic disorders

Intrinsic abnormalities of platelet function occur in many patients with essential thrombocythaemia and other myeloproliferative and myelodysplastic diseases and in paroxysmal nocturnal haemoglobinuria.

Uraemia

This is associated with various abnormalities of platelet function. Heparin, dextrans, alcohol and radiographic contrast agents may also cause defective function.

Diagnosis of platelet disorders

Patients with suspected platelet or blood vessel abnormalities should initially have a blood count and blood film examination (Fig. 23.10). Bone marrow examination is often needed in thrombocytopenic patients to determine whether or not there is a failure of platelet production. The marrow may also reveal one of the conditions associated with defective production (Table 23.2). In children and young adults with isolated thrombocytopenia, the marrow test is often not performed. In the elderly, the test is needed particularly to exclude myelodysplasia. In patients with thrombocytopenia, a negative drug history, normal or excessive numbers of marrow megakaryocytes and no other marrow abnormality or splenomegaly, ITP is the usual




Fig. 23.10 Laboratory tests for platelet disorders. NB. Some intrinsic platelet functional disorders are associated with thrombocytopenia (e.g. Bernard-Soulier syndrome). ADP, adenosine diphosphate; DIC, disseminated intravascular coagulation.




Fig. 23.11 Defective platelet aggregation in a patient on aspirin therapy. There is no secondary phase aggregation with adenosine diphosphate (ADP) and reduced responses to both adrenaline and collagen. Similar results are obtained in α -storage granule deficiency and cyclo-oxygenase deficiency.

diagnosis. Testing for platelet antibodies in serum or on the surface of platelets has not proved reliable in distinguishing ITP from other causes of thrombocytopenia. Screening tests for DIC are also useful, as are tests for an underlying disease (e.g. SLE or HIV infection).

When the blood count, including platelet count and blood film examination, are normal, a PFA-100 (platelet function analysis) or, much less frequently, a bleeding time is used to detect abnormal platelet function. In most patients with abnormal platelet function demonstrated by prolonged bleeding time, or the PFA-100 test, the defect is acquired and associated either with systemic disease (e.g. uraemia) or with aspirin therapy. The very rare hereditary

defects of platelet function require more elaborate *in vitro* tests to define the specific abnormality. These include platelet aggregation studies (Fig. 23.11) and nucleotide pool measurements. If von Willebrand disease is suspected, assay of VWF and coagulation factor VIII are required.

Platelet transfusions

Transfusion of platelet concentrates are indicated in the following circumstances:

1 Thrombocytopenia or abnormal platelet function when bleeding or before invasive procedures and there is no alternative therapy available (e.g. steroids or high-dose immunoglobulin). The platelet count

should be above $50 \times 10^9/\text{L}$ before, for example, liver biopsy or lumbar puncture.

2 Prophylactically in patients with platelet counts of less than $5\text{--}10 \times 10^9/\text{L}$. If there is infection, potential bleeding sites or coagulopathy, the count should be kept above $20 \times 10^9/\text{L}$.

The indications for transfusion of platelet concentrates are discussed further on p. 349.

Bibliography

- Balduini C.L., Cattaneo M., Fabris F. *et al.* (2003) Inherited thrombocytopenias: a proposed diagnostic algorithm from the Italian Gruppo di Studio delle Piastrine. *Haematologica* 88,582–92.
- British Committee for Standards in Haematology, Blood Transfusion Task Force (2003) Guidelines for the use of platelet transfusions. *Br J Haematol* 122,10–23.
- British Committee for Standards in Haematology, General Haematology Task Force (2003) Guidelines for the investigation and management of idiopathic thrombocytopenic purpura in adults, children and in pregnancy. *Br J Haematol* 120,574–96.
- Fakhouri F., Vernant J.P., Veyradier A. *et al.* (2005) Efficiency of curative and prophylactic treatment with rituximab in ADAMTS13-deficient thrombotic thrombocytopenic purpura: a study of 11 cases. *Blood* 106,1932–7.
- George J.N. (2006) Clinical practice. Thrombotic thrombocytopenic purpura. *N Engl J Med* 354,1927–35.
- Keeling D., Davidson S., Watson H. (2006) The management of heparin-induced thrombocytopenia. *Br J Haematol* 133,258–69.
- Levy G.G., Motto D.G., Ginsburg D. (2005) ADAMTS13 turns 3. *Blood* 106,11–7.
- Mannucci P.M., Canciani M.T., Forza I. *et al.* (2001) Changes in health and disease of the metalloprotease that cleaves von Willebrand factor. *Blood* 98,2730–5.
- Moake J.L. (2002) Thrombotic microangiopathies. *N Engl J Med* 347,589–600.
- Provan D. and Newland A. (2002) Fifty years of idiopathic thrombocytopenic purpura (ITP): management of refractory ITP in adults. *Br J Haematol* 118,933–44.
- Vesely S.K., George J.N., Lamule B. *et al.* (2003) ADAMTS13 activity in thrombotic thrombocytopenic purpura-hemolytic uremic syndrome: relation to presenting features and clinical outcomes in a prospective cohort of 142 patients. *Blood* 102,60–8.

CHAPTER 24

24

Coagulation disorders

Hereditary coagulation disorders, 290

Haemophilia A, 290

Acquired coagulation disorders, 296

Thromboelastography, 301

Bibliography, 302

Hereditary coagulation disorders

Hereditary deficiencies of each of the coagulation factors have been described. Haemophilia A (factor VIII deficiency), haemophilia B (Christmas disease, factor IX deficiency) and von Willebrand disease (VWD) are the most common; the others are rare.

Haemophilia A

Haemophilia A is the most common of the hereditary clotting factor deficiencies. The prevalence is of the order of 30–100 per million population. The inheritance is sex-linked (Fig. 24.1) but up to 33% of patients have no family history and result from spontaneous mutation.

Molecular genetics

The factor VIII gene is situated near the tip of the long arm of the X chromosome (Xq2.6 region). It is extremely large and consists of 26 exons. The factor VIII protein includes a triplicated region A₁A₂A₃ with 30% homology with each other, a duplicated homology region C₁C₂ and a heavy glycosylated B domain which is removed when factor VIII is activated by thrombin. The protein is synthesized in the liver and spleen.

The defect is an absence or low level of plasma factor VIII. Approximately half of the patients have missense or frameshift mutations or deletions in the




Fig. 24.1 A typical family tree in a family with haemophilia. Note the variable levels of factor VIII activity in carriers (*) because of random inactivation of X chromosome (Lyonization). The percentages show the degree of factor VIII activity as a percentage of normal.

factor VIII gene. In others a characteristic 'flip-tip' inversion is seen in which the factor VIII gene is broken by an inversion at the end of the X chromosome (Fig. 24.2). This mutation leads to a severe clinical form of haemophilia A.

Clinical features

Infants may suffer from profuse post-circumcision haemorrhage or develop joint and soft tissue bleeds




Fig. 24.2 The mechanism of the flip-tip inversion leading to disruption of the factor VIII gene. (Left) The orientation of the factor VIII gene is shown with the three copies of gene A in this region (one within intron 22 and two near the telomere). (Middle) During spermatogenesis at meiosis, the single X pairs with the Y chromosome in the homologous regions. The X chromosome is longer than

the Y and there is nothing to pair with most of the long arm of X. The chromosome undergoes homologous recombination between the A genes. (Right) The final result is that the factor VIII gene is disrupted. cen, centromeric end; tel, telomere; the arrows indicate the direction of transcription from the A gene.



Fig. 24.3 Haemophilia A: acute haemarthrosis of the left knee joint with swelling of the suprapatellar region. There is wasting of the quadriceps muscles, particularly on the right.

and excessive bruising when they start to be active. Recurrent painful haemarthroses and muscle haematomas dominate the clinical course of severely affected patients and if poorly treated may lead to progressive joint deformity and disability (Figs 24.3–24.6). Local pressure can cause entrap-

ment neuropathy or ischaemic necrosis. Prolonged bleeding occurs after dental extractions. Spontaneous haematuria and gastrointestinal haemorrhage, sometimes with obstruction resulting from intramucosal bleeding, can also occur. The clinical severity of the disease correlates with the extent of the factor VIII deficiency (Table 24.1). Operative and post-traumatic haemorrhage are life-threatening both in severely and mildly affected patients. Although not common, spontaneous intracerebral haemorrhage occurs more frequently than in the general population and is an important cause of death in patients with severe disease.

Haemophilic pseudotumours are large encapsulated haematomas with progressive cystic swelling from repeated haemorrhage. They may occur in fascial and muscle planes, large muscle groups and in the long bones, pelvis and cranium. These result from repeated subperiosteal haemorrhages with bone destruction and new bone formation.

As a result of human immunodeficiency virus (HIV) present in concentrates made from human plasma during the early 1980s, over 50% of haemophiliacs treated in the USA or Western Europe became infected with HIV. Acquired immune deficiency syndrome (AIDS) has been a common cause of death in severe haemophilia. Thrombocytopenia from HIV infection may exacerbate bleeding episodes.




Fig. 24.4 Haemophilia A showing severe disability. The left knee is swollen with posterior subluxation of the tibia on the femur. The ankles and feet show residual deformities of talipes equinus, with some cavus and associated toe clawing. There is generalized muscle wasting. The scar on the medial side of the left lower thigh is the site of a previously excised pseudotumour.

Many patients were infected with hepatitis C virus before testing of donors and blood products became possible. This is causing an increasing amount of morbidity including chronic hepatitis, cirrhosis and hepatoma. Hepatitis B transmission may also be a risk.

Laboratory findings (Table 24.2)

The following tests are abnormal:

- 1 Activated partial thromboplastin time (APTT).
- 2 Factor VIII clotting assay.

The bleeding time and prothrombin time (PT) tests are normal.

Carrier detection and antenatal diagnosis

Until recently, carrier detection and antenatal diagnosis were limited to measuring plasma levels of factor VIII and von Willebrand factor (VWF). Carriers are now better detected with DNA probes. A known specific mutation can be identified or restriction fragment length polymorphisms within or close to the factor VIII gene allows the mutant allele to be tracked. Chorionic biopsies at 8–10 weeks' gestation provide sufficient fetal DNA for analysis. Antenatal diagnosis is also possible following the demonstration of low levels of factor VIII in fetal blood obtained at 16–20 weeks' gestation from the umbilical vein by ultrasound-guided needle aspiration.



Fig. 24.5 Haemophilia A: massive haemorrhage in the area of the right buttock.



Fig. 24.6 Haemophilia A: radiographic appearances of the right elbow joint in a 25-year-old male. The joint space has been destroyed and there is bony ankylosis. Subchondral cystic areas are prominent.

Treatment

Most patients attend specialized haemophilia centres where there is a multidisciplinary team dedicated to their care. Bleeding episodes are treated with factor VIII replacement therapy and spontaneous bleeding is usually controlled if the patient's factor VIII level is raised to 30–50% of

normal. Guidelines exist for the plasma level to be achieved for different types of haemorrhage. For major surgery, serious post-traumatic bleeding or when haemorrhage is occurring at a dangerous site, the factor VIII level should be elevated to 100% and then maintained above 50% when acute bleeding has stopped, until healing has occurred. On average, factor VIII infusion produces a plasma increment of 2 U/dL per unit infused per kilogram body weight. Roughly, the dose to be infused (units) = (weight (kg) × increment needed (U/dL))/2.

Recombinant factor VIII (five different commercial preparations) and immunoaffinity-purified factor VIII preparations, which are heat or solvent-detergent treated, are now available for clinical use and virtually eliminate the risk of viral transmission.

DDAVP (desmopressin) provides an alternative means of increasing the plasma factor VIII level in milder haemophiliacs. Following the intravenous administration of this drug, there is a 2–4-fold rise maximum at 30–60 min in the patient's own factor VIII by release from endothelial cells and this rise is proportional to the resting level. DDAVP may also be taken subcutaneously or nasally—this has been used as immediate treatment for mild haemophilia after accidental trauma or haemorrhage. DDAVP has an antidiuretic action and should be avoided in the elderly; fluid restitution is advised after its use.

Local supportive measures used in treating haemarthroses and haematomas include resting the affected part and the prevention of further trauma.

Table 24.1 Correlation of coagulation factor activity and disease severity in haemophilia A or B.

Coagulation factor activity (percentage of normal)	Clinical manifestations
<2	Severe disease Frequent spontaneous bleeding into joints, muscles, internal organs from early life Joint deformity and crippling if not adequately treated
2–10	Moderate disease Bleeding after minor trauma Occasional spontaneous episodes
>10–30	Mild disease Minor bleeding after significant trauma, surgery

Table 24.2 Main clinical and laboratory findings in haemophilia A, factor IX deficiency (haemophilia B, Christmas disease) and von Willebrand disease.

	Haemophilia A	Factor IX deficiency	von Willebrand disease
Inheritance	Sex-linked	Sex-linked	Dominant (incomplete)
Main sites of haemorrhage	Muscle, joints, post-trauma or postoperative	Muscle, joints, post-trauma or postoperative	Mucous membranes, skin cuts, post-trauma or postoperative
Platelet count	Normal	Normal	Normal
Bleeding time	Normal	Normal	Prolonged
Prothrombin time	Normal	Normal	Normal
Partial thromboplastin time	Prolonged	Prolonged	Prolonged or normal
Factor VIII	Low	Normal	May be moderately reduced
Factor IX	Normal	Low	Normal
VWF	Normal	Normal	Low or abnormal function (Table 24.3)
Ristocetin-induced platelet aggregation	Normal	Normal	Impaired

VWF, von Willebrand factor.

Prophylactic treatment

The increased availability of factor VIII concentrates that may be stored in domestic refrigerators has dramatically altered haemophilia treatment. At the earliest suggestion of bleeding, the haemophilic child may be treated at home. This advance has reduced the occurrence of crippling haemarthroses and the need for inpatient care. Severely affected patients are now reaching adult life with little or no arthritis. After the first spontaneous joint bleed, most boys with severe haemophilia are started on prophylactic factor VIII three times a week, aiming to keep their factor VIII levels above 1%. This may require the placement of a vascular access device such as Port-a-Cath if venous access is difficult.

Haemophiliacs are advised to have regular conservative dental care. Haemophilic children and their parents often require extensive help with social and psychological matters. With modern treatment the lifestyle of a haemophilic child can be almost normal but certain activities such as body contact sports are to be avoided.

Gene therapy

Because it is only necessary to maintain factor levels >1% to prevent most of the mortality and morbidity of factor VIII or IX deficiency, there is great interest in gene-based therapy. Various viral vectors (retro-

viral, adeno-associated) as well as non-viral vectors are being explored. Phase 1 trials have been carried out for both haemophilia A and B.

Inhibitors

One of the most serious complications of haemophilia is the development of antibodies (inhibitors) to infused factor VIII which occurs in 5–10% of (mainly severe) patients, especially children. This renders the patient refractory to further replacement therapy so that tremendous doses have to be given to achieve a significant rise in plasma factor VIII activity. Immunosuppression and immune tolerance regimes have been used in an attempt to eradicate the antibody. Recombinant activated factor VII (VIIa) and activated prothrombin complex concentrates (FEIBA—factor VIII inhibitor bypassing activity) can be useful in the treatment of bleeding episodes.

Factor VIIa complexes with tissue factor exposed at the site of injury and produces local haemostasis. The process is independent of factor VIII or IX and is not affected by their inhibitors. Factor VIIa has a short half-life and therefore frequent doses may be needed. In the longer term, immunosuppression with cyclophosphamide, intravenous immunoglobulin and high-dose factor VIII has also been successful.

Factor IX deficiency

The inheritance and clinical features of factor IX deficiency (Christmas disease, haemophilia B) are identical to those of haemophilia A. Indeed, the two disorders can only be distinguished by specific coagulation factor assays. The incidence is one-fifth that of haemophilia A. Factor IX is coded by a gene close to the gene for factor VIII near the tip of the long arm of the X chromosome. Its synthesis, like that of prothrombin, factor VII, factor X and protein C, is vitamin K-dependent. Carrier detection and antenatal diagnosis is performed as for haemophilia A. The principles of replacement therapy are similar to those of haemophilia A. Bleeding episodes are treated with high-purity factor IX concentrates. Because of its longer biological half-life, infusions do not have to be given as frequently as factor VIII concentrates in haemophilia A. Recombinant factor IX is available. Higher doses are needed compared with plasma-derived factor IX.

Laboratory findings (Table 24.2)

The following tests are abnormal:

1 APTT

2 Factor IX clotting assay

As in haemophilia A, the bleeding time and PT tests are normal.

Von Willebrand disease

In this disorder there is either a reduced level or

abnormal function of VWF resulting from a point mutation or major deletion. VWF is produced in endothelial cells and megakaryocytes. It has two roles (Chapter 22). It promotes platelet adhesion to damaged endothelium and it is the carrier molecule for factor VIII, protecting it from premature destruction. The latter property explains the occasional reduced factor VIII levels found in VWD.

Chronic elevation of VWF is part of the acute phase response to injury, inflammation, neoplasia or pregnancy. VWF is synthesized as a large 300-kDa protein which then forms multimers up to 10^6 Da in weight. Three types of VWD have been described (Table 24.3). Type 2 is divided into four subtypes depending on the type of functional defect. Type 1 accounts for 75% of cases.

VWD is the most common inherited bleeding disorder. Usually, the inheritance is autosomal dominant with varying expression. The severity of the bleeding is variable. Typically, there is mucous membrane bleeding (e.g. epistaxes, menorrhagia), excessive blood loss from superficial cuts and abrasions, and operative and post-traumatic haemorrhage. The severity is variable in the different types. Haemarthroses and muscle haematomas are rare, except in type 3 disease.

Laboratory findings (Table 24.2)

1 The bleeding time can be prolonged. This test is usually replaced by the PFA-100 test (see p. 277).

2 Factor VIII levels are often low. If low, a factor VIII VWF binding assay is performed.

Table 24.3 Classification of von Willebrand disease.

Type 1	Quantitative partial deficiency
Type 2	Functional abnormality
Type 3	Complete deficiency

Secondary classification of type 2 VWD			
Subtype	Platelet-associated function	Factor VIII binding capacity	High MW VWF multimers
2A	Decreased	Normal	Absent
2B	Increased affinity for GPIb	Normal	Usually reduced/absent
2M	Decreased	Normal	Normal or ultra large
2N	Normal	Reduced	Normal

GPIb, glycoprotein Ib; MW, molecular weight; VWD, von Willebrand disease; VWF, von Willebrand factor.

- 3 The APTT may be prolonged.
- 4 VWF levels are usually low.
- 5 There is defective platelet aggregation by patient plasma in the presence of ristocetin (VWF: Rco). Aggregation to other agents (adenosine diphosphate (ADP), thrombin or adrenaline) is usually normal.
- 6 Collagen-binding function (VWF: CB) is usually reduced.
- 7 Multimer analysis is useful for diagnosing different subtypes (Table 24.3).
- 8 The platelet count is normal except for type 2B disease (where it is low).

Treatment

Options are as follows:

- 1 Local measures and antifibrinolytic agent (e.g. tranexamic acid for mild bleeding).
- 2 DDAVP infusion for those with type 1 VWD. This releases VWF from endothelial stage sites 30 min after intravenous infusion.
- 3 High-purity factor VWF concentrates for patients with very low VWF levels. Factor VIII concentrate may also be given for more rapid correction.

Hereditary disorders of other coagulation factors

All these disorders (deficiency of fibrinogen, prothrombin, factors V, VII, combined V and VIII, factors X, XI, XIII) are rare. In most the inheritance is autosomal recessive. Factor XI deficiency is seen mainly in Ashkenazi Jews and occurs in either sex. The bleeding risk is not related to severity of the deficiency, and is absent from muscles and joints. The severity is poorly related to the factor XI level in plasma. It usually causes excess bleeding only after trauma such as surgery, and is treated by factor XI concentrate or fresh frozen plasma. Factor XIII deficiency produces a severe bleeding tendency, characteristically with umbilical stump bleeding. Specific concentrates or recombinant preparation of factors VII, XI and XIII are now available.

Acquired coagulation disorders

The acquired coagulation disorders (Table 24.4) are more common than the inherited disorders. Unlike

Table 24.4 The acquired coagulation disorders.

<i>Deficiency of vitamin K-dependent factors</i>
Haemorrhagic disease of the newborn
Biliary obstruction
Malabsorption of vitamin K (e.g. tropical sprue, gluten-induced enteropathy)
Vitamin K-antagonist therapy (e.g. coumarins, indandiones)
Liver disease
Disseminated intravascular coagulation
<i>Inhibition of coagulation</i>
Specific inhibitors (e.g. antibodies against factor VIII)
Non-specific inhibitors (e.g. antibodies found in systemic lupus erythematosus, rheumatoid arthritis)
<i>Miscellaneous</i>
Diseases with M-protein production
L-Asparaginase
Therapy with heparin, defibrinating agents or thrombolytics
Massive transfusion syndrome

the inherited disorders, multiple clotting factor deficiencies are usual.

Vitamin K deficiency

Fat-soluble vitamin K is obtained from green vegetables and bacterial synthesis in the gut. Deficiency may present in the newborn (haemorrhagic disease of the newborn) or in later life.

Deficiency of vitamin K is caused by an inadequate diet, malabsorption or inhibition of vitamin K by drugs such as warfarin which act as vitamin K antagonists. Warfarin is associated with a decrease in the functional activity of factors II, VII, IX and X and proteins C and S, but immunological methods show normal levels of these factors. The non-functional proteins are called PIVKA (proteins formed in vitamin K absence). Conversion of PIVKA factors to their biologically active forms is a post-translational event involving carboxylation of glutamic acid residues in the N-terminal region where these factors show strong sequence homology (Fig. 24.7). Gamma-carboxylated glutamic acid binds calcium ions through which it forms




Fig. 24.7 The action of vitamin K in γ -carboxylation of glutamic acid in coagulation factors which are then able to bind Ca^{2+} and attach to the platelet phospholipid.

a complex with phospholipid. In the process of carboxylation, vitamin K is converted to vitamin K epoxide which is cycled back to the reduced form by reductases. Warfarin interferes with the reduction of vitamin K epoxide leading to a functional vitamin K deficiency.

Haemorrhagic disease of the newborn

Vitamin K-dependent factors are low at birth and fall further in breast-fed infants in the first few days of life. Liver cell immaturity, lack of gut bacterial synthesis of the vitamin and low quantities in breast milk may all contribute to a deficiency which may cause haemorrhage, usually on the second to fourth day of life, but occasionally during the first 2 months.

Diagnosis

The PT and APTT are both abnormal. The platelet count and fibrinogen are normal with absent fibrin degradation products.

Treatment

1 Prophylaxis. For many years vitamin K has been given to all newborn babies as a single intramuscular injection of 1 mg. This remains the most appropriate and safest treatment. Following epidemiological evidence suggesting a possible link between intramuscular vitamin K and an increased risk of childhood tumours (which has not been substantiated), some centres recommended an oral regimen but this is less effective in prevention and repeat doses are needed up to day 28.

2 In bleeding infants: vitamin K 1 mg intramuscularly is given every 6 h with, initially, fresh frozen plasma if haemorrhage is severe.

Vitamin K deficiency in children or adults

Deficiency resulting from obstructive jaundice, pancreatic or small bowel disease occasionally causes a bleeding diathesis in children or adults.

Diagnosis

Both PT and APTT are prolonged. There are low plasma levels of factors II, VII, IX and X.

Treatment

1 Prophylaxis: vitamin K 5 mg/day orally.
2 Active bleeding or prior to liver biopsy: vitamin K 10 mg slowly intravenously. Some correction of PT is usual within 6 h. The dose should be repeated on the next 2 days after which optimal correction is usual.

Liver disease

Multiple haemostatic abnormalities contribute to a bleeding tendency and may exacerbate haemorrhage from oesophageal varices.

1 Biliary obstruction results in impaired absorption of vitamin K and therefore decreased synthesis of factors II, VII, IX and X by liver parenchymal cells.

2 With severe hepatocellular disease, in addition to a deficiency of these factors, there are often reduced

levels of factor V and fibrinogen and increased amounts of plasminogen activator.

3 Functional abnormality of fibrinogen (dysfibrinogenemia) is found in many patients.

4 Decreased thrombopoietin production from the liver contributes to thrombocytopenia.

5 Hypersplenism associated with portal hypertension frequently results in thrombocytopenia.

6 Disseminated intravascular coagulation (DIC; see below) may be related to release of thromboplastins from damaged liver cells and reduced concentrations of antithrombin, protein C and α_2 -antiplasmin. In addition, there is impaired removal of activated clotting factors and increased fibrinolytic activity.

Disseminated intravascular coagulation

Widespread inappropriate intravascular deposition of fibrin with consumption of coagulation factors and platelets occurs as a consequence of many disorders which release procoagulant material into the circulation or cause widespread endothelial damage or platelet aggregation (Table 24.5). It may be associated with a fulminant haemorrhagic or thrombotic syndrome with organ dysfunction or run a less severe and more chronic course. The main clinical presentation is with bleeding but 5–10% of patients manifest microthrombotic lesions (e.g. with gangrene of limbs).

Pathogenesis (Fig. 24.8)

The key event underlying DIC is increased activity of tissue factor. This can come from its release into the circulation from damaged tissues present on tumour cells or from up-regulation of tissue factor on circulating monocytes or endothelial cells

Table 24.5 Causes of disseminated intravascular coagulation.

Infections

Gram-negative and meningococcal septicaemia

Clostridium welchii septicaemia

Severe Falciparum malaria

Viral infection—varicella, HIV, hepatitis, cytomegalovirus

Malignancy

Widespread mucin-secreting adenocarcinoma

Acute promyelocytic leukaemia

Obstetric complications

Amniotic fluid embolism

Premature separation of placenta

Eclampsia; retained placenta

Septic abortion

Hypersensitivity reactions

Anaphylaxis

Incompatible blood transfusion

Widespread tissue damage

Following surgery or trauma

After severe burns

Vascular abnormalities

Kasabach–Merritt syndrome

Leaking prosthetic valves

Cardiac bypass surgery

Vascular aneurysms

Miscellaneous

Liver failure

Pancreatitis

Snake and invertebrate venoms

Hypothermia

Heat stroke

Acute hypoxia

Massive blood loss




Fig. 24.8 The pathogenesis of disseminated intravascular coagulation and the changes in clotting factors, platelets and fibrin degradation products (FDPs) that occur in this syndrome.

in response to pro-inflammatory cytokines (e.g. interleukin-1, tumour necrosis factor, endotoxin).

1 DIC may be triggered by the entry of procoagulant material into the circulation in the following situations: severe trauma, amniotic fluid embolism, premature separation of the placenta, widespread mucin-secreting adenocarcinomas, acute promyelocytic leukaemia (AML type M₃), liver disease, severe falciparum malaria, haemolytic transfusion reaction and some snake bites.

2 DIC may also be initiated by widespread endothelial damage and collagen exposure (e.g. endotoxaemia, Gram-negative and meningococcal septicaemia, septic abortion), certain virus infections and severe burns or hypothermia.

In addition to its role in the deposition of fibrin in the microcirculation, intravascular thrombin formation produces large amounts of circulating fibrin monomers which form complexes with fibrinogen. Intense fibrinolysis is stimulated by thrombi on vascular walls and the release of split products interferes with fibrin polymerization, thus contributing to the coagulation defect. The combined action of thrombin and plasmin normally causes depletion of fibrinogen and all coagulation factors. Intravascular thrombin also causes widespread platelet aggregation and deposition in the vessels. The bleeding problems which may be a feature of DIC are compounded by thrombocytopenia caused by consumption of platelets.

Clinical features

These are dominated by bleeding, particularly from venepuncture sites or recent wounds (Fig. 24.9a). There may be generalized bleeding in the gastrointestinal tract, the oropharynx, into the lungs, urogenital tract and in obstetric cases vaginal bleeding may be particularly severe. Less frequently, microthrombi may cause skin lesions, renal failure, gangrene of the fingers or toes (Fig. 24.9b) or cerebral ischaemia.

Some patients may develop subacute or chronic DIC, especially with mucin-secreting adenocarcinoma. Compensation by the liver may render some of the coagulation tests normal.

Laboratory findings (Table 24.6)

In many acute syndromes the blood may fail to clot because of gross fibrinogen deficiency.



(a)



(b)

Fig. 24.9 Clinical features of disseminated intravascular coagulation: (a) indurated and confluent purpura of the arm; (b) peripheral gangrene with swelling and discoloration of the skin of the feet in fulminant disease.

Tests of haemostasis

- 1 The platelet count is low.
- 2 Fibrinogen concentration low.
- 3 The thrombin time is prolonged.
- 4 High levels of fibrin degradation products such as D-dimers are found in serum and urine.
- 5 The PT and APTT are prolonged in the acute syndromes.

Blood film examination

In many patients there is a haemolytic anaemia ('microangiopathic') and the red cells show prominent fragmentation because of damage caused when passing through fibrin strands in small vessels (p. 69).

Table 24.6 Haemostasis tests: typical results in acquired bleeding disorders.

	Platelet count	Prothrombin time	Activated partial thromboplastin time	Thrombin time
Liver disease	Low	Prolonged	Prolonged	Normal (rarely prolonged)
DIC	Low	Prolonged	Prolonged	Grossly prolonged
Massive transfusion	Low	Prolonged	Prolonged	Normal
Oral anticoagulants	Normal	Grossly prolonged	Prolonged	Normal
Heparin	Normal (rarely low)	Mildly prolonged	Prolonged	Prolonged
Circulating anticoagulant	Normal	Normal or prolonged	Prolonged	Normal

DIC, Disseminated intravascular coagulation.

Table 24.7 Indications for the use of fresh frozen plasma (National Institutes of Health Consensus Guidelines).

Coagulation factor deficiency (where specific or combined factor concentrate is not available)
Reversal of warfarin effect
Multiple coagulation defects (e.g. in patients with liver disease, DIC)
Massive blood transfusion with coagulopathy and clinical bleeding
Thrombotic thrombocytopenic purpura
Deficiencies of antithrombin*, protein C* or protein S
Some patients with immunodeficiency syndromes

DIC, disseminated intravascular coagulation.

* Antithrombin and protein C concentrates now available.

Treatment

Treatment of the underlying cause is most important. The management of patients who are bleeding differs from that of patients with thrombotic problems.

Bleeding

Supportive therapy with fresh frozen plasma (Table 24.7) and platelet concentrates is indicated in patients with dangerous or extensive bleeding. Cryoprecipitate provides a more concentrated source of fibrinogen and red cell transfusions may be required.

Thrombosis

The use of heparin or antiplatelet drugs to inhibit the coagulation process is considered in those with

thrombotic problems such as dermal ischaemia. Fibrinolytic inhibitors should not be considered because failure to lyse thrombi in organs such as the kidney may have adverse effects. Antithrombin concentrates or recombinant human protein C may be used to inhibit DIC in severe cases with sepsis (e.g. meningococcal septicaemia). There is reduced activated protein C (APC) in severe sepsis and recombinant human APC has been found to reduce mortality in this setting.

Coagulation deficiency caused by antibodies

Circulating antibodies to coagulation factors are occasionally seen with an incidence of approximately 1 per million per year. Alloantibodies to factor VIII occur in 5–10% of haemophiliacs. Factor VIII autoantibodies may also result in a bleeding syndrome. These immunoglobulin G (IgG) antibodies occur rarely post-partum, in certain immunological disorders (e.g. rheumatoid arthritis), in cancer and in old age. Treatment usually consists of a combination of immunosuppression and treatment with factor replacement, usually as human factor VIII, recombinant VIIa or activated prothrombin complex concentrate (FEIBA).

Another protein known as the lupus anticoagulant interferes with lipoprotein-dependent stages of coagulation and is usually detected by prolongation of the APTT test (Table 24.6). This inhibitor is detected in 10% of patients with systemic lupus erythematosus (SLE) and in patients with other autoimmune diseases who frequently have antibodies

to other lipid-containing antigens (e.g. cardiolipin). The antibody is not associated with a bleeding tendency but there is an increased risk of arterial or venous thrombosis and, as with other causes of thrombophilia, an association with recurrent miscarriage (Chapter 25).

Massive transfusion syndrome

Many factors may contribute to a bleeding disorder following massive transfusion. Blood loss results in reduced levels of platelets, coagulation factors and inhibitors. Further dilution of these factors occurs during replacement with red cells.

Management

Platelet concentrates are given to maintain a platelet count $>50 \times 10^9/L$ or $80\text{--}100 \times 10^9/L$ in cerebral injury or polytrauma. The PT and APTT should be

kept to less than 1.5 times normal with fresh frozen plasma given initially at 15 mL/kg. Cryoprecipitate is given to keep fibrinogen at least 1 g/L. Recombinant VIIa is increasingly used in patients with massive blood loss after trauma or surgery to reduce haemorrhage.

The results of haemostasis screening tests in acquired bleeding disorders are shown in Table 24.6 and a summary of the indications for use of fresh frozen plasma in Table 24.7.

Thromboelastography: near-patient testing

Thromboelastography (TEG) is a technique for a global assessment of haemostatic function of a single blood sample in which the reaction of platelets with the protein coagulation cascade is observed from the time of the initial platelet fibrin interaction





Fig. 24.10 Thromboelastography (TEG): normal trace and appearances in different pathological states. α -angle, speed of solid clot formation; A_{60} , measure of clot lysis or retraction at 60 min; k, clot formation time; MA, absolute strength of fibrin clot; r, rate of initial fibrin formation. (From S.V. Mallett and D.J.A. Cox (1992). Thromboelastography. *Br J Anaesth* 69,307–13 with permission)



through platelet aggregation, clot strengthening and fibrin cross-linkage to eventual clot lysis. It is suited as a monitor of haemostasis in surgery (e.g. of the liver or heart) associated with haemostatic defects. Freshly drawn blood is placed in a cuvette which is oscillated, the motion being transferred to a pin which writes on heat-sensitive paper. As fibrin strands form, the fibrin clot affects movement of the pin. The normal trace shows the rate of initial fibrin formation, the time to formation of a clot (coagulation time), strength of the fibrin clot, clot lysis index or retraction. Typical patterns showing results in fibrinolysis, hypercoagulability, haemophilia and thrombocytopenia are shown in Fig. 24.10.

The platelet function analyser (PFA-100) may also be useful in testing platelet function before and during surgery (p. 277).

Bibliography

- Bernard G.R., Vincent J.L., Laterre P.F. *et al.* (2001) Efficacy and safety of recombinant human activated protein C for severe sepsis. *N Engl J Med* 344,699–709.
- Blumberg N., Francis C., Heal J. (eds) (2006) Clinical uses of plasma and plasma fractions. *Best Pract Res Clin Haematol* 19,1–258.
- Deldado J., Jimenez-Yuste V., Hernandez-Navarro F. *et al.* (2003) Acquired haemophilia: review and meta-analysis focused on therapy and prognostic factors. *Br J Haematol* 121,21–35.
- Erber W.N. and Perry D.J. (2006) Plasma and plasma products in the treatment of massive haemorrhage. *Best Pract Res Clin Haematol* 19,97–112.
- Favaloro E.J. (2001) Utility of the PFA-100® for assessing bleeding disorders and monitoring therapy: a review of analytical variables, benefits and limitations. *Haemophilia* 7,170–9.
- Haemophilia A database: <http://europium.csc.mrc.ac.uk>
- Haemophilia database: <http://www.unds.ac.uk/mdgen>
- Hay C.R.M., Brown S., Collins P.W. *et al.* (2006) The diagnosis and management of factor VIII and IX inhibitors: a guideline from the United Kingdom Haemophilia Centre Doctors Organisation. *Br J Haematol* 133,591–605.
- Josephson C.D. and Abshire T.C. (2006) Clinical uses of plasma and plasma fractions: plasma-derived products for hemophilias A and B, and for von Willebrand disease. *Best Pract Res Clin Haematol* 19,35–49.
- Laffan M., Brown S.A., Collins P.W. *et al.* (2004) The diagnosis of von Willebrand disease: a guideline from the UK Haemophilia Centre Doctors' Organization. *Haemophilia* 10,199–217.
- Lee C.A., Berntorp E.E., Hoots W.K. (eds) (2005) *Textbook of Haemophilia*. Blackwell Publishing, Oxford.
- Levi M. (2004) Current understanding of disseminated intravascular coagulation. *Br J Haematol* 124,567–76.
- Mannucci P.M. (2003) Hemophilia: treatment options in the twenty-first century. *J Thromb Haemost* 1,1349–55.
- Mannucci P.M. (2004) Treatment of von Willebrand's Disease. *N Engl J Med* 351,683–94.
- Mittal S. and Watson H.G. (2006) A critical appraisal of the use of recombinant factor VIIa in acquired bleeding conditions. *Br J Haematol* 133,355–63.
- O'Shaughnessy D.F., Atterbury C., Bolton Maggs P. *et al.* for the British Committee for Standards in Haematology, Blood Transfusion Task Force. (2004) Guidelines for the use of fresh-frozen plasma, cryoprecipitate and cryosupernatant. *Br J Haematol* 126,11–28.
- Pasi K.J., Collins P.W., Keeling D.M. *et al.* (2004) Management of von Willebrand disease: a guideline from the UK Haemophilia Centre Doctors' Organization. *Haemophilia* 10,218–31.
- Peyvandi F. and Mannucci P.M. (2005) Congenital bleeding: autosomal recessive disorders. In: Hoffbrand A.V., Tuddenham E., Catovsky D. (eds) *Postgraduate Haematology*, 5th edn. Blackwell Publishing, Oxford: pp. 842–58.
- Plug I., Mausev-Bunshoten P., Bröcker-Vriends A.H.J.T. *et al.* (2006) Bleeding in causes of haemophilia. *Blood* 108,52–6.
- Roberts H.R., Monroe D.M., White G.C. (2004) The use of recombinant factor VIIa in the treatment of bleeding disorders. *Blood* 104,3858–64.
- Salooja N. and Perry D.J. (2002) Thromboelastography. *Blood Coagul Fibrinolysis* 12,327–37.
- United Kingdom Haemophilia Centre Doctors' Organization. (2003) Guidelines on the selection and use of therapeutic products to treat haemophilia and other hereditary bleeding disorders. *Haemophilia* 9,1–23.

CHAPTER 25

25

Thrombosis and antithrombotic therapy

Arterial thrombosis, 303
Venous thrombosis, 304
Investigation of thrombophilia, 310
Diagnosis of venous thrombosis, 310
Anticoagulant drugs, 311
Heparin, 311

Oral anticoagulants, 313
Graduated compression stockings, 316
Inferior vena cava filter, 316
Fibrinolytic agents, 317
Antiplatelet drugs, 317
Bibliography, 319

Thrombi are solid masses or plugs formed in the circulation from blood constituents. Platelets and fibrin form the basic structure. Their clinical significance results from ischaemia from local vascular obstruction or distant embolization. Thrombi are involved in the pathogenesis of myocardial infarction, cerebrovascular disease, peripheral arterial disease and deep vein occlusion.

Thrombosis, both arterial and venous, is more common as age increases and is frequently associated with risk factors (e.g. surgery or pregnancy). The term thrombophilia is used to describe the inherited or acquired disorders of the haemostatic mechanism that predispose to thrombosis.

Arterial thrombosis

Pathogenesis

Atherosclerosis of the arterial wall, plaque rupture and endothelial injury expose blood to subendothelial collagen and tissue factor. This initiates the formation of a platelet nidus on which platelets adhere and aggregate.

Platelet deposition and thrombus formation are important in the pathogenesis of atherosclerosis. Platelet-derived growth factor (PDGF) stimulates the migration and proliferation of smooth muscle

cells and fibroblasts in the arterial intima. Regrowth of endothelium and repair at the site of arterial damage and incorporated thrombus result in thickening of the vessel wall.

As well as blocking arteries locally, emboli of platelets and fibrin may break away from the primary thrombus to occlude distal arteries. Examples are carotid artery thrombi leading to cerebral thrombosis and transient ischaemic attacks (TIAs) and heart valve and chamber thrombi leading to systemic emboli and infarcts (Fig. 25.1).

Clinical risk factors

The risk factors for arterial thrombosis are related to the development of atherosclerosis and are listed in Table 25.1. The identification of patients at risk is largely based on clinical assessment. A number of epidemiological studies have resulted in the construction of coronary artery thrombosis risk profiles based on sex, age, elevated blood pressure, high levels of serum cholesterol, glucose intolerance, cigarette smoking and electrocardiogram (ECG) abnormalities. These profiles have allowed pre-symptomatic assessment of young and apparently fit subjects and are valuable in counselling a change in lifestyle or for recommending medical therapy in individuals at risk. The Northwick Park Heart Study showed




Fig. 25.1 Arteriogram showing saddle embolus at the aortic bifurcation (dotted arrow) and embolus in the left common iliac artery (solid arrow).

Table 25.1 Risk factors in arterial thrombosis (atherosclerosis).

Positive family history
Male sex
Hyperlipidaemia
Hypertension
Diabetes mellitus
Gout
Polycythaemia
Hyperhomocysteinaemia
Cigarette smoking
ECG abnormalities
Elevated factor VII
Elevated fibrinogen
Lupus anticoagulant
Collagen vascular diseases
Behçet's disease

ECG, electrocardiogram.

that elevated plasma levels of factor VII and fibrinogen are the strongest independent predictors of coronary events. Hyperhomocysteinaemia has been recognized more recently as a risk factor for peripheral and coronary arterial disease and stroke.

Venous thrombosis

Pathogenesis and risk factors

Virchow's triad suggests that there are three components that are important in thrombus formation:

Table 25.2 Risk factors for venous thrombosis.

Related to coagulation abnormality

Hereditary haemostatic disorders

Factor V Leiden
Prothrombin G20210A variant
Protein C deficiency
Antithrombin deficiency
Protein S deficiency
Abnormal fibrinogen
Abnormal plasminogen

Hereditary or acquired haemostatic disorders

Raised plasma levels of factor VII, VIII, IX or XI
Raised plasma levels of fibrinogen
Raised plasma levels of homocysteine
Glucosylceramide deficiency
Coagulation factor IX concentrates
Lupus anticoagulant
Oestrogen therapy (oral contraceptive and HRT)
Heparin-induced thrombocytopenia
Pregnancy and puerperium
Surgery, especially abdominal and hip
Major trauma
Malignancy
Myocardial infarct
Thrombocythaemia

Related to stasis

Cardiac failure
Stroke
Prolonged immobility
Pelvic obstruction
Nephrotic syndrome
Dehydration
Hyperviscosity, polycythaemia
Varicose veins

Related to unknown factors

Age
Obesity
Sepsis
Paroxysmal nocturnal haemoglobinuria
Behçet's disease

HRT, hormone replacement therapy.


- 1 Slowing down of blood flow;
- 2 Hypercoagulability of the blood; and
- 3 Vessel wall damage.

For venous thrombosis, increased systemic coagulability and stasis are most important, vessel wall damage being less important than in arterial thrombosis, although it may be important in patients with sepsis and indwelling catheters. Stasis allows the completion of blood coagulation at the site of initiation of the thrombus (e.g. behind the valve pockets of the leg veins in immobile patients).


Table 25.2 lists a number of recognized risk factors.

Hereditary disorders of haemostasis


The prevalence of inherited disorders associated with increased risk of thrombosis is at least as high as that of hereditary bleeding disorders. A hereditary 'thrombophilia' should be particularly suspected in young patients who suffer from spontaneous thrombosis, recurrent deep vein thromboses (Fig. 25.2) or



(a)



(b)



(c)

Fig. 25.2 Diagnostic imaging of deep vein thrombosis (DVT) and pulmonary embolus (PE). (a) Colour power Doppler ultrasound of the right femoral vessels with compression shows normal flow in the femoral artery but absent flow in the vein because of thrombus. A normal vein would collapse with compression of the probe. (Courtesy of Dr Tony Young) (b) Femoral venogram demonstrating extensive thrombus within the right external iliac vein. (Courtesy of Drs I.S. Francis and A.F. Watkinson) (c) Computed tomography (CT) pulmonary angiography: a coronal image shows bilateral filling defects (green crosses) in the central pulmonary arteries indicating pulmonary emboli. (Courtesy of Dr Tony Young)

an unusual site of thrombosis (e.g. axillary, splanchnic veins, sagittal sinus). Several abnormalities are now well characterized (Table 25.2).

Factor V Leiden gene mutation (activated protein C resistance)

This is the most common inherited cause of an increased risk of venous thrombosis. It occurs in approximately 4% of Caucasian factor V alleles. It was first recognized because of a failure to see prolongation in the activated partial thromboplastin time (APTT) test when activated protein C was added to the plasma of certain patients. Protein C breaks down activated factor V so activated protein C should slow the clotting reaction and prolong the APTT. In 1994, the underlying reason for this phenomenon was recognized to be a genetic polymorphism in the factor V gene (replacement of arginine at position 506 with glutamine—Arg506Gln) which makes factor V less susceptible to cleavage by activated protein C (Fig. 25.3). This is called the factor V Leiden mutation. The frequency of factor V Leiden in the general population in Western countries means that it cannot be regarded as a rare mutation but as a genetic polymorphism that is maintained in the population (Fig. 25.4). Presumably, individuals with this allele have been 'selected', probably because of reduced bleeding tendency. It does not increase the risk of arterial thrombosis.




Fig. 25.3 The genetic basis of factor V Leiden.

(a) Activated protein C inactivates factor Va by proteolytic cleavage at three sites in the Va heavy chain. (b) In the factor V Leiden mutation the Arg506Gln polymorphism leads to glutamine at position 506 with less efficient inactivation of factor V and increased risk of thrombosis.

Patients who are heterozygous for factor V Leiden are at an approximately 5–8-fold increased risk of thrombosis compared to the general population. Individuals who are homozygous have a 30–140-fold risk. Following venous thrombosis they have a higher risk of re-thrombosis compared to individuals with deep vein thrombosis (DVT) but normal factor V.

The incidence of factor V Leiden in patients with venous thrombosis is approximately 20–40%. Polymerase chain reaction (PCR) screening for the mutation is relatively simple and the test is widely performed. The absolute risk of thrombosis will depend on many other factors and it is difficult to advise individual patients of their risk. At present it is not recommended to start anticoagulation therapy in individuals with the Leiden mutation, even if homozygous, with no history of thrombosis. A small minority of patients with activated protein C resistance do not have factor V Leiden and have other mutations of factor V.

Antithrombin deficiency

Inheritance is autosomal dominant. There are recurrent venous thromboses usually starting in early adult life. Arterial thrombi occur occasionally. Antithrombin concentrates are available and are used to prevent thrombosis during surgery or childbirth. Many molecular variants of antithrombin have been categorized and are associated with varying degrees of risk of thrombosis.

Protein C deficiency

Inheritance is autosomal dominant with variable penetrance. Protein C levels in heterozygotes are approximately 50% of normal. Characteristically, many patients develop skin necrosis as a result of dermal vessel occlusion when treated with warfarin, thought to be caused by reduction of protein C levels even further in the first day or two of warfarin therapy before reduction in the levels of the vitamin K-dependent clotting factors, especially factor VII. Rarely, infants may be born with homozygous deficiency and characteristically present with severe disseminated intravascular coagulation (DIC) or purpura fulminans in infancy. Protein C concentrates are available.




Fig. 25.4 The incidence of carriers of factor V Leiden in different countries.

Protein S deficiency

Protein S deficiency has been found in a number of families with a thrombotic tendency. It is a cofactor for protein C and the clinical features are similar to protein C deficiency, including a tendency to skin necrosis with warfarin therapy. The inheritance is autosomal dominant.

Prothrombin allele G20210A

Prothrombin allele G20210A is a variant (prevalence 2–3% in the population) that leads to increased plasma prothrombin levels and increases thrombotic risk by at least twofold. It is probable that the cause of venous thrombosis with this mutation and with high levels of factors VIII, IX and XI is that sustained generation of thrombin results in prolonged down-regulation of fibrinolysis through activation of thrombin-activated fibrinolysis inhibitor (p. 274).

Hyperhomocyst(e)inaemia

Higher levels of plasma homocysteine may be genetic or acquired and are associated with increased risk

for both venous and arterial thrombosis. However, recent large trials show no evidence that lowering the levels reduces these risks.

Homocysteine is derived from dietary methionine and is removed by either remethylation to methionine or conversion to cysteine via a *trans*-sulphuration pathway (Fig. 25.5). Classic homocystinuria is a rare autosomal recessive disorder caused by deficiency of cystathione β -synthase, the enzyme responsible for *trans*-sulphuration. Vascular disease and thrombosis are major features of the disease. Heterozygous cystathione β -synthase deficiency is present in approximately 0.5% of the population and leads to a moderate increase in homocysteine. Methylene tetrahydrofolate reductase (MTHFR) is involved in the remethylation pathway and a common thermolabile variant of the enzyme may be responsible for mild homocysteinaemia (above 15 $\mu\text{mol/L}$) although this may only be seen in the presence of folate deficiency (p. 46). Acquired risk factors for hyperhomocysteinaemia include deficiencies of folate, vitamin B₁₂ or vitamin B₆, drugs (e.g. ciclosporin), renal damage




Fig. 25.5 The metabolism of homocysteine. Homocysteine is derived from dietary methionine and is metabolized either by the *trans*-sulphuration or remethylation pathways. *Trans*-sulphuration proceeds using the cystathionine β -synthase (CBS) enzyme with vitamin B_6 as a cofactor. Remethylation involves the action of methionine synthase (MS) on 5-methyl-THF with vitamin B_{12} as a cofactor. In addition, methylene-tetrahydrofolate reductase (MTHFR) is involved in this cycle.

and smoking. The levels also increase with age and are higher in men and post-menopausal females.

Defects of fibrinogen

Defects of fibrinogen are usually clinically silent or cause excess bleeding. Thrombosis is a rare association.

Hereditary or acquired disorders of haemostasis

High factor VIII or fibrinogen levels are also associated arterial thrombosis.

The combination of multiple risk factors is associated with increased risk of thrombosis. If these are persistent they may represent a reason for extended anticoagulation.

Acquired risk factors

These may cause thrombosis in patients without another identifiable abnormality but are most likely to do so if an inherited predisposing abnormality (e.g. factor V Leiden) is also present.

Postoperative venous thrombosis

This is more likely to occur in the elderly, obese, those with a previous or family history of venous

thrombosis, and in those in whom major abdominal or hip operations are performed.

Venous stasis and immobility

These factors are probably responsible for the high incidence of postoperative venous thrombosis and for venous thrombosis associated with congestive cardiac failure, myocardial infarction and varicose veins. In atrial fibrillation, thrombin generation from accumulation of activated clotting factors leads to a high risk of systemic embolization. The use of muscle relaxants during anaesthesia may also contribute to venous stasis. Venous thrombosis also has a higher frequency after prolonged aeroplane journeys.

Malignancy

Patients with carcinoma of the ovary, brain and pancreas have a particularly increased risk of venous thrombosis but there is an increased risk with all cancers. The tumours produce tissue factor and a pro-coagulant that directly activates factor X. Mucin-secreting adenocarcinomas may be associated with DIC.

Inflammation

This up-regulates procoagulant factors, down-regulates anticoagulant pathways, particularly

protein C. Thrombosis is particularly likely in inflammatory bowel disease, Behçet's disease, systemic tuberculosis, systemic lupus erythematosus (SLE) and diabetes.

Blood disorders

Increased viscosity, thrombocytosis, altered platelet membrane receptors and responses are possible factors for the high incidence of thrombosis in patients with polycythaemia vera and essential thrombocythaemia. There is a high incidence of venous thrombosis, including thrombi in large veins (e.g. the hepatic vein) in patients with paroxysmal nocturnal haemoglobinuria. An increased tendency to venous thrombosis has been observed in patients with sickle cell disease and patients with post splenectomy thrombocytosis.

Oestrogen therapy

Oestrogen therapy, particularly high-dose therapy, is associated with increased plasma levels of factors II, VII, VIII, IX and X and depressed levels of anti-thrombin and tissue plasminogen activator in the vessel wall. There is a high incidence of postoperative venous thrombosis in women on high-dose oestrogen therapy and full-dose oestrogen-containing oral contraceptives. The risk is much less with low-dose oestrogen contraceptive preparations. Hormone replacement therapy also increases the risk of thrombosis, largely obviated by the use of low oestrogen preparations.

The antiphospholipid syndrome

This may be defined as the occurrence of thrombosis or recurrent miscarriage in association with laboratory evidence of persistent antiphospholipid antibody. One antiphospholipid antibody is the 'lupus anticoagulant' (LA) which was initially detected in patients with SLE, and is identified by a prolonged plasma APTT which does not correct with a 50 : 50 mixture of normal plasma. A second test dependent on limiting quantities of phospholipid (e.g. the dilute Russell's viper venom test) is also used in diagnosis. The antibody represents part of the spectrum of antiphospholipid antibody syndrome (APS) and whereas lupus anticoagulants are reactive in the fluid phase, the other antiphospholipid antibodies (APAs), such as anticardiolipin antibodies

Table 25.3 Clinical associations of lupus anticoagulant and anticardiolipin antibodies.

Venous thrombosis deep venous thrombosis/pulmonary embolism renal, hepatic, retinal veins
Arterial thrombosis
Recurrent fetal loss
Thrombocytopenia
Livedo reticularis

NB. Recurrent fetal loss may also occur in other types of thrombophilia.

(ACA) and antibodies to β_2 -glycoprotein (β_2 -GPI), are identified by solid phase immunoassay. Both solid phase assays and coagulation tests for LA should be used in the diagnosis of APS.

As well as patients with SLE, APS is also found in other autoimmune disorders particularly of connective tissues, lymphoproliferative diseases, post-viral infections, with certain drugs including phenothiazines and as an 'idiopathic' phenomenon in otherwise healthy subjects. Paradoxically, in view of its name, it is associated with venous and arterial thrombosis. The arterial thrombosis may cause peripheral limb ischaemia, stroke or myocardial infarct. As with other causes of thrombophilia, recurrent abortion caused by placental infarction is also associated (Table 25.3). Thrombocytopenia may be present and livedo reticularis is a frequent dermal manifestation. Treatment is with anticoagulation where indicated. It is usual to maintain an international normalized ratio (INR) of between 2.0 and 3.0 with warfarin but higher levels may be needed if previous arterial or major DVT has occurred or recurrence of thrombosis occurs on warfarin therapy. Low-dose heparin and aspirin are useful in the management of recurrent miscarriage.

Collagen vascular diseases and Behçet's syndrome are also associated with arterial and venous thrombosis, whether or not the lupus anticoagulant is present (p. 300).

Factor IX concentrates

Venous thrombosis may complicate the use of factor IX concentrates which contain trace amounts of activated coagulation factors. Patients with liver disease who are unable to clear these activated factors are especially at risk.

Glucosylceramide deficiency

Reduced plasma levels of the glycolipid glucosylceramide are a potential risk factor for venous thrombosis, particularly in young male patients. The glycolipid modulates the protein C pathway.

Investigation of thrombophilia

Many of the conditions associated with an increased thrombotic risk are obvious following clinical examination. A full assessment is indicated, particularly in patients who have recurrent or spontaneous DVT or pulmonary emboli, in patients who have thrombosis at a young age and in those patients with a familial tendency to thrombosis or thrombosis at an unusual site. It is also needed in women with recurrent fetal loss. With the increasing recognition of systemic causes of thrombophilia, the indications for thrombophilia screening are widening. The following laboratory tests are used in diagnosis.

Screening tests

1 Blood count and erythrocyte sedimentation rate (ESR)—to detect elevation in haematocrit, white cell count, platelet count, fibrinogen and globulins.

2 Blood film examination—may provide evidence of myeloproliferative disorder; leucoerythroblastic features may indicate malignant disease.

3 Prothrombin time (PT) and APTT—a shortened APTT is often seen in thrombotic states and may indicate the presence of activated clotting factors. A prolonged APTT test, not corrected by the addition of normal plasma, suggests a 'lupus anticoagulant' or an acquired inhibitor to a coagulation factor.

4 Anticardiolipin and anti- β_2 -GPI antibodies.

5 Thrombin time and reptilase time—prolongation suggests an abnormal fibrinogen.

6 Fibrinogen assay.

7 Activated protein C (APC) resistance test and DNA analysis for factor V Leiden.

8 Antithrombin—immunological and functional assays.

9 Protein C and protein S—immunological and functional assays.

10 Prothrombin gene analysis for the G20210A variant.

11 Plasma homocysteine estimation.

In many patients even full investigation yields no abnormalities and treatment with oral anticoagulants may remain empirical.

12 Test for CD59 and CD55 expression (paroxysmal nocturnal haemoglobinuria) in red cells if paroxysmal nocturnal haemoglobinuria is suspected.

Diagnosis of venous thrombosis

Deep vein thrombosis

Clinical suspicion DVT is suspected in those with previous DVT, cancer or confined to bed. In the leg, unilateral thigh or calf swelling or tenderness, pitting oedema and the presence of collateral superficial non-varicose veins are important signs.

Serial compression ultrasound This is the most reliable and practical method for patients with first suspicion of DVT in the legs and other sites (Fig. 25.2a). It can be combined with spectral, colour (Fig. 25.2) or power Doppler (duplex) scanning which improves accuracy by focusing on individual veins. It does not distinguish between acute and chronic thrombi. Persisting venous obstruction detected by ultrasonography at the completion of warfarin therapy is associated with an increased risk of recurrent thrombosis.

Contrast venography This most sensitive procedure is reserved for patients with highly suggestive clinical findings but negative ultrasonography. Iodinated contrast medium is injected into a vein peripheral to the suspected DVT. This permits direct demonstration by X-ray of the site, size and extent of the thrombus (Fig. 25.2b). However, it is a painful invasive technique, with a risk of contrast reaction and procedure-induced DVT.

Plasma D-dimer concentration The concentration of these fibrin breakdown products is raised when there is a fresh venous thrombosis. It is a useful assay when recurrent thrombosis is suspected and also may be combined with the tests above for diagnosis of a first event. A negative result is useful in emergency departments for excluding DVT in patients with equivocal clinical features. D-dimer elevation in cancer, inflammation after surgery or trauma limits its usefulness.

Magnetic resonance imaging (MRI) This may also be used but is expensive. Impedance plethysmography is less sensitive and accurate and is falling out of use.

Pulmonary embolus

Clinical suspicion This is particularly suspected in patients with chest symptoms, especially if there are signs of or previous history of DVT, immobilization for more than 2 days or recent (<4 weeks) surgery, haemoptysis or cancer.

Chest X-ray This is often normal but may show evidence of pulmonary infarction or pleural effusion.

Ventilation perfusion (VQ) scintigraphy This detects areas of the lung being ventilated but not perfused.

Computed tomography (CT) pulmonary angiography Fine slices of the lung are scanned by spiral CT so that filling defects in the pulmonary arteries are visualized (Fig. 25.2c).

Magnetic resonance pulmonary angiography Gadolinium-enhanced MRI is a relatively new, expensive but accurate technique.

Pulmonary angiography This is the traditional reference method but is invasive with complications, albeit uncommon, such as arrhythmia or contrast reaction.

Electrocardiogram This is performed to determine whether there is right heart 'strain' which occurs only in relatively severe cases.

Anticoagulant drugs

Anticoagulant drugs are used widely in the treatment of venous thromboembolic disease. Their value in the treatment of arterial thrombosis is less well established.

Heparin

This acidic, unfractionated mucopolysaccharide of average molecular weight (MW) 15 000–18 000 is an inhibitor of blood coagulation because of its action in potentiating the activity of antithrombin (see below). As it is not absorbed from the gastrointestinal tract it must be given by injection. It is inactivated by the liver and excreted in the urine. The effective biological half-life is approximately 1 h (Table 25.4).

Table 25.4 Comparison of unfractionated heparin with low molecular weight heparin.

	Unfractionated heparin	Low molecular weight heparin
Mean molecular weight in kilodaltons (range)	15 (4–30)	4.5 (2–10)
Anti-Xa : anti-IIa	1 : 1	2 : 1 to 4 : 1
Inhibits platelet function	Yes	No
Bioavailability	50%	100%
Half-life		
intravenous	1 h	2 h
subcutaneous	2 h	4 h
Elimination	Renal and hepatic	Renal
Monitoring	APTT	Xa assay (usually not needed)
Frequency of heparin-induced thrombocytopenia	High	Low
Osteoporosis	Yes	Less frequent

APTT, activated partial thromboplastin time.




Fig. 25.6 The action of heparin. This activates antithrombin which then forms complexes with activated serine protease coagulation factors (thrombin, Xa, IXa and XIa) and so inactivates them.

Mode of action

Heparin dramatically potentiates the formation of complexes between antithrombin and activated serine protease coagulation factors, thrombin (IIa) and factors IXa, Xa and XIa (Fig. 25.6). This complex formation inactivates these factors irreversibly. In addition, heparin impairs platelet function.

Low molecular weight heparin (LMWH) preparations (MW 2000–10 000) are produced by enzymatic or chemical depolymerization of unfractionated heparin. They have a greater ability to inhibit factor Xa than to inhibit thrombin and interact less with platelets compared with standard heparin, and so may have a lesser tendency to cause bleeding. They also have greater bioavailability and a more prolonged half-life in plasma, making once-daily administration in prophylaxis or treatment feasible (Table 25.4).

Indications

Heparin is routinely used in DVT, pulmonary embolism and unstable angina pectoris. It is also widely used in the prophylaxis of venous thrombosis and is the drug of choice for women requiring anticoagulation in pregnancy because it does not cross the placenta. It is also used during cardiopulmonary

bypass surgery, for maintaining the patency of indwelling venous lines and in some case of DIC if the manifestations are predominantly vaso-occlusive. There is some evidence that LMWH (or warfarin) may improve the survival in cancer patients over and above the protection from venous thrombosis but the exact mechanisms, types of cancer improved and optimum treatment regimes remain to be elucidated.

Administration and laboratory control

Standard heparin

Continuous intravenous infusion This provides the smoothest control of heparin therapy and is the treatment of choice where rapid reversal of anticoagulation by protamine sulphate may be required (e.g. in surgical patients or late pregnancy). It is still preferred by many for treatment of acute pulmonary embolus. In an adult, dosage of 30 000–40 000 units over 24 h (1000–2000 units/h with a loading dose of 5000 units) is usually satisfactory. Therapy is monitored by maintaining the APTT at 2–3 times the upper limit of the normal value. It is usual to start warfarin therapy within 2 days of starting heparin therapy and to discontinue heparin when the INR has been above 2.0 on two successive days. For acute coronary syndromes, both unfractionated heparin and LMWH are of benefit when used with aspirin in the prevention of mural thrombosis, systemic embolization and venous thrombosis.

Subcutaneous heparin Intermittent subcutaneous injections are preferred when heparin is given as prophylaxis against venous thrombosis (e.g. for surgical procedures). The usual dosage is 5000 units 12-hourly preoperatively followed by this dosage 8–12-hourly for 7 days or until the patient is mobile. LMWH (see below) is usually preferred as it can be given once daily.

Low molecular weight heparin

LMWH is given by subcutaneous injection and, as it has a longer half-life than standard heparin, it can be given once a day in prophylaxis, or once or twice a day in treatment (Table 25.4). Compared with unfractionated heparin, LMWH has a more

predictable dose-response which avoids the need for routine monitoring. There is less risk of thrombocytopenia or osteoporosis. It is the treatment of choice for preventing or treating DVT. It is used increasingly for treatment of pulmonary embolus and unstable angina. Although routine monitoring is not required, measurement of anti-Xa peak levels 4 h after injection allows dose adjustment in selected patients (e.g. in pregnancy, renal failure, gross obesity and in children). Typical treatment regimens are 200 anti-Xa units/kg once daily or 100 anti-Xa units/kg twice daily). LMWH is preferred to unfractionated heparin for therapy of DVT including pulmonary embolism. Many patients with uncomplicated DVTs may now be managed at home with regular LMWH injections once or twice daily according to the preparation. LMWH is the preferred treatment for the prevention of DVT in both medical and surgical patients. It is also the preferred anticoagulant in pregnancy because it does not cross the placenta. Typical once daily subcutaneous dosage in prophylaxis is 2000–2500 units (moderate risk patients) or 4000–5000 units (high-risk patients).

Complications

Unfractionated heparin is associated with a number of side-effects. The risk of these complications appears to be reduced by approximately 50% by the use of LMWH.

Bleeding during heparin therapy

Bleeding may be because of excessive prolonged anticoagulation or to an antiplatelet functional effect of heparin. Intravenous heparin has a half-life of less than 1 h and it is usually only necessary to stop the infusion. Protamine is able to inactivate heparin immediately and for severe bleeding a dose of 1 mg/100 units heparin provides effective neutralization. However, protamine itself may act as an anticoagulant when in excess.

Heparin-induced thrombocytopenia

A mild lowering of the platelet count may occur in the first 24 h as a result of platelet clumping. This is of no clinical consequence (heparin-induced thrombocytopenia, HIT, type 1). The important HIT, type 2, may occur in up to 5% of patients who are treated

with unfractionated heparin and paradoxically presents with thrombosis. It results from the binding of heparin to platelets followed by the generation of an immunoglobulin G (IgG) antibody to the heparin-platelet factor 4 (PF4) complex, which leads to platelet activation (Fig. 25.7). Typically, it presents as a fall of >50% in the platelet count 5 or more days after starting heparin treatment or earlier if heparin has been given previously. Diagnosis is difficult but assays have recently been developed to allow the detection of antibodies to immobilized heparin-PF4 complex. Heparin therapy must be discontinued. Thrombin inhibitors such as hirudin or lepirudin or argatroban appear promising as alternatives and the heparinoid danaparoid may also be used. LMWH is less likely than unfractionated heparin to cause HIT but there is cross-reactivity of the antibody. Warfarin therapy in some cases causes skin necrosis and should be delayed until alternative anticoagulation has been achieved.

Osteoporosis

This occurs with long-term (>2 months) heparin therapy, especially in pregnancy. The drug complexes minerals from the bones but the exact pathogenesis is unknown.

Oral anticoagulants

These are derivatives of coumarin or indandione. Warfarin, a coumarin, is most widely used. The drugs are vitamin K antagonists (p. 297) and so treatment results in decreased biological activity of the vitamin K-dependent factors II, VII, IX and X. Oral anticoagulants block the post-ribosomal γ -carboxylation of glutamic acid residues of these proteins (Fig. 24.7). After warfarin is given, factor VII levels fall considerably within 24 h but prothrombin has a longer plasma half-life and only falls to 50% of normal at 3 days; the patient is fully anticoagulated only after this period.

Principles of oral anticoagulation

A typical starting regimen for warfarin would be 10 mg on day 1, 5 mg on day 2 and then 5 mg on the third day. After this the dosage should be adjusted according to the PT. The usual maintenance dosage




Fig. 25.7 Mechanism of heparin-induced thrombocytopenia (HIT). Platelet factor 4 (PF4) is released from α granules and forms a complex on the platelet surface with heparin. Immunoglobulin G antibodies (usually IgG₂) develop against this complex and once bound can activate the platelet through the platelet immunoglobulin receptor Fc- γ RII. This leads to platelet stimulation, further release of PF4 and the platelet release reaction with consequent thrombocytopenia and thrombus development.

Table 25.5 Oral anticoagulant control tests. Target levels recommended by the British Society for Haematology (2000).

Target INR	Clinical state
2.5 (2.0–3.0)	Treatment of DVT, pulmonary embolism, atrial fibrillation, recurrent DVT off warfarin; symptomatic inherited thrombophilia, cardiomyopathy, mural thrombus, cardioversion
3.5 (3.0–4.0)	Recurrent DVT while on warfarin, mechanical prosthetic heart valves, antiphospholipid syndrome (some cases)

DVT, deep vein thrombosis; INR, international normalized ratio.

of warfarin is 3–9 mg/day but individual responses vary greatly. Lower loading dosage is recommended for the elderly or those with liver disease.

The indications and recommended ranges for INR with warfarin treatment are summarized in Table 25.5. The effect of oral anticoagulants is monitored by the PT. The INR is used and is based on the ratio of the patient's PT to a mean normal PT with correction for the 'sensitivity' of the thromboplastin used. This is calibrated against a primary World Health Organization (WHO) standard thromboplastin.

Warfarin crosses the placenta and is teratogenic. Heparin is preferred for pregnant patients because it does not cross the placenta and its action is short-lived.

It is usual to continue warfarin for 3–6 months for established DVT, pulmonary embolism and following xenograft heart valves. Long-term therapy is given for recurrent venous thrombosis, for embolic

complications of rheumatic heart disease or atrial fibrillation, and with prosthetic valves and arterial grafts. It is also given long term in patients with a severe cause of thrombophilia (e.g. the lupus anticoagulant and a history of thrombosis).

Drug interactions

Approximately 97% of warfarin in the circulation is bound to albumin and only a small fraction of warfarin is free and can enter the liver parenchymal cells; it is this free fraction that is active. In the liver cells, warfarin is degraded in microsomes to an inactive water-soluble metabolite which is conjugated and excreted in the bile and partially reabsorbed to be also excreted in urine. Drugs that affect the albumin binding or excretion of warfarin (or of other oral anticoagulants) or those that decrease the absorption of vitamin K will interfere with the control of therapy (Table 25.6).

Table 25.6 Drugs and other factors that interfere with the control of anticoagulant therapy.

Potentialiation of oral anticoagulants	Inhibition of oral anticoagulants
Drugs that increase the effect of coumarins <i>Reduced coumarin binding to serum albumin</i> Sulfonamides <i>Inhibition of hepatic microsomal degradation of coumarin</i> Cimetidine Allopurinol Tricyclic antidepressants Metronidazole Sulfonamides <i>Alteration of hepatic receptor site for drug</i> Thyroxine Quinidine <i>Decreased synthesis of vitamin K factors</i> High doses of salicylates Some cephalosporins Liver disease Decreased synthesis of vitamin K factors Decreased absorption of vitamin K e.g. malabsorption, antibiotic therapy, laxatives	Drugs that depress the action of coumarins <i>Acceleration of hepatic microsomal degradation of coumarin</i> Barbiturates Rifampicin <i>Enhanced synthesis of clotting factors</i> Oral contraceptives Hereditary resistance to oral anticoagulants Pregnancy

NB. Patients are also more likely to bleed if taking antiplatelet agents (e.g. NSAIDs, dipyridamole or aspirin); alcohol in large amounts enhances warfarin action.

Management of warfarin overdose

If the INR is in excess of 4.5 without bleeding, warfarin should be stopped for 1 or 2 days and the dose adjusted according to the INR. The long half-life of warfarin (40 h) delays the full impact of dose changes for 4–5 days. If the INR is very high (e.g. >8) without bleeding, an oral dose of 0.5–2.5 mg vitamin K may be given. Mild bleeding usually only needs an INR assessment, drug withdrawal and subsequent dosage adjustment (Table 25.7). More serious bleeding may need cessation of therapy, vitamin K therapy or the infusion of fresh frozen plasma or prothrombin concentrates. Vitamin K is the specific antidote; an oral or intravenous dose of 2.5 mg is usually effective. Higher doses result in resistance to further warfarin therapy for 2–3 weeks.

Management of surgery

For minor surgery (e.g. dental extraction) antico-

agulation can be maintained and mouth rinses with tranexamine acid given. For major surgery, warfarin is stopped to get an INR <1.5 and LMWH given when the INR falls to <2.0 (except on the day or surgery) and continued until the INR is >2.0 after restarting warfarin.

New anticoagulants

Traditional anticoagulant therapy has disadvantages. LMWH has to be given subcutaneously and warfarin requires frequent monitoring and dose adjustment and there is interaction with drugs and food.

Factor Xa inhibitors: Fondaparinux, a synthetic analogue of the antithrombin-binding pentasaccharide of heparin, is an indirect irreversible factor Xa inhibitor. It is given subcutaneously, has a plasma half-life of 17 h and does not require laboratory monitoring. Extensive trials have demonstrated that

Table 25.7 Recommendations on the management of bleeding and excessive anticoagulation by the British Committee for Standards in Haematology (third edition 1998; 2005 update).

INR 3.0–6.0 (target INR 2.5)	Reduce warfarin dose or stop
INR 4.0–6.0 (target INR 3.5)	Restart warfarin when INR <5.0
INR 6.0–8.0	Stop warfarin*
No bleeding or minor bleeding	Restart when INR <5.0
INR >8.0	Stop warfarin
No bleeding or minor bleeding	Restart warfarin when INR <5.0
	If other risk factors for bleeding give 0.5–2.5 mg vitamin K orally
Major bleeding	Stop warfarin
	Give prothrombin complex concentrate 50 units/kg, in preference
	FFP 15 mL/kg (when available)
	Give 5 mg vitamin K (i.v. or oral)

FFP, fresh frozen plasma; INR, international normalized ratio; i.v., intravenous.

* 1 mg vitamin K may be given orally to rapidly reduce the INR to the therapeutic range within 24 h in all patients with an INR above the therapeutic range and no bleeding.

it is an effective antithrombotic agent in selected patients for prevention and treatment of both venous and arterial thromboembolic disorders. It also reduces major bleeding and may improve long-term mortality and morbidity e.g. in acute coronary syndromes. *Idraparinux*, a chemically modified analogue of fondaparinux has a longer half-life and, as a once weekly injection, it may be more convenient than LMWH in treatment and prophylaxis, and has the potential to replace warfarin in selected patients.

Direct thrombin inhibitors These include recombinant hirudins, bivalirudin, argatroban, melagraton and ximelagraton.

Bivalirudin has been used as an alternative to heparin in patients undergoing percutaneous coronary interventions and is associated with less bleeding and a reduced need for adjunctive treatment with glycoprotein IIb/IIIa antagonists in patients undergoing stenting.

Ximelagraton, a lipophilic pro-drug may be given orally and following absorption is rapidly converted to melagraton. *Melagraton* may be given subcutaneously. Routine anticoagulation monitoring is unnecessary. Approximately 6–10% of patients develop reversible increases in liver enzymes. *Ximelagraton* has been compared with either LMWH or warfarin in large numbers of clinical

trials. It is as effective as conventional therapy in preventing venous thrombosis in orthopaedic and general surgery without an increased risk of major haemorrhage. It has had similar success in the treatment of acute venous thrombosis, myocardial infarction and severe recurrent myocardial ischaemia and in the prevention of stroke and systemic arterial embolism in patients with atrial fibrillation.

Before they are licensed, more data on their effectiveness and safety in a more diverse spectrum of patients are necessary. No specific antidote is available for either the pentasaccharides or ximelagraton.

Graduated compression stockings

These are used postoperatively and during long aeroplane flights to reduce the risk of DVT. After a DVT, they reduce the risk of post-thrombotic syndrome.

Inferior vena cava filter

This can provide short-term protection against pulmonary embolism when a DVT in the legs is diagnosed but anticoagulation is contraindicated (e.g. ongoing or very recent intracranial or gastrointestinal bleeding or where there is recurrent pulmonary embolism despite adequate anticoagulation).



Table 25.8 Fibrinoly

Streptokinase (SK)
Tissue plasminogen
Single-chain urokinase
(SCU-PA)
Acylated plasminogen
(APSAC)

Fibrinolytic agents

A number of fibrinolytic agents are available for patients with acute myocardial infarction, major pulmonary embolism, and locally arterial occlusion.

Administration is simplified with the use of these agents. The therapy is most effective if symptoms begin within 6 hours of onset. Streptokinase is given in a total dose of 1.5 million units followed by 0.5 million units/h for 12–48 h. Tissue plasminogen activator (alteplase) is given in a total dose of 100 mg over 60 min. The value of alteplase compared with streptokinase is still under evaluation.

The use of fibrinolytic agents for the control of short-term bleeding is still considered unnecessary. Complications such as hemorrhage and hypotension are rare with these agents (Table 25.9).

Table 25.9 Contraindications

Absolute contraindications

Active gastrointestinal bleeding
Aortic dissection
Head injury or cerebrovascular disease in the past 2 months
Neurosurgery in the past 3 months
Intracranial aneurysms or arteriovenous malformations
Proliferative diabetic retinopathy

Table 25.10 Antiplatelet therapy in patients with an acute coronary syndrome and in those undergoing percutaneous coronary intervention (PCI). (After Lange R.A. and Hillis L.D. (2004) Antiplatelet therapy for ischemic heart disease. *N Engl J Med* 350,277–80)

Drug	Target/patient group	Duration
<i>Acute coronary syndrome</i>		
Aspirin	All	Life long
Clopidogrel	All	9–12 months
<i>Glycoprotein IIb/IIIa inhibitors</i>		
Abciximab	None	–
Eptifibatide	High risk	48–72 h
Tirofiban	High risk	48–72 h
<i>Patients undergoing PCI</i>		
Aspirin	All	Life long
Clopidogrel	All	9–12 months
Abciximab	High risk	12 h after PCI
Eptifibatide	High risk	18–24 h after PCI
Tirofiban	None	–

monophosphate (cAMP) levels in circulating platelets which decreases their sensitivity to activating stimuli. Dipyridamole has been shown to reduce thromboembolic complications in patients with prosthetic heart valves and to improve the results in coronary bypass operations.

Sulfinpyrazone This drug is a competitive inhibitor of cyclo-oxygenase. It has been effective in reducing the frequency of blockage in arteriovenous shunts in chronic dialysis patients.

Ticlopidine This is an antiplatelet drug that was used following coronary angioplasty. Side-effects include neutropenia or thrombocytopenia. It has been replaced by clopidogrel except in patients intolerant of clopidogrel.

Clopidogrel This adenosine diphosphate (ADP) receptor antagonist is an antiplatelet agent in use for reduction of ischaemic events in patients with




Fig. 25.8 Sites of action of antiplatelet drugs. Aspirin acetylates the enzyme cyclo-oxygenase irreversibly. Sulfinpyrazone inhibits cyclo-oxygenase reversibly. Dipyridamole inhibits phosphodiesterase, increases cyclic adenosine monophosphate (cAMP) levels and inhibits aggregation. Inhibition of adenosine uptake by red cells allows adenosine accumulation in plasma which

stimulates platelet adenylate cyclase. Prostacyclin (epoprostenol) stimulates adenylate cyclase. The lipid-soluble β -blockers inhibit phospholipase. Calcium-channel antagonists block the influx of free calcium ions across the platelet membrane. Dextran coats the surface interfering with adhesion and aggregation. GP, glycoprotein.

ischaemic stroke, myocardial infarction or peripheral vascular disease. It is used after coronary artery stenting or angioplasty and in patients requiring long-term antiplatelet therapy who are intolerant or allergic to aspirin.

Glycoprotein IIb/IIIa inhibitors: abciximab, eptifibatide, tirofiban These drugs are monoclonal antibodies that inhibit the platelet GPIIb/IIIa receptor. They are used in conjunction with heparin and aspirin for the prevention of ischaemic complications in high-risk patients undergoing percutaneous transluminal coronary angioplasty. They can be used once only.

Specific antithrombotic agents Intravenous prostacyclin has been used in clinical trials in patients with peripheral vascular disease and thrombotic thrombocytopenic purpura. It has also reduced arteriovenous shunt blockage in haemodialysis patients.

Bibliography

- Baglin T.P., Keeling D.M., Watson H.G. (2006) Guidelines on oral anticoagulation (warfarin): third edition – 2005 update. *Br J Haematol* 132,277–85.
- Baglin T., Barrowcliffe T.W., Cohen A., Greave M. (2006) Guidelines on the use and monitoring of heparin. *Br J Haematol* 133,19–34.
- Bates S.M. and Weitz J.I. (2006) The status of new anticoagulants. *Br J Haematol* 134,3–19.
- Bønaa K.H., Njølstad I., Ueland P.M. *et al.* (2006) Homocysteine lowering and myocardial infarction. *N Engl J Med* 354,1573–88.
- Bounameaux H. and de Moerloose P. (2004) Is laboratory monitoring of low-molecular-weight heparin therapy necessary? No. *J Thromb Haemost* 2,551–4.
- Fedullo P.F. and Tapson V.F. (2003) The evaluation of suspected pulmonary embolism. *N Engl J Med* 349,1247–56.
- The Fifth Organization to assess strategies in acute ischemic syndromes investigators (2006) Comparison of fondaparinux and enoxaparin in acute coronary syndromes. *N Engl J Med* 354,1464–76.
- Geerts W.H., Pineo G.F., Heit J.A. *et al.* (2004) Prevention of venous thromboembolism: the Seventh ACCP Conference on antithrombotic and thrombolytic therapy. *Chest* 126,338S–400S.
- Greaves M., Cohen H., MacHinn S.J., Mackie I. (2000) Guidelines on the investigation and management of the antiphospholipid syndrome *Br J Haematol* 109,704–15.
- Hasan H.I. and Matson M.B. (2006) Inferior vena cava filters: current concepts. *Thrombus* 10,9–11.
- Hirsh J., Guyatt, G., Albers G.W., Schünemann H.J. (eds) (2004) The Seventh ACCP Conference on antithrombotic and thrombolytic therapy: evidence-based guidelines. *Chest* 126(Suppl.),172S–3S.
- Hirsh J., O'Donnell M., Weitz J.I. (2005) New anticoagulants. *Blood* 105,453–63.
- The HOPE-2 Investigators (2006) Homocysteine lowering and cardiovascular events after acute myocardial infarction. *N Engl J Med* 354,1567–77.
- Keeling D.M., Mackie I.J., Moody A., Watson H.G. (2004) The diagnosis of deep vein thrombosis in symptomatic outpatients and the potential for clinical assessment and D-dimer assays to reduce the need for diagnostic imaging. *Br J Haematol* 124,15–25.
- Kujovich J.L. (2004) Hormones and pregnancy: thromboembolic risks for women. *Br J Haematol* 126,443–54.
- Kyrle P.A. and Eichinger S. (2005) Deep vein thrombosis. *Lancet* 365,1163–74.
- Lane D.A. and Grant P.J. (2000) Role of hemostatic gene polymorphisms in venous and arterial thrombotic disease. *Blood* 95,1517–32.
- Lange R.A. and Hillis L.D. (2004) Antiplatelet therapy for ischemic heart disease. *N Engl J Med* 350,277–80. Erratum in (2004) *N Engl J Med* 351,200.
- Lecumberri R., Paramo J.A., Rocha E. (2005) Anticoagulant treatment and survival in cancer patients: the evidence from clinical studies. *Haematologica* 90,1258–66.
- Lissalde-Lavigne G., Cochery-Nouvellon E., Mercier E. *et al.* (2005) The association between hereditary thrombophilias and pregnancy loss. *Haematologica* 90,1223–30.
- Lockshin M.D. and Erkan D. (2003) Treatment of the antiphospholipid syndrome. *N Engl J Med* 349,1177–9.
- Love C.D.O. (2006) Can haematological tests predict cardiovascular risk? The 2005 Kettle Lecture. *Br J Haematol* 133,232–50.
- Schulman S. (2003) Clinical practice. Care of patients receiving long-term anticoagulant therapy. *N Engl J Med* 349,675–83.
- Stein P.D., Fowler S.E., Goodman L.R. *et al.* (2006) Multidetector computed tomography for acute pulmonary embolism. *N Engl J Med* 354,2317–27.
- Stein P.D., Hull R.D., Patel K.C. *et al.* (2004) D-dimer for the exclusion of acute venous thrombosis and pulmonary embolism: a systematic review. *Ann Intern Med* 140,589–602.
- Wells P.S., Anderson D.R., Rodger M. *et al.* (2003) Evaluation of D-dimer in the diagnosis of suspected deep-vein thrombosis. *N Engl J Med* 349,1227–35.
- Wood K. (ed) For the British Committee for Standards in Haematology. (2000) *Standard Haematology Practice 2: Guidelines in Oral Anticoagulation*. Blackwell Science, Oxford: pp. 104–29.

CHAPTER 26

26

Haematological changes in systemic disease

Anaemia of chronic disorders, 320
Malignant diseases, 320
Rheumatoid arthritis, 323
Renal failure, 324
Liver disease, 325

Hypothyroidism, 326
Infections, 326
Inborn errors of metabolism, 331
Non-specific monitoring of systemic disease, 333
Bibliography, 336

Anaemia of chronic disorders

Many of the anaemias seen in clinical practice occur in patients with systemic disorders and are the result of a number of contributing factors. The anaemia of chronic disorders (also discussed on p. 39) is of central importance and occurs in patients with a variety of chronic inflammatory and malignant diseases (Table 26.1). Usually, both the erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP) are raised. It may be complicated by additional features which may be because of disease affecting particularly one or other system. The characteristic features are described in Chapter 3.

The pathogenesis of this anaemia appears to be related to the decreased release of iron from

macrophages to plasma and so to erythroblasts caused by hepcidin, reduced red cell lifespan and an inadequate erythropoietin response to anaemia. The plasma levels of various cytokines, especially interleukin-1 (IL-1), IL-6 and tumour necrosis factor (TNF) are raised and reduce erythropoietin secretion. The anaemia is corrected by the successful treatment of the underlying disease. It does not respond to iron therapy despite the low serum iron. Responses to recombinant erythropoietin therapy may be obtained (e.g. in rheumatoid arthritis or cancer). In many conditions the anaemia is complicated by anaemia from other causes (e.g. iron or folate deficiency, renal failure, bone marrow infiltration, hypersplenism or endocrine abnormality).

Malignant diseases (other than primary bone marrow diseases)

Anaemia

Contributing factors include anaemia of chronic disorders, blood loss and iron deficiency, marrow infiltration (Fig. 26.1) often associated with a leucoerythroblastic blood film (nucleated red cells and granulocyte precursors in the blood film), folate deficiency, haemolysis and marrow suppression from radiotherapy or chemotherapy (Table 26.2). Other causes of a leucoerythroblastic anaemia include

Table 26.1 Causes of anaemia of chronic disorders.

Chronic inflammatory diseases

Infectious (e.g. pulmonary abscess, tuberculosis, osteomyelitis, pneumonia, bacterial endocarditis)
Non-infectious (e.g. rheumatoid arthritis, systemic lupus erythematosus and other connective tissue diseases, sarcoid, Crohn's disease, cirrhosis)

Malignant disease

(e.g. carcinoma, lymphoma, sarcoma, myeloma)




Fig. 26.1 Metastatic carcinoma in bone marrow aspirates: (a) breast; (b) stomach; (c) colon; bone marrow trephine biopsies: (d) prostate; (e) stomach; (f) kidney.

myelofibrosis, acute and chronic leukaemias, and severe haemolytic or megaloblastic anaemia.

Microangiopathic haemolytic anaemia (p. 69) occurs with mucin-secreting adenocarcinoma (Fig. 26.2), particularly of the stomach, lung and breast. Less common forms of anaemia with malignant disease include autoimmune haemolytic anaemia with malignant lymphoma and rarely with other

tumours; primary red cell aplasia with thymoma or lymphoma; and myelodysplastic syndromes secondary to chemotherapy. There is also an association of pernicious anaemia with carcinoma of the stomach.

The anaemia of malignant disease may respond partly to erythropoietin. Folic acid should only be given if there is definite megaloblastic anaemia caused by the deficiency; it might 'feed' the tumour.

Table 26.2 Haematological abnormalities in malignant disease.

Haematological abnormality	Tumour or treatment associated
<i>Pancytopenia</i>	
Marrow hypoplasia	Chemotherapy, radiotherapy
Myelodysplasia	Chemotherapy, radiotherapy
Leucoerythroblastic	Metastases in marrow
Megaloblastic	Folate deficiency B ₁₂ deficiency (carcinoma of stomach)
<i>Red cells</i>	
Anaemia of chronic disorders	Most forms
Iron deficiency anaemia	Especially gastrointestinal, uterine
Pure red cell aplasia	Thymoma
Immune haemolytic anaemia	Lymphoma, ovary, other tumours
Microangiopathic haemolytic anaemia	Mucin-secreting carcinoma
Polycythaemia	Kidney, liver, cerebellum, uterus
<i>White cells</i>	
Neutrophil leucocytosis	Most forms
Leukaemoid reaction	Disseminated tumours, those with necrosis
Eosinophilia	Hodgkin's disease, others
Monocytosis	Various tumours
<i>Platelets and coagulation</i>	
Thrombocytosis	Gastrointestinal tumours with bleeding, others
Disseminated intravascular coagulation	Mucin-secreting carcinoma, prostate
Activation of fibrinolysis	Prostate
Acquired inhibitors of coagulation	Most forms
Paraprotein interfering with platelet function	Lymphomas, myeloma
Tumour cell procoagulants—tissue factor and cancer procoagulant (density activates factor X)	Especially ovarian, pancreas, brain, colon




Fig. 26.2 Peripheral blood film in metastatic mucin-secreting adenocarcinoma of the stomach showing red cell polychromasia and fragmentation and thrombocytopenia. The patient had disseminated intravascular coagulation.

Polycythaemia

Secondary polycythaemia is occasionally associated with renal, hepatic, cerebellar and uterine tumours (p. 236).

White cell changes

Leukaemoid reactions (p. 102) may occur with tumours showing widespread necrosis and inflammation. Hodgkin's disease is associated with a variety of white cell abnormalities including eosinophilia, monocytosis and leucopenia. In non-Hodgkin's lymphoma, malignant cells may circulate in the blood (p. 194).

Platelet and blood coagulation abnormalities

Patients with malignant disease may show either thrombocytosis or thrombocytopenia. Disseminated tumours, particularly mucin-secreting adenocarcinomas, are associated with disseminated intravascular coagulation (DIC; p. 298) and generalized haemostatic failure. Activation of fibrinolysis occurs in some patients with carcinoma of the prostate. Occasional patients with malignant disease have spontaneous bruising or bleeding caused by an acquired inhibitor of one or other coagulation factor, most frequently factor VIII, or to a paraprotein interfering with platelet function.

Cancer patients have a high incidence (estimated at 15%) of venous thromboembolism. This is increased by surgery and some drugs, most recently thalidomide. It is most common in ovarian, brain, pancreatic and colon cancers. It may be difficult to manage with oral anticoagulation because of bleeding, interruptions with chemotherapy and thrombocytopenia, anorexia or vomiting. Liver disease and drug interactions can cause further complications so daily low molecular weight heparin injections may be preferable to warfarin.

Rheumatoid arthritis (and other connective tissue disorders)

In patients with rheumatoid arthritis, the anaemia of chronic disorders is proportional to the severity of the disease. It is complicated in some patients by



(a)



(b)

Fig. 26.3 Felty's syndrome: (a) the typical deformities of rheumatoid arthritis of the hand and (b) splenomegaly.

iron deficiency caused by gastrointestinal bleeding related to therapy with salicylates, non-steroidal anti-inflammatory agents or corticosteroids. Bleeding into inflamed joints may also be a factor. Marrow hypoplasia may follow therapy with gold. In Felty's syndrome, splenomegaly is associated with neutropenia (Fig. 26.3). Anaemia and thrombocytopenia may also be present. In systemic lupus erythematosus (SLE) there may be anaemia of chronic disorders and 50% of patients are leucopenic with reduced neutrophil and lymphocyte counts often associated with circulating immune complexes. Renal impairment and drug-induced gastrointestinal blood loss also contribute to the anaemia. Autoimmune haemolytic anaemia (typically with immunoglobulin G (IgG) and the C3 component of complement on the surface of the red cells) occurs in 5% of patients and may be the presenting feature of the syndrome. There may be autoimmune

thrombocytopenia in 5% of patients. The lupus anticoagulant is described on p. 300. This circulating anticardiolipin interferes with blood coagulation by altering the binding of coagulation factors to platelet phospholipid and predisposes to both arterial and venous thrombosis and recurrent abortions. The antibody may be responsible for a false positive Wassermann reaction. Tests for antinuclear factor (ANF) and anti-DNA antibodies are usually positive.

Patients with temporal arteritis and polymyalgia rheumatica have a markedly elevated ESR, pronounced red cell rouleaux in the blood film and a polyclonal immunoglobulin response. These and other collagen vascular disorders are associated with anaemia of chronic disorders.

Renal failure

Anaemia

A normochromic anaemia is present in most patients with chronic renal failure. Generally, there is a 2 g/dL fall in haemoglobin level for every 10 mmol/L rise in blood urea. There is impaired red cell production as a result of defective erythropoietin secretion (see Fig. 2.5). Uraemic serum has also been shown to contain factors that inhibit proliferation of erythroid progenitors but, in view of the excellent response to erythropoietin in most patients, the clinical relevance of these is doubtful. Variable shortening of red cell lifespan occurs and

in severe uraemia the red cells show abnormalities including spicules (spurs) and 'burr' cells (Fig. 26.4). Increased red cell 2,3-diphosphoglycerate (2,3-DPG) levels in response to the anaemia and hyperphosphataemia result in decreased oxygen affinity and a shift of the haemoglobin oxygen dissociation curve to the right (p. 18), which is augmented by uraemic acidosis. The patient's symptoms are therefore relatively mild for the degree of anaemia.

Other factors may complicate the anaemia of chronic renal failure (Table 26.3): the anaemia of chronic disorders, iron deficiency from blood loss during dialysis or caused by bleeding because of defective platelet function, and folate deficiency in some chronic dialysis patients. Aluminium excess in patients on chronic dialysis also inhibits erythropoiesis. Patients with polycystic kidneys usually have retained erythropoietin production and may have less severe anaemia for the degree of renal failure.

Treatment

Erythropoietin therapy has been found to correct the anaemia in patients on dialysis or in chronic renal failure, providing that iron and folate deficiency, aluminium excess and infections have been corrected. The dosage of erythropoietin usually required is 50–150 units/kg three times a week intravenously or by subcutaneous infusion. The response is faster after intravenous administration, but greater with the subcutaneous route.




Fig. 26.4 Peripheral blood film in chronic renal failure showing red cell acanthocytosis and numerous 'burr' cells.

Table 26.3 Haematological abnormalities in renal failure.

Anaemia

Reduced erythropoietin production

Aluminium excess in dialysis patients

Anaemia of chronic disorders

Iron deficiency

blood loss (e.g. dialysis, venesection, defective platelet function)

Folate deficiency

chronic haemodialysis without replacement therapy

*Abnormal platelet function**Thrombocytopenia*

Immune complex-mediated (e.g. systemic lupus erythematosus, polyarteritis nodosa)

Some cases of acute nephritis and following allograft

Haemolytic uraemic syndrome and thrombotic thrombocytopenic purpura

Thrombosis

Some cases of the nephrotic syndrome

Polycythaemia

In renal allograft recipients

Rarely in renal cell carcinoma, cysts, arterial disease

Maintenance by 75 units/kg/week subcutaneously is typical. Complications of therapy have been initial transient flu-like symptoms, hypertension, clotting of the dialysis lines and, rarely, fits. A poor response to erythropoietin suggests iron or folate deficiency, infection, aluminium toxicity or hyperparathyroidism.

Platelet and coagulation abnormalities

A bleeding tendency with purpura, gastrointestinal or uterine bleeding occurs in 30–50% of patients with chronic renal failure and is marked in patients with acute renal failure. The bleeding is out of proportion to the degree of thrombocytopenia and has been associated with abnormal platelet or vascular function, which can be reversed by dialysis. Correction of the anaemia with erythropoietin also improves the bleeding tendency. Immune complex-mediated thrombocytopenia occurs in some patients with acute nephritis, SLE and polyarteritis nodosa and also following renal allografts. Renal allografts may also lead to polycythaemia in 10–15% of patients.

The haemolytic uraemic syndrome and thrombotic thrombocytopenic purpura are discussed on p. 284. Patients with the nephrotic syndrome have an increased risk of venous thrombosis.

Liver disease

The haematological abnormalities in liver disease are listed in Table 26.4. Chronic liver disease is associated with anaemia that is mildly macrocytic and often accompanied by target cells, mainly as a result of increased cholesterol in the membrane (Fig. 26.5a). Contributing factors to the anaemia may include blood loss (e.g. bleeding varices) with iron deficiency, dietary folate deficiency and direct suppression of haemopoiesis by alcohol. Alcohol may have an inhibiting effect on folate metabolism and is occasionally associated with (ring) sideroblastic changes which disappear when alcohol is withdrawn.

Haemolytic anaemia may occur in patients with alcohol intoxication (Zieve's syndrome) (Fig. 26.5b) and in Wilson's disease (caused by copper oxidation of red cell membranes) and autoimmune haemolytic anaemia is found in some patients with

Table 26.4 Haematological abnormalities in liver disease.

Liver failure ± obstructive jaundice ± portal hypertension

Refractory anaemia—usually mildly macrocytic, often with target cells; may be associated with:

Blood loss and iron deficiency

Alcohol (± ring sideroblastic change)

Folate deficiency

Haemolysis (e.g. Zieve's syndrome, Wilson's disease, immune hypersplenism from portal hypertension)

Bleeding tendency

Deficiency of vitamin K-dependent factors; also of factor V and fibrinogen

Thrombocytopenia hypersplenism, immune platelet function defects

Functional abnormalities of fibrinogen

Increased fibrinolysis

Portal hypertension—haemorrhage from varices

Viral hepatitis

Aplastic anaemia

Tumours

Polycythaemia

Neutrophil leucocytosis and leukaemoid reactions



Fig. 26.5 Liver disease: peripheral blood film showing: (a) macrocytosis and target cells; and (b) marked acanthocytosis in Zieve's syndrome.

chronic immune hepatitis. Haemolysis may also occur in end-stage liver disease because of abnormal red cell membranes resulting from lipid changes. Viral hepatitis (usually non-A, non-B, non-C) is associated with aplastic anaemia.

The acquired coagulation abnormalities associated with liver disease are described on p. 297. There are deficiencies of vitamin K-dependent factors (II, VII, IX and X) and, in severe disease, of factor V and fibrinogen. Thrombocytopenia may occur from hypersplenism or from immune complex-mediated platelet destruction. Abnormalities of platelet function may also be present. Dysfibrinogenaemia with abnormal fibrin polymerization may occur as a result of excess sialic acid in the fibrinogen molecules. A consumptive coagulopathy may be superimposed. These haemostatic defects may contribute to major blood loss from bleeding varices caused by portal hypertension.

Hypothyroidism

A moderate anaemia is usual and may be caused by lack of thyroxine. T_3 and T_4 potentiate the action of erythropoietin. There is also a reduced oxygen need and thus reduced erythropoietin secretion. The anaemia is often macrocytic and the mean corpuscular volume (MCV) falls with thyroxine therapy. Autoimmune thyroid disease, especially myxoedema or Hashimoto's disease, is associ-

ated with pernicious anaemia. Iron deficiency may also be present, particularly in women with menorrhagia.

Infections

Haematological abnormality is usually present in patients with infections of all types (Table 26.5). The effect of inflammation as a prothrombotic stimulus is also discussed on p. 308.

Bacterial infections

Acute bacterial infections are the most common cause of neutrophil leucocytosis. Toxic granulation, Döhle bodies and metamyelocytes may be present in the blood. Leukaemoid reactions with a white cell count $>50 \times 10^9/L$ and granulocyte precursors in the blood may occur in severe infections, particularly in infants and young children. The neutrophil alkaline phosphatase (NAP) score is raised in contrast to the low NAP score in chronic myeloid leukaemia. Mild anaemia is common if the infection is prolonged. Severe haemolytic anaemia occurs in bacterial septicaemias, particularly those caused by Gram-negative organisms, where there is usually associated DIC (p. 298). DIC dominates the clinical picture with certain infections (e.g. bacterial meningitis). The acute phase response to infections is accompanied by a rise in coagulation factors

Table 26.5 Blood abnormalities associated with infections.

Haematological abnormality	Infection associated
<i>Anaemia</i>	
Anaemia of chronic disorders	Chronic infections especially tuberculosis
Aplastic anaemia	Viral hepatitis
Transient red cell aplasia	Human parvovirus
Marrow fibrosis	Tuberculosis
Immune haemolytic anaemia	Infectious mononucleosis, <i>Mycoplasma pneumoniae</i>
Direct red cell damage or microangiopathic	Bacterial septicaemia (associated DIC), <i>Clostridium perfringens</i> , malaria, bartonellosis
	Viruses—haemolytic uraemic syndrome and TTP
Hypersplenism	Chronic malaria, tropical splenomegaly syndrome, leishmaniasis, schistosomiasis
<i>White cell changes</i>	
Neutrophil leucocytosis	Acute bacterial infections
Leukaemoid reactions	Severe bacterial infections particularly in infants
	Tuberculosis
Eosinophilia	Parasitic diseases (e.g. hookworm, filariasis, schistosomiasis, trichinosis)
	Recovery from acute infections
Monocytosis	Chronic bacterial infections: tuberculosis, brucellosis, bacterial endocarditis, typhoid
Neutropenia	Viral infections—HIV, hepatitis, influenza
	Fulminant bacterial infections (e.g. typhoid, miliary tuberculosis)
Lymphocytosis	Infectious mononucleosis, toxoplasmosis, cytomegalovirus, rubella, viral hepatitis, pertussis, tuberculosis, brucellosis
Lymphopenia	HIV infection
	<i>Legionella pneumonophila</i>
<i>Thrombocytopenia</i>	
Megakaryocytic depression, immune complex-mediated and direct interaction with platelets	Acute viral infections particularly in children (e.g. measles, varicella, rubella, malaria, severe bacterial infection)
Prothrombotic state	All with prolonged inflammation

DIC, disseminated intravascular coagulation; HIV, human immunodeficiency virus; TTP, thrombotic thrombocytopenic purpura.

and a fall in natural anticoagulants. Recombinant human activated protein C reduces the death rate of severe DIC.

Clostridium perfringens organisms produce an α toxin, a lecithinase acting directly on the circulating red cells (Fig. 26.6). Haemolysis in bartonellosis (Oroya fever) is caused by direct red cell infection. With severe acute bacterial infections there may be thrombocytopenia. *Mycoplasma pneumoniae* infections are associated with autoimmune haemolytic anaemia of the 'cold' type (p. 67).

Chronic bacterial infections are associated with the anaemia of chronic disorders. In tuberculosis, additional factors in the pathogenesis of anaemia include marrow replacement and fibrosis associated with miliary disease and reactions to anti-tuberculous therapy (e.g. isoniazid is a pyridoxine antagonist and may cause sideroblastic anaemia). Disseminated tuberculosis is associated with leukaemoid reactions and patients with involvement of bone marrow may show leucoerythroblastic changes in the peripheral blood film.




Fig. 26.6 Peripheral blood film in a patient with haemolytic anaemia in clostridial septicaemia showing red cell contraction and spherocytosis.

Viral infections

Acute viral diseases are often associated with a mild anaemia. An immune haemolytic anaemia with an anti-i autoantibody is associated with infectious mononucleosis (p. 68). Viral infections, as well as syphilis, have been associated with paroxysmal cold haemoglobinuria (p. 70). Viruses have also been linked to the pathogenesis of the haemolytic uraemic syndrome, thrombotic thrombocytopenic purpura (pp. 284 and 285) and the haemophagocytic syndrome (p. 106). Aplastic anaemia may occur with viral A or more usually non-A, non-B, non-C hepatitis. Transient red cell aplasia is associated with human parvovirus infection and this may result in severe anaemia in patients with a haemolytic anaemia because of the shortened red cell survival (e.g. in hereditary spherocytosis or sickle cell disease; p. 247).

Acute thrombocytopenia is not uncommon in rubella, morbilli and varicella infections. Rubella and cytomegalovirus (CMV) infections may cause a reactive lymphocytosis similar to that found in infectious mononucleosis. CMV may be responsible for a post-transfusion mononucleosis-like syndrome, CMV being transmitted by leucocytes. CMV infections in infants are associated with massive hepatosplenomegaly. In bone marrow transplant recipients or other immunosuppressed patients, CMV infections may cause pancytopenia as well as other severe disorders (e.g. pneumonitis or hepatitis; p. 258).

HIV infection


This is associated with a wide range of haematological changes. These are caused by marrow defects and immune cytopenias directly resulting from HIV infection, the effects of opportunistic infections or lymphoma and the side-effects of drugs used to treat HIV itself or drugs for the complicating infection or lymphoma.

Thrombocytopenia and neutropenia may be immune or secondary to marrow dysfunction. The marrow may be hypercellular with prominent plasma cells and lymphocytes, normocellular, hypocellular or fibrotic. Dysplastic features are common, with ineffective thrombopoiesis or granulocyte formation accounting at least in part for the cytopenia. The myelodysplasia does not show the chromosome abnormalities found in classic myelodysplasia and does not appear to be pre-leukaemic. Thrombocytopenia is treated if necessary by corticosteroids, high-dose gammaglobulin infusions or by other therapies for immune thrombocytopenia (p. 281), or by antiretroviral therapy.


Anaemia is also common, like the other cytopenias, and more severe as the disease progresses. It is usually multifactorial in origin—the anaemia of chronic disorders, marrow dysplasia and drug therapy, especially zidovudine. Serum vitamin B₁₂ is often low, most likely because of intestinal malabsorption, but the anaemia does not respond to vitamin B₁₂ therapy. Blood transfusions or recombinant erythropoietin injections are needed.

Increased plasma cells in the marrow and polyclonal increase in immunoglobulins are frequent. A paraprotein is present in 5–10% but appears benign. Non-Hodgkin's lymphoma, in over 90% high grade, both systemically and in the central nervous system, occurs in HIV infected individuals with over 100 times the frequency expected in the general population. Diffuse large B-cell lymphomas are most common, with 20% confined to the central nervous system. Approximately one-third are

Burkitt's lymphoma. Hodgkin's disease, usually of poor prognosis type, is also increased in frequency. EBV infection appears to underlie this as well as Burkitt's lymphoma. Treatment of lymphomas in the setting of HIV is with standard regimens. Continuation of antiretroviral therapy exaggerates the tendency to cytopenia induced by chemotherapy so prophylaxis against opportunistic infections is important. The bone marrow may indeed reveal the presence of opportunistic infection (Fig. 26.7a,b).



(a)



(b)

Fig. 26.7 Human immunodeficiency virus (HIV) infection: bone marrow trephine biopsy. (a) Granuloma showing positivity with Ziehl-Nielsen stain. (b) Higher power shows large numbers of acid-fast bacilli




Fig. 26.8 Malaria: peripheral blood in severe *Plasmodium falciparum* infection showing: (a) many ring forms and a meront; and at higher magnification: (b) a meront; and (c) a gametocyte.

Malaria

Some degree of haemolysis is seen in all types of malarial infection. The most severe abnormalities are found in *Plasmodium falciparum* infections (Fig. 26.8). In the worst cases, DIC occurs and intravascular haemolysis is marked with haemoglobinuria. This may be associated with quinine therapy ('blackwater fever'). Thrombocytopenia is commonly found in acute malaria. Patients with chronic malaria have an anaemia of chronic disorders; hypersplenism may contribute to the anaemia and result in moderate thrombocytopenia and neutropenia. Tropical splenomegaly (p. 125) is

probably a chronic immune reaction to malaria. Dyserythropoiesis in the marrow, folate deficiency and protein-calorie malnutrition may contribute to the anaemia.

Toxoplasmosis

Toxoplasmosis in children and adults is associated with lymphadenopathy and large numbers of atypical lymphocytes in the blood. Congenital disease may be confused with hydrops fetalis in a severely anaemic hydropic infant with gross hepatosplenomegaly, thrombocytopenia or a leucoerythroblastic blood film.




Fig. 26.9 Kala-azar: bone marrow aspirates showing macrophages containing Leishman-Donovan bodies.

Kala-azar (visceral leishmaniasis)

The visceral form of leishmaniasis is associated with pancytopenia, hepatosplenomegaly and lymphadenopathy. Bone marrow or splenic aspirates may show large numbers of parasitized macrophages (Fig. 26.9).

Other parasitic diseases

In the acute phase of both African and South American trypanosomiasis, organisms are found in the peripheral blood (Fig. 26.10). Microfilariae of bancroftian filariasis and loiasis are also detected during blood film examination (Fig. 26.11). In chronic schistosomiasis, hypersplenism follows the splenic enlargement associated with portal hypertension. In many parasitic diseases there is eosinophilia (Table 26.5).

Inborn errors of metabolism

Gaucher's, Tay-Sachs and Niemann-Pick diseases all result from hereditary deficiency of the enzymes required for glycolipid breakdown.

Gaucher's disease

Gaucher's disease is an uncommon autosomal recessive disorder characterized by an accumulation of glucosylceramide in the lysosomes of



Fig. 26.10 African trypanosomiasis: blood film showing *Trypanosoma brucei*.




Fig. 26.11 Peripheral blood films showing microfilariae of: (a) *Wuchereria bancrofti*; and (b) *Loa loa*.

reticuloendothelial cells as a result of deficiency of glucocerebrosidase (Fig. 26.12). Three types occur: a chronic adult type (type I); an acute infantile neuronopathic type (type II); and a subacute neuronopathic type with onset in childhood or adolescence (type III). Type I is caused by a variety of mutations in the glucocerebrosidase gene, one type of which (a single base pair substitution in codon 444) is particularly common in Ashkenazi Jews and explains the high incidence of the disease in this group. In type I the outstanding physical sign is splenomegaly. Moderate liver enlargement and pingueculae (conjunctival deposits) are other characteristic findings. In many cases, bone deposits cause bone pain and pathological fractures. Expansion of the lower end of the femur may produce the 'Erlenmeyer flask deformity' (Fig. 26.13c).

The clinical manifestations are caused by the accumulation of glucocerebroside-laden macrophages in the spleen, liver and bone marrow (Fig. 26.13a,b,e). Gaucher's disease at all ages is commonly associated with marked anaemia, leucopenia and thrombocytopenia occurring singly or in combination. Polyclonal hypergammaglobulinemia or monoclonal gammopathy are frequent. Diagnosis is made by assay of white cell glucocerebrosidase and DNA analysis. Lysosomal enzymes, chitotriosidase and acid phosphatase are raised and useful in monitoring therapy. Pulmonary activation-regulated cytokine (PARC), angiotensin-converting enzyme (ACE) and ferritin are also elevated.

Enzyme replacement therapy with glucocerebrosidase made by recombinant technology and given intravenously is very effective in treating the




Fig. 26.12 Gaucher's disease results from a deficiency of glucocerebrosidase. Gal, Galactose; Glc, glucose.

disease with shrinkage of spleen, rise in blood count and improved bone structure (Fig. 26.13d,f). An oral drug, miglustat, is useful in mild forms. It reduces the amount of substrate being produced in lysosomes and may be used in combination with the intravenous enzyme. Splenectomy can result in haematological improvement but following this operation there is often increased deposition of cerebroside in extrasplenic tissue, particularly bones. Stem cell transplantation has been carried out successfully in severely affected patients, usually with type II or III disease.

Niemann–Pick disease

Niemann–Pick disease shows certain clinical and pathological similarities to Gaucher's disease. It is caused by a sphingomyelinase deficiency. The majority of patients are infants who die in the first few years of life although occasional patients survive to adult life. Massive hepatosplenomegaly occurs and there is usually lung and nervous system involvement with retarded physical and mental development. A 'cherry-red' spot is commonly seen in the retina of affected infants. Pancytopenia is a regular feature and in marrow aspirates 'foam cells' of similar size to Gaucher cells are seen. Chemical analysis of the tissues reveals that the disorder is caused by an accumulation of sphingomyelin and cholesterol.

Non-specific monitoring of systemic disease

The inflammatory response to tissue injury includes changes in plasma concentrations of proteins

known as acute phase proteins. These proteins include fibrinogen and other clotting factors, complement components and CRP (p. 334), haptoglobin, serum amyloid A (SAA) protein, ferritin and others. The rise in these liver-derived proteins is part of a wider response which includes fever, leucocytosis and increased immune reactivity. The acute phase response is mediated by cytokines (e.g. IL-1; see Fig. 7.4) and TNF, released from macrophages and possibly other cells. Patients with chronic disease may show periodic or continuous evidence of the acute phase response depending upon the extent of inflammation. Quantitative measurements of acute phase proteins are valuable indicators of the presence and extent of inflammation and of its response to treatment. When short-term (less than 24 h) changes in the inflammatory response are expected CRP is the test of choice (Table 26.6). Long-term changes in the acute phase proteins are monitored by either the ESR or plasma viscosity. These tests are influenced by plasma proteins which are either slowly responding acute phase reactants (e.g. fibrinogen) or are not acute phase proteins (e.g. immunoglobulins).

Erythrocyte sedimentation rate

This commonly used but non-specific test measures the speed of sedimentation of red cells in plasma over a period of 1 h. The speed is mainly dependent on the plasma concentration of large proteins (e.g. fibrinogen and immunoglobulins). The normal range in men is 1–5 mm/h and in women 5–15 mm/h but there is a progressive increase in old age. The ESR is raised in a wide variety of systemic inflammatory and neoplastic diseases and in pregnancy.




Fig. 26.13 Gaucher's disease: (a) bone marrow aspirate—a Gaucher cell with 'fibrillar' cytoplasmic pattern; (b) spleen histology—pale clusters of Gaucher cells in the reticuloendothelial cord; (c) magnetic resonance imaging (MRI) scan of the left knee of a patient before treatment showing Erlenmeyer flask deformity with expansion of the marrow and thinning of the cortical bone; (d) following a year of glucocerebrosidase therapy with subsequent remodelling of bone; and bone marrow trephine biopsy before (e) and after (f) 2 years of glucocerebrosidase therapy.

Table 26.6 Advantages and disadvantages of the tests used to monitor the acute phase response.

Advantages	Disadvantages
<i>CRP*</i>	
Specific test of acute phase protein	More than one protein required to measure acute (CRP) and chronic inflammation
Fast response (6 h) to change in disease activity	Costly when assayed in small numbers
High sensitivity—owing to large incremental change	Sophisticated equipment and antisera required
Can be measured on stored serum	
Small sample volumes	
Automated analysis	
<i>ESR and plasma viscosity</i>	
Useful in chronic disease	Not sensitive to acute changes (<24 h)
ESR inexpensive, easy, no electrical power required	Not specific for acute phase response
Plasma viscosity—result obtained quickly (15 min)	Slow to change with alteration in disease activity and insensitive to small changes in activity
Plasma viscosity not affected by anaemia	Fresh samples (<2 h) required for ESR

ESR, erythrocyte sedimentation rate.

* C-reactive protein (CRP) is normally present in low concentrations (<5 mg/L). Levels are not influenced by anaemia, pregnancy or heart failure. During severe acute infection the plasma concentration may rise 100-fold.

It is useful for diagnosing and monitoring temporal arteritis and polymyalgia rheumatica and for monitoring patients with Hodgkin's disease. High values (>100 mm/h) have a 90% predictive value for serious disease including infections, collagen vascular disease or malignancy (particularly myeloma). A raised ESR is associated with marked rouleaux formation of red cells in the peripheral blood film (see Fig. 18.4). Changes in the ESR can be used to monitor the response to therapy.

Lower than expected readings occur in polycythaemia vera because of the high red cell concentration. Higher than expected values may occur in severe anaemia because of the low red cell concentration.

Plasma viscosity

In many laboratories measurement of ESR has been replaced by plasma viscosity measurement. Plasma viscosity is affected by the concentration of plasma proteins of large molecular size, especially those with pronounced axial asymmetry—fibrinogen and some immunoglobulins. Normal values at room temperature are usually in the range of 1.50–1.70 mPa/s. Lower levels are found in neonates

because of lower levels of proteins, particularly fibrinogen. Viscosity increases only slightly in the elderly as fibrinogen increases. There is no difference in values between men and women. Other advantages over the ESR test include independence from the effects of anaemia and results that are available within 15 min.

C-reactive protein

Phylogenetically CRP is a crude 'early' immunoglobulin which initiates the inflammatory reaction. CRP–antigen complexes can substitute for antibody in the fixation of C1q and trigger the complement cascade initiating the inflammatory response to antigens or tissue damage. Subsequent binding of C3b on the surface of microorganisms opsonizes them for phagocytosis.

After tissue injury, an increase in CRP, SAA protein and other acute phase reactants may be detected within 6–10 h. Increase in fibrinogen may not occur until 24–48 h following injury. Immunoassays of CRP are now widely used for early detection of acute inflammation or tissue injury and for the monitoring of remission (e.g. response of infection to an antibiotic).

Table 26.6 lists the advantages and disadvantages of the tests used to assess the acute phase response.

Bibliography

- Desnick R.J. and Schuchman E.H. (2003) Enzyme replacement and enhancement therapies: lessons from lysosomal disorders. *Nat Rev Genet* 3,954–66. Erratum in: (2003) *Nat Rev Genet* 4,157.
- Ganz T. (2003) Hepcidin, a key regulator of iron metabolism and mediator of anemia of inflammation. *Blood* 102,783–8.
- Ludwig H., Rai K., Blade J. *et al.* (2002) Management of disease-related anemia in patients with multiple myeloma or chronic lymphocytic leukemia: epoetin treatment recommendations. *Hematol J* 3,121–30.
- Spivak J.L. (2000) The blood in systemic disorders. *Lancet* 355,1707–12.
- Vellodi A. (2005) Lysosomal storage disorders. *Br J Haematol* 128,413–31.
- Weiss G. and Goodnough L.T. (2005) Anemia of chronic disease. *N Engl J Med* 352,1011–23.
- Wraith J.E. (2001) Advances in the treatment of lysosomal storage disease. *Dev Med Child Neurol* 43,639–46.

CHAPTER 27

27

Blood transfusion

Blood donor, 337

Red cell antigens and blood group antibodies, 337

Hazards of allogeneic blood transfusion, 341

Techniques in blood group serology, 343

Cross-matching and pre-transfusion tests, 343

Complications of blood transfusion, 344

Reduction of blood product use, 346

Blood components, 347

Preparations from human plasma, 349

Acute blood loss, 350

Bibliography, 350

Blood transfusion consists of the 'safe' transfer of blood components (Fig. 27.1) from a donor to a recipient.

Blood donor

This should be voluntary. The measures to protect donors are listed in Table 27.1.

Red cell antigens and blood group antibodies

Approximately 400 red blood cell group antigens have been described. The clinical significance of blood groups in blood transfusion is that individuals who lack a particular blood group antigen may produce antibodies reacting with that antigen which may




Fig. 27.1 The preparation of blood components from whole blood. FFP, fresh frozen plasma; SAGM, saline-adenine-glucose-mannitol. * Cryoprecipitate is mainly a source of fibrinogen. Cryosupernatant is used for plasma exchange in thrombotic thrombocytopenic purpura.

Table 27.1 Measures to protect the donor.

Age 17–70 years (maximum 60 at first donation)
Weight above 50 kg (7 st 12 lb)
Haemoglobin >13 g/dL for men, >12 g/dL for women
Minimum donation interval of 12 weeks (16 weeks advised) and three donations per year maximum
Pregnant and lactating women excluded because of high iron requirements
Exclusion of those with:
known cardiovascular disease, including hypertension
significant respiratory disorders
epilepsy and other CNS disorders
gastrointestinal disorders with impaired absorption
Insulin-dependent diabetes
Chronic renal disease
Ongoing medical investigation or clinical trials
Exclusion of any donor returning to occupations such as driving bus, plane or train, heavy machine or crane operator, mining, scaffolding, etc. because delayed faint would be dangerous

CNS, central nervous system.

lead to a transfusion reaction. The different blood group antigens vary greatly in their clinical significance with the ABO and Rh (formerly Rhesus) groups being the most important. Some other systems are listed in Table 27.2.

Blood group antibodies

Naturally occurring antibodies occur in the plasma of subjects who lack the corresponding antigen and

who have not been transfused or been pregnant. The most important are anti-A and anti-B. They are usually immunoglobulin M (IgM), and react optimally at cold temperatures (4°C) so, although reactive at 37°C, are called cold antibodies.

Immune antibodies develop in response to the introduction—by transfusion or by transplacental passage during pregnancy—of red cells possessing antigens that the subject lacks. These antibodies are commonly IgG, although some IgM antibodies may also develop—usually in the early phase of an immune response. Immune antibodies react optimally at 37°C (warm antibodies). Only IgG antibodies are capable of transplacental passage from mother to fetus. The most important immune antibody is the Rh antibody, anti-D.

ABO system

This consists of three allelic genes: A, B and O. The A and B genes control the synthesis of specific enzymes responsible for the addition of single carbohydrate residues (*N*-acetyl galactosamine for group A and *D*-galactose for group B) to a basic antigenic glycoprotein or glycolipid with a terminal sugar *L*-fucose on the red cell, known as the H substance (Fig. 27.2). The O gene is an amorph and does not transform the H substance. Although there are six possible genotypes, the absence of a specific anti-O prevents the serological recognition of more than four phenotypes (Table 27.3). The two major subgroups of A (A_1 and A_2) complicate the issue but are of minor clinical significance. A_2 cells react more

Table 27.2 Clinically important blood group systems.

Systems	Frequency of antibodies	Cause of haemolytic transfusion reaction	Cause of haemolytic disease of newborn
ABO	Almost universal	Yes (common)	Yes (usually mild)
Rh	Common	Yes (common)	Yes
Kell	Occasional	Yes (occasional)	Anaemia not haemolysis
Duffy	Occasional	Yes (occasional)	Yes (occasional)
Kidd	Occasional	Yes (occasional)	Yes (occasional)
Lutheran	Rare	Yes (rare)	No
Lewis	Occasional	Yes (rare)	No
P	Occasional	Yes (rare)	Yes (rare)
MN	Rare	Yes (rare)	Yes (rare)
Li	Rare	Unlikely	No




Fig. 27.2 Structure of ABO blood group antigens. Each consists of a chain of sugars attached to lipids or proteins which are an integral part of the cell membrane. The H antigen of the O blood group has a terminal fucose (fuc). The A antigen has an additional *N*-acetyl galactosamine (galnac), and the B antigen has an additional galactose (gal). glu, glucose.

Table 27.3 The ABO blood group system.

Phenotype	Genotype	Antigens	Naturally occurring antibodies	Frequency (UK) (%)
O	OO	O	Anti-A, anti-B	46
A	AA or AO	A	Anti-B	42
B	BB or BO	B	Anti-A	9
AB	AB	AB	None	3

weakly than A₁ cells with anti-A and patients who are A₂B can be wrongly grouped as B.

The A, B and H antigens are present on most body cells including white cells and platelets. In the 80% of the population who possess secretor genes, these antigens are also found in soluble form in secretions and body fluids (e.g. plasma, saliva, semen and sweat).


Naturally occurring antibodies to A and/or B antigens are found in the plasma of subjects whose red cells lack the corresponding antigen (Table 27.3; Fig. 27.3).

Rhesus system


The Rh blood group locus is composed of two related structural genes, *RhD* and *RhCE*, which encode the membrane proteins that carry the D, Cc and Ee

antigens. The *RhD* gene may be either present or absent, giving the Rh D+ or Rh D- phenotype, respectively. Alternative RNA splicing from the *RhCE* gene generates two proteins, which encode the C, c, E or e antigens (Fig. 27.4). A shortened nomenclature for Rh phenotype is commonly used (Table 27.4).

Rh antibodies rarely occur naturally; most are immune (i.e. they result from previous transfusion or pregnancy). Anti-D is responsible for most of the clinical problems associated with the system and a simple subdivision of subjects into Rh D+ and Rh D- using anti-D is sufficient for routine clinical purposes. Anti-C, anti-c, anti-E and anti-e are occasionally seen and may cause both transfusion reactions and haemolytic disease of the newborn. Anti-d does not exist. Rh haemolytic disease of the newborn is described in Chapter 28.



(a)



(b)

Fig. 27.3 (a) The ABO grouping in a group A patient. The red cells suspended in saline agglutinate in the presence of anti-A or anti-A + B (serum from a group O patient). (b) Routine grouping in a 96-well microplate. Positive reactions show as sharp agglutinates; in negative reactions the cells are dispersed. Rows 1–3, patient cells against antisera; rows 4–6, patient sera against known cells; rows 7–8, anti-D against patient cells.




Fig. 27.4 Molecular genetics of the rhesus blood group. The locus consists of two closely linked genes, *RhD* and *RhCcEe*. The *RhD* gene codes for a single protein which contains the RhD antigen whereas *RhCcEe* mRNA undergoes alternative splicing to three transcripts. One of these encodes the E or e antigen whereas the other two

(only one is shown) contain the C or c epitope. A polymorphism at position 226 of the *RhCcEe* gene determines the Ee antigen status whereas the C or c antigens are determined by a four amino acid allelic difference. Some individuals do not have an *RhD* gene and are therefore RhD–.

Other blood group systems

Other blood group systems are less frequently of clinical importance. Although naturally occurring antibodies of the P, Lewis and MN system are not

uncommon, they usually only react at low temperatures and hence are of no clinical consequence. Immune antibodies against antigens of these systems are detected infrequently. Many of the antigens are of low antigenicity and others (e.g. Kell),

Table 27.4 The most common Rh genotypes in the UK population.

CDE nomenclature	Short symbol	Caucasian frequency (%)	Rh D status
cde/cde	rr	15	Negative
CDe/cde	R ₁ r	31	Positive
CDe/CDe	R ₁ R ₁	16	Positive
cDE/cde	R ₂ r	13	Positive
CDe/cDE	R ₁ R ₂	13	Positive
cDE/cDE	R ₂ R ₂	3	Positive
Other genotypes		9	Positive (almost all)

although comparatively immunogenic, are of relatively low frequency and therefore provide few opportunities for isoimmunization except in multiply transfused patients.

Hazards of allogeneic blood transfusion

A large number of measures are taken to protect the recipient (Table 27.5).

Infection

Donor selection and testing of all donations are designed to prevent transmission of diseases (Table 27.6). The main risk is from viruses that have long incubation periods and especially those that are carried for many years by asymptomatic individuals. Some viruses that are transfusion transmissible show cell-associated latency and, if in white cells, can cause infection in the recipient after allogeneic transfusion. Live viruses causing acute infection can

Table 27.6 Microbiological testing in England and Wales.

- 1 Human immunodeficiency virus (HIV) 1 and 2
- 2 Hepatitis B virus (HBV)
- 3 Hepatitis C virus (HCV)
- 4 Human T-cell leukaemia viruses (HTLV)
- 5 Cytomegalovirus (CMV)—for immunosuppressed recipients
- 6 Malaria—antibody screening of potentially exposed donors
- 7 Chagas' disease—antibody screening of potentially exposed donors
- 8 Bacteria—all donations tested for antibody to syphilis17p

be transmitted in the pre-symptomatic viraemic phase if blood is collected during that short period.

Individual infections (Table 27.7)

Hepatitis

Donors with a history of hepatitis are deferred for 12 months. If there is a history of jaundice, they can be accepted if markers for HBV and HCV are negative.

Human immunodeficiency virus (HIV)

This can be transmitted by cells or plasma. Male homosexuals, bisexuals, intravenous drug users and prostitutes are excluded, as are their sexual partners and partners of haemophiliacs. Inhabitants of large areas of sub-Saharan Africa and South-East Asia where HIV infection is particularly common are also excluded. Rare transmission occurs when the donor is incubating the infection but is not yet positive for the antigen-antibody test used (window period transmission).

Table 27.5 Measures to protect recipient.

Donor selection
Donor deferral/exclusion
Stringent arm cleaning
Microbiological testing of donations (Table 27.6)
Immunohaematological testing of donations
Diversion of first 20–30 mL of blood collected
Leucodepletion of cellular products
Post-collection viral inactivation
Monitoring and testing for bacterial contamination
Pathogen inactivation
Safest possible sources of donor for plasma products

Table 27.7 Infections agents reported to have been transmitted by blood transfusion.

Viruses	
Hepatitis viruses	Hepatitis A virus (HAV) Hepatitis B virus (HBV) Hepatitis C virus (HCV) Hepatitis D virus (HDV) (requires coinfection with HBV)
Retroviruses	Human immunodeficiency virus (HIV) 1 + 2 (+ other subtypes)
Herpes viruses	Human T-cell leukaemia virus (HTLV) I + II Human cytomegalovirus (CMV) Epstein-Barr virus (EBV) Human herpesvirus 8 (HHV-8)
Parvoviruses	Parvovirus B19
Miscellaneous viruses	GBV-C—previously referred to as hepatitis G virus (HGV) Transfusion transmitted virus (TTV) West Nile virus
Bacteria	
Endogenous	<i>Treponema pallidum</i> (syphilis) <i>Borrelia burgdorferi</i> (Lyme disease) <i>Brucella melitensis</i> (brucellosis) <i>Yersinia enterocolitica</i> / <i>Salmonella</i> spp.
Exogenous	Environmental species—staphylococcal spp./pseudomonas/ <i>Serratia</i> spp.
Rickettsiae	<i>Rickettsia rickettsii</i> (Rocky Mountain spotted fever) <i>Coxiella burnetii</i> (Q fever)
Protozoa	
	<i>Plasmodium</i> spp. (malaria) <i>Trypanosoma cruzi</i> (Chagas' disease) <i>Toxoplasma gondii</i> (toxoplasmosis) <i>Babesia microti/divergens</i> (babesiosis) <i>Leishmania</i> spp. (leishmaniasis)
Prions	
	New variant Creutzfeldt-Jacob disease (nvCJD)

Human T-cell leukaemias viruses

Human T-cell leukaemias virus type I (HTLVI) is associated with adult T-cell leukaemia or tropical spastic paraparesis. Human T-cell leukaemia virus type II (HTLVII) has no known association with any clinical condition. Screening for both is mandatory in the UK despite the low prevalence, approximately 1 in 50 000 untested donors.

Cytomegalovirus

Post-infusion cytomegalovirus (CMV) infection is usually subclinical but may give an infectious mononucleosis syndrome. Immunosuppressed individuals are at risk of pneumonitis and a potentially fatal disease. These are premature babies (<1500 g),

bone marrow and other organ transplant recipients, and pregnant women (the fetus is at risk). For such recipients, CMV negative blood or blood components must be given.

Other infections

Syphilis is more likely to be transmitted by platelets (stored at room temperature) than blood (stored at 4°C). However, all donations are tested. Malarial parasites are viable in blood stored at 4°C, so in endemic areas all recipients are given antimalarial drugs. In non-endemic areas, donors are carefully vetted for travel to tropical areas and in some centres tests for malarial antibodies are performed. Chagas' disease is a significant problem with blood

transfusion in Latin America. **Bacterial infections** resulting from skin commensals are most frequently transmitted by platelets stored for more than 3 days.

Prions The risk of new variant Creutzfeldt–Jacob disease (nvCJD) is considered a threat to blood safety only in the UK. Plasma for fractionation and fresh frozen plasma for infants or children is obtained from the USA. It is unknown how many people could be infected with nvCJD. There are three reports of possible transmission by blood transfusion, so recipients of blood or blood components are now excluded as blood donors in the UK. No screening tests for prions are yet available.

Techniques in blood group serology

The most important technique is based on the agglutination of red blood cells. Saline agglutination is important in detecting IgM antibodies, usually at room temperature and 4°C (e.g. anti-A, anti-B; Fig. 27.3). Addition of colloid to the incubation or proteolytic enzyme treatment of red cells increases the sensitivity of the indirect antiglobulin test, as does low ionic strength saline (LISS). These latter methods can detect a range of IgG antibodies.

The antiglobulin (Coombs') test is a fundamental and widely used test in both blood group serology and general immunology. Antihuman globulin (AHG) is produced in animals following the injection of human globulin, purified complement or specific immunoglobulin (e.g. IgG, IgA or IgM). Monoclonal preparations are also now available. When AHG is added to human red cells coated with immunoglobulin or complement components, agglutination of the red cells indicates a positive test (Fig. 27.5).

The antiglobulin test may be either direct or indirect. The direct antiglobulin test (DAT) is used for detecting antibody or complement on the red cell surface where sensitization has occurred *in vivo*. The AHG reagent is added to washed red cells and agglutination indicates a positive test. A positive test occurs in haemolytic disease of the newborn, autoimmune or drug-induced immune haemolytic anaemia and haemolytic transfusion reactions.

The indirect antiglobulin test (IAT) is used to detect antibodies that have coated the red cells *in*




Fig. 27.5 The antiglobulin test for antibody or complement on the surface of red blood cells (RBC). The antihuman globulin (Coombs') reagent may be broad spectrum or specific for immunoglobulin G (IgG), IgM, IgA or complement (C3).

vitro. It is a two-stage procedure: the first step involves the incubation of test red cells with serum; in the second step, the red cells are washed and the AHG reagent is added. Agglutination implies that the original serum contained antibody which has coated the red cells *in vitro*. This test is used as part of the routine antibody screening of the recipient's serum prior to transfusion and for detecting blood group antibodies in a pregnant woman.

Most of the above methods were originally developed for tube techniques. These were replaced by 96-well microplates but most laboratories now use gel-based technology (Fig. 27.6).

Cross-matching and pre-transfusion tests

A number of steps are taken to ensure that patients receive compatible blood at the time of transfusion.

From the patient

- 1 The ABO and Rh blood group is determined.
- 2 Serum is screened for important antibodies by an indirect antiglobulin test on a large panel of antigenically-typed group O red cells.

If a red cell alloantibody is discovered, donor blood is selected lacking the relative antigen. The most likely are Rh D, C, c, E, e and K.




Fig. 27.6 Patient antibody screening using the microcolumn (gel) system: 10 tests with two controls (tube 11 is the positive control and tube 12 the negative control) are shown. The patient's serum is tested against screening cells with known red cell phenotype. Tubes 1, 3, 5–8 and 10 show positive results. The patient's serum contained anti-Fy^a. (Courtesy of Mr G. Hazlehurst)

Table 27.8 Techniques used in compatibility testing. Donor cells tested against recipient serum and agglutination detected visually or microscopically after mixing and incubation at the appropriate temperature.

For detecting clinically significant IgM antibodies
Saline 37°C

For detecting immune antibodies (mainly IgG)
Indirect antiglobulin test at 37°C
Low ionic strength saline at 37°C
Enzyme-treated red cells at 37°C

Ig, immunoglobulin.

From the donor

An appropriate ABO and Rh unit is selected. Donor (blood) testing is described on p. 341.

The cross-match

The techniques that may be used are described in Table 27.8.

Electronic cross-match

In this, a patient has group and antibody screen performed as two separate occasions. If both are negative and no blood has been transfused between the

test, ABO and Rh compatible blood is issued directly without wet testing.

Complications of blood transfusion

(Table 27.9)

Haemolytic transfusion reactions

Haemolytic transfusion reactions may be immediate or delayed. Immediate life-threatening reactions associated with massive intravascular haemolysis are the result of complement-activating antibodies of IgM or IgG classes, usually with ABO specificity. Reactions associated with extravascular haemolysis (e.g. immune antibodies of the Rh system which are unable to activate complement) are generally less severe but may still be life-threatening. The cells become coated with IgG and are removed in the reticuloendothelial system. In mild cases, the only signs of a transfusion reaction may be a progressive unexplained anaemia with or without jaundice. In some cases where the pre-transfusion level of an antibody was too low to be detected in a cross-match, a patient may be reimmunized by transfusion of incompatible red cells and this will lead to a delayed transfusion reaction with accelerated clearance of the red cells. There may be rapid appearance of anaemia with mild jaundice.

Clinical features of a major haemolytic transfusion reaction

Haemolytic shock phase This may occur after only a few millilitres of blood have been transfused or up to 1–2 h after the end of the transfusion. Clinical features include urticaria, pain in the lumbar region, flushing, headache, precordial pain, shortness of breath, vomiting, rigours, pyrexia and a fall in blood pressure. If the patient is anaesthetized this shock phase is masked. There is increasing evidence of red cell destruction and haemoglobinuria, jaundice and disseminated intravascular coagulation (DIC) may occur. Moderate leucocytosis (e.g. $15\text{--}20 \times 10^9/\text{L}$) is usual.

The oliguric phase In some patients with a haemolytic reaction there is renal tubular necrosis with acute renal failure.

Table 27.9 Complications of blood transfusion

Early	Late
Haemolytic reactions: immediate or delayed	Transmission of infection (Table 27.7)
Reactions caused by infected blood	Transfusional iron overload
Allergic reactions to white cells, platelets or proteins	Immune sensitization e.g. to red cells, platelets or Rh D antigen
Pyrogenic reactions (to plasma proteins or caused by HLA antibodies)	Transfusion-associated graft-versus-host disease
Circulatory overload	
Bacterial contamination	
Air embolism	
Thrombophlebitis	
Citrate toxicity	
Hyperkalaemia	
Clotting abnormalities (after massive transfusion)	
TRALI	
Post-transfusion purpura	

CMV, cytomegalovirus; HIV, human immunodeficiency virus; HLA, human leucocyte antigen; TRALI, transfusion-related acute lung injury.

Diuretic phase Fluid and electrolyte imbalance may occur during the recovery from acute renal failure.

Investigation of an immediate transfusion reaction

If a patient develops features suggesting a severe transfusion reaction the transfusion should be stopped and investigations for blood group incompatibility and bacterial contamination of the blood must be initiated.

1 Most severe reactions occur because of clerical errors in the handling of donor or recipient blood specimens. Therefore it must be established that the identity of the recipient (from the patient's wristband) is the same as that on the compatibility label and that this corresponds with the actual unit being transfused.

2 The unit of donor blood and post-transfusion samples of the patient's blood should be sent to the laboratory who will:

- repeat the group on pre- and post-transfusion samples and on the donor blood, and repeat the cross-match;
- perform a direct antiglobulin test on the post-transfusion sample;
- check the plasma for haemoglobinaemia;

(d) perform tests for DIC; and

(e) examine the donor sample directly for evidence of gross bacterial contamination and set up blood cultures from it at 20 and 37°C. If the clinical picture is suggestive of bacterial infection blood cultures must be taken from the patient and broad spectrum intravenous antibiotics started.

3 A post-transfusion sample of urine must be examined for haemoglobinuria.

4 Further samples of blood are taken 6 h and/or 24 h after transfusion for a blood count and bilirubin, free haemoglobin and methaemalbumin estimations.

5 In the absence of positive findings, the patient's serum is examined 5–10 days later for red cell or white cell antibodies.

Management of patients with major haemolysis

The principal object of initial therapy is to maintain the blood pressure and renal perfusion. Intravenous dextran, plasma or saline and frusemide are sometimes needed. Hydrocortisone 100 mg intravenously and an antihistamine may help to alleviate shock. In the event of severe shock, support with intravenous adrenaline 1 : 10 000 in small incremental doses may be required. Further compatible

transfusions may be required in severely affected patients. If acute renal failure occurs this is managed in the usual way, if necessary with dialysis until recovery occurs.

Other transfusion reactions

Febrile reactions because of white cell antibodies Human leucocyte antigen (HLA) antibodies (see below and Chapter 21) are usually the result of sensitization by pregnancy or a previous transfusion. They produce rigors, pyrexia and, in severe cases, pulmonary infiltrates. They are minimized by giving leucocyte-depleted (i.e. filtered) packed cells (see below).

Febrile or non-febrile non-haemolytic allergic reactions These are usually caused by hypersensitivity to donor plasma proteins and if severe can result in anaphylactic shock. The clinical features are urticaria, pyrexia and, in severe cases, dyspnoea, facial oedema and rigors. Immediate treatment is with antihistamines and hydrocortisone. Adrenaline is also useful. Washed red cells or frozen red cells may be needed for further transfusions if the majority of plasma-removed blood (e.g. saline, adenine, glucose, mannitol (SAGM) blood) causes reactions.

Post-transfusion circulatory overload The management is that of cardiac failure. These reactions are prevented by a slow transfusion of packed red cells or of the blood component required, accompanied by diuretic therapy.

Transfusion of bacterially contaminated blood This is very rare but may be serious. It can present with circulatory collapse.

Graft-versus-host disease This may occur when live lymphocytes are transfused to an immunocompromised patient. It is prevented by irradiation of the blood product. It is uniformly fatal.

Transfusion related acute lung injury (TRALI) This presents as pulmonary infiltrates with chest symptoms depending on severity. It is caused by positive transfer of leucoagglutins in donor plasma causing endothelial and epithelial injury. Most

of the donors are multiparous women. Management is supportive.

Post-transfusion purpura This is a rare problem of severe thrombocytopenia 7–10 days after transfusion of a platelet-containing product, usually red cells. It is caused by an antibody in the recipient (previously transfused or pregnant) anti HPA-1a against a platelet-specific antigen HPA-1a (PI^A). Both transfused and recipient platelets are destroyed by the immune complexes. It is usually self-limiting but immunoglobulin or plasma exchange may be needed.

Viral transmission Post-transfusion hepatitis may be caused by one of the hepatitis viruses, although CMV and Epstein–Barr virus (EBV) have also been implicated. Post-transfusion hepatitis and HIV infection is seen rarely now because of routine screening of all blood donations.

Other infections Toxoplasmosis, malaria and syphilis may be transmitted by blood transfusion. Transfusion-transmitted nvCJD has probably occurred in three cases in the UK.

Post-transfusional iron overload Repeated red cell transfusions over many years, in the absence of blood loss, cause deposition of iron initially in reticuloendothelial tissue at the rate of 200–250 mg/unit of red cells. After 50 units in adults, and lesser amounts in children, the liver, myocardium and endocrine glands are damaged with clinical consequences. This is a major problem in thalassaemia major and other severe chronic refractory anaemias (Chapter 6).

Reduction of blood product use

In light of transfusion risks, and limited resources, appropriate use of blood component is of ever increasing importance.

Preoperative correction of anaemia (particularly iron deficiency) and cessation of anti-platelet therapies (e.g. aspirin) where possible, together with lower trigger levels for red cell transfusions (7–8 g/dL in most surgical patients) can all help to reduce blood use.

In surgery the use of alternative fluid replacement, intraoperative or post-operative cell salvage, biological alternatives (e.g. erythropoietin, recombinant clotting factors, recombinant activated clotting factor VII (VIIa) or fibrin glue) all may help.

Blood components

A blood donation is taken by an aseptic technique into plastic bags containing an appropriate amount of anticoagulant—usually citrate, phosphate, dextrose (CPD). The citrate anticoagulates the blood by combining with the blood calcium. Three components are made by initial centrifugation of whole blood: red cells, buffy coat and plasma (Fig. 27.1).

Red cells are stored at 4–6°C for up to 35 days, depending on the preservative. After the first 48 h there is a slow progressive K⁺ loss from the red cells into the plasma. In cases where infusion of K⁺ could be dangerous, fresh blood should be used (e.g. for exchange transfusion in haemolytic disease of the newborn). During red cell storage there is a fall in 2,3-diphosphoglycerate (2,3-DPG) but after transfusion 2,3-DPG levels return to normal within 24 h. Optimum additive solutions have been developed to increase the shelf life of plasma-depleted red cells by maintaining both adenosine triphosphate (ATP) and 2,3-DPG levels.

Platelets and plasma may also be collected by apheresis.

Leucodepletion

In many countries, including Britain, blood products are now routinely filtered to remove the majority of white cells, a process known as leucodepletion. This is usually performed soon after collection and prior to processing and is more effective than filtration of blood at the bedside. A blood component is defined as leucocyte depleted if there are less than 5×10^6 /L white cells present.

Leucodepletion reduces the incidence of febrile transfusion reactions and HLA alloimmunization. It is effective at preventing transmission of CMV infection and in addition should reduce the theoretical possibility of transmission of nvCJD in countries where this has been reported.

Red cells

Packed (plasma-depleted) red cells are the treatment of choice for most transfusions (Fig. 27.7a). In older subjects, a diuretic is often given simultaneously and the infusion should be sufficiently slow to avoid circulatory overload. Iron chelation therapy should be considered with patients on a regular transfusion programme to avoid iron overload.

Recombinant erythropoietin is widely used to reduce transfusion requirements (e.g. in patients on dialysis, cancer patients and myelodysplasia). Factor VIIa can reduce transfusion need in patients with major haemorrhage (e.g. at surgery or after trauma).

Red cell substitutes are under development but have not yet proven clinically valuable. These synthetic oxygen-carrying substitutes are often fluorinated hydrocarbons and stromal-free pyridoxylated and polymerized haemoglobin solutions.

Autologous donation and transfusion

Anxiety over acquired immune deficiency syndrome (AIDS) and other infections has increased the demand for autotransfusion. There are three ways of administering an autologous transfusion:

- 1 *Predeposit* Blood is taken from the potential recipient in the weeks immediately prior to elective surgery.
- 2 *Haemodilution* Blood is removed immediately prior to surgery once the patient has been anaesthetized and then reinfused at the end of the operation.
- 3 *Salvage* Blood lost during the operation is collected during heavy blood loss and then reinfused.

Autotransfusion is the safest form of transfusion with regard to transmission of viral disease though it has a higher risk of bacterial contamination and of clerical errors. The individual involved must be fit enough to donate blood and the predicted operative replacement transfusion should be 2–4 units. Larger replacement transfusions would require blood to be collected over a longer period and red cells stored in the frozen state, which is both labour intensive and expensive. The high cost and initial restriction of its use to patients undergoing elective surgery means that it will benefit only a minor proportion of the total number of blood recipients.



(a)



(b)



(c)

Fig. 27.7 Blood components: (a) plasma-depleted red cells; (b) platelets; and (c) fresh frozen plasma.

Preoperative autotransfusion is largely reserved for those patients with multiple antibodies.

Granulocyte concentrates

These are prepared as buffy coats or on blood cell separators from normal healthy donors or from patients with chronic myeloid leukaemia. They have been used in patients with severe neutropenia ($<0.5 \times 10^9/\text{L}$) who are not responding to antibiotic therapy but it is not usually possible to give sufficient amounts. They may transmit CMV infection and must be irradiated to eliminate the risk of causing GVHD.

Platelet concentrates

These are harvested by cell separators or from individual donor units of blood (Fig. 27.7b). They are stored at room temperature. Platelet transfusion is used in patients who are thrombocytopenic or have disordered platelet function and who are actively bleeding (therapeutic use) or are at serious risk of bleeding (prophylactic use).

For prophylaxis, the platelet count should be kept above $5\text{--}10 \times 10^9/\text{L}$ unless there are additional risk factors such as sepsis, drug use or coagulation disorders for which the threshold should be higher. For invasive procedures (e.g. liver biopsy or lumbar puncture) the platelet count should be raised to above $50 \times 10^9/\text{L}$. For brain or eye surgery the count should be $>100 \times 10^9/\text{L}$.

Therapeutic use is indicated in bleeding associated with platelet disorders. In massive haemorrhage the count should be kept above $50 \times 10^9/\text{L}$.

Platelet transfusions should be avoided in autoimmune thrombocytopenic purpura unless there is serious haemorrhage. They are contraindicated in heparin-induced thrombocytopenia, thrombotic thrombocytopenic purpura and haemolytic uraemic syndrome (p. 284).

Refractoriness to platelet transfusions is defined by a poor platelet increment post-transfusion ($<7.5 \times 10^9/\text{L}$ at 1 h or $<4.5 \times 10^9/\text{L}$ at 24 h). The causes are either immunological (mostly HLA alloimmunization) or non-immunological (sepsis, hypersplenism, DIC, drugs). Platelets express HLA class I (but not class II) antigens and HLA-matched

or cross-match-compatible platelets are needed for patients with HLA antibodies.

Preparations from human plasma

Fresh frozen plasma (FFP)

Rapidly frozen plasma separated from fresh blood is stored at less than -30°C . Frozen plasma is usually prepared from single donor units though pooled products are also available. Its main use is for the replacement of coagulation factors (e.g. when specific concentrates are unavailable) or after massive transfusions, in liver disease and DIC, after cardiopulmonary bypass surgery, to reverse a warfarin effect, and in thrombotic thrombocytopenic purpura (see p. 284). Virally inactivated forms of FFP are now available.

Human albumin solution (4.5%)

It is a useful plasma volume expander when a sustained osmotic effect is required prior to the administration of blood, but it should not be given in excess. It is also used for fluid replacement in patients undergoing plasmapheresis and sometimes for fluid replacement in selected patients with hypoalbuminaemia.

Human albumin solution (20%) (salt-poor albumin)

It may be used in severe hypoalbuminaemia when it is necessary to use a product with minimal electrolyte content. Principal indications for its use are patients with nephrotic syndrome or liver failure.

Cryoprecipitate

This is obtained by thawing FFP at 4°C and contains concentrated factor VIII and fibrinogen. It is stored at less than -30°C or, if lyophilized, at $4\text{--}6^\circ\text{C}$, and was used widely as replacement therapy in haemophilia A and von Willebrand disease before more purified preparations of factor VIII became available. Its main use is in fibrinogen replacement in disseminated intravascular coagulation (DIC) or massive transfusion or hepatic failure.

Freeze-dried factor VIII concentrates

These are also used for treating haemophilia A or von Willebrand disease. The small volume makes them ideal for children, surgical cases, patients at risk from circulatory overload and for those on home treatment. Their use is declining as recombinant forms of factor VIII become widely available.

Freeze-dried factor IX–prothrombin complex concentrates

A number of preparations are available that contain variable amounts of factors II, VII, IX and X. They are mainly used for treating factor IX deficiency (Christmas disease) but are also used in patients with liver disease or in haemorrhage following overdose with oral anti-coagulants or in patients with factor VIII inhibitors. There is a risk of thrombosis.

Protein C concentrate

This is used in severe sepsis with disseminated intravascular coagulation (e.g. meningococcal septicaemia) to reduce thrombosis resulting from depletion of protein C.

Immunoglobulin

Pooled immunoglobulin is a valuable source of antibodies against common viruses. It is used in hypogammaglobulinaemia for protection against viral and bacterial disease. It may also be used in immune thrombocytopenia and other acquired immune disorders (e.g. post-transfusion purpura or alloimmune neonatal thrombocytopenia).

Specific immunoglobulin

This may be obtained from donors with high titres of antibody (e.g. anti-RhD, antihepatitis B, anti-herpes zoster or antirubella).

Acute blood loss

After a single episode of blood loss, there is initial vasoconstriction with a reduction in total blood volume. The plasma volume rapidly expands and

the haemoglobin and packed cell volume fall and there is a rise in neutrophils and platelets. The reticulocyte response begins on the second or third day and lasts 8–10 days. The haemoglobin begins to rise by about the seventh day but, if iron stores have become depleted, the haemoglobin may not rise subsequently to normal. Clinical assessment is needed to gauge whether blood transfusion is needed. This is usually unnecessary in adults at losses less than 500 mL unless haemorrhage is continuing and may not be needed with losses of up to 1.5 L. Blood transfusion is not without risks and should not be undertaken lightly. The problems of massive blood loss and massive transfusion are considered on p. 301.

Bibliography

- American Association of Blood Banks (2002) *American Association of Blood Banks Technical Manual*, 14th edn. AABB, Bethesda, MD.
- Ashford P., Gozzard D., Jones J. *et al.* for the British Committee for Standards in Haematology, Blood Transfusion Task Force. (2000) Guidelines for blood bank computing. *Transfus Med* 10,307–14.
- Barbara J.A.J. (2002) Transfusion transmitted infections. Transfusion alternatives. *Transfus Med* 4,42–7.
- Blumberg N., Francis C., Heal J. (eds) (2006) Clinical, use of plasma and plasma fractions. *Clin Haematol* 19,1–258.
- British Committee for Standards in Haematology, Blood Transfusion Task Force (1996) Guidelines for pretransfusion compatibility procedures in blood transfusion laboratories. *Transfus Med* 6,273–83.
- British Committee for Standards in Haematology, Blood Transfusion Task Force (1998) Guidelines on the clinical use of leucocyte-depleted blood components. *Transfus Med* 8,59–72.
- British Committee for Standards in Haematology, Blood Transfusion Task Force (2003) Guidelines for the use of platelet transfusions. *Br J Haematol* 122,10–23.
- British Committee for Standards in Haematology, Blood Transfusion Task Force (2004) Guidelines for the use of fresh-frozen plasma, cryoprecipitate and cryosupernatant. *Br J Haematol* 126,11–28.
- British Committee for Standards in Haematology Guidelines (1999) The administration of blood and blood components and the management of transfused patients. *Transfus Med* 9,227–38.
- Chang T.M.S. (2000) Red blood cell substitutes. *Clin Haematol* 13,651–68.
- Daniels G. (2002) *Human Blood Groups*, 2nd edn. Blackwell Science, Oxford.

- Daniels G., Poole J., de Silva M. *et al.* (2002) The clinical significance of blood group antibodies. *Transfus Med* 12,287–95.
- Gibson B.E., Todd A., Roberts I. *et al.* for the British Committee for Standards in Haematology Transfusion Task Force: Writing Group (2004) Transfusion guidelines for neonates and older children. *Br J Haematol* 124,433–53.
- Llewelyn C.A., Hewitt P.E., Knight R.S. *et al.* (2004) Possible transmission of variant Creutzfeldt–Jakob disease by blood transfusion. *Lancet* 363,417–21.
- Klein H. and Anstee D. (2005) *Mollison's Blood Transfusion in Clinical Medicine*, 11th edn. Blackwell Publishing, Oxford.
- Murphy M.F. and Pamphilon D.H. (eds) (2005) *Practical Transfusion Medicine*, 2nd edn. Blackwell Publishing, Oxford.
- Murphy M.F., Wallington T.B., Kelsey P. *et al.* for the British Committee for Standards in Haematology, Blood Transfusion Task Force (2001) Guidelines for the clinical use of red cell transfusions. *Br J Haematol* 113,24–31.
- National Institute for Clinical Excellence (2003) Full guidance on the use of antenatal anti-D prophylaxis for RhD–negative women (<http://www.nice.org.uk>).
- Roberts H.R., Monroe D.M., White G.C. (2004) The use of recombinant factor VIIa in the treatment of bleeding disorders. *Blood* 104,3858–64.
- Stainsky D., Cohen H., Jones H. *et al.* (2004) *Serious Hazards of Blood Transfusions Annual Report 2003* (<http://www.shotuk.org/SHOT%20Report%202003.pdf>).
- UK Blood Transfusion Services (2002) *Guidelines for the Blood Transfusion Services in the United Kingdom*, 6th edn. The Stationery Office of London (or <http://www.transfusionguidelines.org.uk>).
- Webert K.E. and Blajchman M.A. (2003) Transfusion-related acute lung injury. *Transfus Med Rev* 17,252–62.

CHAPTER 28

28

Pregnancy and neonatal haematology

Haematology of pregnancy, 352
Neonatal haematology, 354

Haemolytic disease of the newborn, 356
Bibliography, 358

Haematology of pregnancy

Pregnancy places extreme stresses on the haematological system and an understanding of the physiological changes that result is obligatory in order to interpret any need for therapeutic intervention.

Physiological anaemia

Physiological anaemia is the term often used to describe the fall in haemoglobin (Hb) concentration that occurs during normal pregnancy (Fig. 28.1). Blood plasma volume increases by approximately 1250 mL, or 45%, above normal by the end of gestation and although the red cell mass itself increases

by some 25% this still leads to a fall in Hb concentration. Values below 10 g/dL are probably abnormal and require investigation.

Iron deficiency anaemia

Up to 600 mg iron is required for the increase in red cell mass and a further 300 mg for the fetus. Despite an increase in iron absorption, few women avoid depletion of iron reserves by the end of pregnancy.

In uncomplicated pregnancy, the mean corpuscular volume (MCV) typically rises by approximately 4 fL. A fall in red cell MCV is the earliest sign of iron deficiency. Later, the mean corpuscular haemoglobin




Fig. 28.1 Haematological changes during pregnancy.




Fig. 28.2 Causes of thrombocytopenia during pregnancy.

(MCH) falls and finally anaemia results. Early iron deficiency is likely if the serum ferritin is below $15 \mu\text{g/L}$ together with serum iron $<10 \mu\text{mol/L}$ and should be treated with oral iron supplements. The use of routine iron supplementation in pregnancy is debated but iron is probably better avoided until the Hb falls below 10 g/dL or MCV below 82 fL in the third trimester.

Folate deficiency

Folate requirements are increased approximately twofold in pregnancy and serum folate levels fall to approximately half the normal range with a less dramatic fall in red cell folate. In some parts of the world, megaloblastic anaemia during pregnancy is common because of a combination of poor diet and exaggerated folate requirements. Given the protective effect of folate against neural tube defects, folic acid $400 \mu\text{g/day}$ should be taken periconceptually and throughout pregnancy. Food fortification with folate is now being practised in many countries. Vitamin B_{12} deficiency is rare during pregnancy although serum vitamin B_{12} levels fall to below normal in 20–30% of pregnancies and low values are sometimes the cause of diagnostic confusion.

Thrombocytopenia

The platelet count typically falls by approximately 10% in an uncomplicated pregnancy. In approxi-

mately 7% of women this fall is more severe and can result in thrombocytopenia (platelet count $<140 \times 10^9/\text{L}$). In over 75% of cases this is mild and of unknown cause, a condition referred to as *incidental thrombocytopenia of pregnancy*. Approximately 21% of cases are secondary to a hypertensive disorder and 4% are associated with immune thrombocytopenic purpura (ITP; Fig. 28.2).

Incidental thrombocytopenia of pregnancy This is a diagnosis of exclusion and is usually detected at the time of delivery. The platelet count is always $>70 \times 10^9/\text{L}$ and recovers within 6 weeks. No treatment is required and the infant is not affected.

Thrombocytopenia of hypertensive disorders This is variable in severity but the platelet count rarely falls to $<40 \times 10^9/\text{L}$. It is more severe when associated with pre-eclampsia and if severe the primary treatment is as rapid delivery as possible. The platelet count falls for a day or two after delivery and then recovers rapidly. The HELLP syndrome (haemolysis, elevated liver enzymes and low platelets) is a subtype of this category.

Idiopathic thrombocytopenic purpura (p. 281) In pregnancy, ITP represents a particular problem, both to the mother and to the fetus, as the antibody crosses the placenta and the fetus may become severely thrombocytopenic.

Like all adults, pregnant women with ITP and

platelet counts $>50 \times 10^9/L$ do not usually need treatment. Treatment is required for women with platelet counts $<10 \times 10^9/L$ and for those with platelet counts of $10\text{--}30 \times 10^9/L$ who are in their second or third trimester or who are bleeding. Treatment is with steroids, intravenous immunoglobulin G (IgG) and splenectomy as appropriate.

At delivery, umbilical vein blood sampling or fetal scalp vein sampling to measure the fetal platelet count may be offered although their exact role is unclear. In general, caesarean section is not indicated when the maternal platelet count is $>50 \times 10^9/L$ unless the fetal platelet count is known to be $<20 \times 10^9/L$. Platelet transfusion may be given to mothers in labour with very low platelet counts or who are actively bleeding.

Newborns of mothers with ITP should have a blood count measured for the first 4 days of life as the platelet count may progressively drop. A count greater than $50 \times 10^9/L$ is reassuring. Cerebral ultrasounds may be performed to look for intracranial haemorrhage (ICH). In newborns without evidence of ICH, treatment with intravenous IgG is appropriate if the infant's platelet count is $<20 \times 10^9/L$. Neonates with thrombocytopenia and ICH should be treated with steroids and intravenous IgG therapy.

Haemostasis and thrombosis

Pregnancy leads to a hypercoagulable state with consequent increased risks of thromboembolism and disseminated intravascular coagulation (DIC; p. 298). There is an increase in plasma factors VII, VIII, X and fibrinogen and fibrinolysis is suppressed. These changes last for up to 2 months into the puerperal period and the incidence of thrombosis during this period is increased. There is an association between thrombophilic conditions in the mother and with recurrent fetal loss. This is presumed to result from placental thrombosis and infarction.

Treatment of thrombosis

Warfarin has no role in management. It crosses the placenta and in addition is associated with embryopathy, especially between 6 and 12 weeks' gestation. Heparin does not cross the placenta but a significant side-effect of prolonged use is maternal

osteoporosis. Low molecular weight heparin is now the treatment of choice because it can be given once daily and is less likely to cause osteoporosis.

Neonatal haematology

Normal blood count

The cord blood Hb varies between approximately 16.5 and 17 g/dL and is influenced by the timing of cord clamping (Fig. 28.3). The reticulocyte count is initially high (2–6%) but falls to below 0.5% at 1 week as erythropoiesis is suppressed in response to the marked increase in the oxygenation of tissues after birth. This is associated with a progressive fall in Hb to approximately 10–11 g/dL at 8 weeks from which point it recovers to 12.5 g/dL at around 6 months. In the blood film, nucleated red cells will be seen for the first 4 days and for up to 1 week in preterm infants. Numbers are increased in cases of hypoxia, haemorrhage or haemolytic disease of the newborn (HDN). MCV averages 119 fL but falls to adult levels by around 9 weeks. By 1 year, the MCV has fallen to around 70 fL and rises throughout childhood again to reach adult levels at puberty. Preterm infants have a more dramatic fall in Hb to 7–9 g/dL at 8 weeks and are more prone to iron and folate deficiency in the first few months of life. Neutrophils are initially high at birth and fall to plateau at 4 days—from this point on the lymphocyte count is higher than neutrophils throughout childhood.




Fig. 28.3 Typical profile of the blood count in the neonatal period.




Fig. 28.4 The investigation of neonatal anaemia. * The DAT test may be negative in HDN, e.g. due to ABO incompatibility. DAT, direct antiglobulin test; HDN, haemolytic disease of the newborn; MCV, mean corpuscular volume.

Anaemia in the neonate

This should be considered for Hb <14 g/dL at birth. The clinical significance of anaemia is compounded by the high (70–80%) levels of HbF at birth, as this is less effective than HbA at releasing oxygen to the tissues (p. 17). Causes include the following (Fig. 28.4):

- 1 *Haemorrhage* Fetomaternal, twin–twin, cord, internal, placenta.

- 2 *Increased destruction* Haemolysis (immune or non-immune) or infection.

- 3 *Decreased production* Congenital red cell aplasia, infection (e.g. parvovirus). Anti-Kell causes allo-immune anaemia of the fetus and newborn with decreased erythropoiesis.

Generally, anaemia at birth is usually secondary to immune haemolysis or haemorrhage; non-immune causes of haemolysis appear within 24 h. Impaired red cell production is usually not apparent for at least 3 weeks. Haemolysis is often associated with severe jaundice and the causes include HDN, autoimmune haemolytic anaemia (AIHA) in

the mother and congenital disorders of the red cell membrane or metabolism.

Red cell transfusion may be needed for symptomatic anaemia with Hb <10.5 g/dL or a higher threshold if there is severe cardiac or respiratory disease.

Anaemia of prematurity

Premature infants have a more marked fall in Hb after birth and this is termed *physiological anaemia of prematurity*. Features include a slowly falling Hb, normal blood film and reticulocytopenia. It can be minimized by ensuring adequate iron and folate replacement and limiting phlebotomy. Erythropoietin is used in some centres.

Neonatal polycythaemia

This is defined as a venous haematocrit over 0.65 and can occur with twin–twin transfusion, intra-uterine growth restriction and maternal hypertension

or diabetes. If symptoms are present it should be treated with partial exchange transfusion using a crystalloid solution.

Fetomaternal alloimmune thrombocytopenia

Fetomaternal alloimmune thrombocytopenia (FMAIT) results from an immunological process similar to that which causes HDN. Fetal platelets that possess a paternally inherited antigen (HPA-1a in 80%; HPA-5b in 15%) that is not present on maternal platelets can sensitize the mother to make antibodies that cross the placenta, coat the platelets which are then destroyed by the reticuloendothelial system and lead to serious bleeding, including intracranial haemorrhage. Alloimmune thrombocytopenia differs from HDN in that 50% of cases occur in the first pregnancy. Its incidence is approximately 1 in 1000–5000 births.

Thrombocytopenia can lead to serious, sometimes fatal, bleeding *in utero* or after birth. Treatment is unsatisfactory. Severe postnatal cases may be treated with a platelet transfusion that is negative for the relevant antigen. Antenatal treatment may be either maternal intravenous immunoglobulin or fetal transfusion with HPA-compatible platelets.

Coagulation

Standard tests need to be interpreted with caution in the neonate. The activated partial thromboplastin time (APTT) and prothrombin time (PT) are prolonged because of reduced levels of the vitamin K-dependent factors II, VII, IX and X, and return to normal at around 6 months. The thrombin time (TT) is comparable with adult values. Neonates have an increased risk of thrombosis. This is a result of physiologically low levels of inhibitors of coagulation and the use of indwelling vascular catheters. Antithrombin (AT) and protein C levels are approximately 60% of normal for the first 3 months. Homozygous protein C deficiency is associated with fulminant purpura fulminans in early life. Therapeutic protein C concentrates are now available. Homozygous AT deficiency usually presents later in childhood but arterial and venous thrombosis may also occur in the neonate.

Haemorrhagic disease of the newborn is discussed on page 297.

Haemolytic disease of the newborn

HDN is the result of *red cell alloimmunization* in which IgG antibodies pass from the maternal circulation across the placenta into the circulation of the fetus where they react with fetal red cells and lead to their destruction. Anti-D antibody is responsible for most cases of severe HDN although anti-c, anti-E, anti-K and a wide range of other antibodies are found in occasional cases (see Table 27.2). Although antibodies against the ABO blood group system are most frequent cause of HDN this is usually mild. Within the UK, approximately 500 fetuses develop haemolytic disease each year and approximately 30 of these cases are fatal.

Rh HDN

When an Rh D-negative (p. 339) woman has a pregnancy with an Rh D-positive fetus, Rh D-positive fetal red cells cross into the maternal circulation (especially at parturition and during the third trimester) and sensitize the mother to form anti-D. The mother could also be sensitized by a previous miscarriage, amniocentesis or other trauma to the placenta or by blood transfusion. Anti-D crosses the placenta to the fetus during the next pregnancy with an Rh D-positive fetus, coats the fetal red cells and results in reticuloendothelial destruction of these cells, causing anaemia and jaundice. If the father is heterozygous for D antigen, there is a 50% probability that the fetus will be D-positive. The fetal Rh D genotype can be established by polymerase chain reaction (PCR) analysis for the presence of Rh D in a maternal blood sample.

The main aim of management is to *prevent* anti-D antibody formation in Rh D-negative mothers. This can be achieved by the administration of small amounts of anti-D antibody which 'mop up' and destroy Rh D-positive fetal red cells before they can sensitize the immune system of the mother to produce anti-D.

Prevention of Rh immunization

At the time of booking, all pregnant women should have their ABO and Rh group determined and serum screened for antibodies at least twice during the pregnancy. All non-sensitized Rh D-negative women should be given at least 500 units (100 µg) of

anti-D at 28 and 34 weeks' gestation to reduce the risk of sensitization from fetomaternal haemorrhage. Fetal Rh D typing from DNA in maternal blood can be used before 28 weeks. If the fetus is Rh D-negative, no further anti-D prophylaxis is needed. In addition, at birth the babies of Rh D-negative women who do not have antibodies must have their cord blood grouped for ABO and Rh. If the baby's blood is Rh D-negative, the mother will require no further treatment. If the baby is Rh D-positive, prophylactic anti-D should be administered at a minimum dose of 500 units intramuscularly within 72 h of delivery. A *Kleihauer test* is performed. If the Kleihauer is positive many centres will perform flow cytometry for a more accurate estimate of the volume of fetomaternal haemorrhage (FMH). This uses differential staining to estimate the number of fetal cells in the maternal circulation (Fig. 28.5). The chance of developing antibodies is related to the number of fetal cells found. The dose of anti-D is increased if there is greater than 4 mL transplacental haemorrhage. Anti-D IgG (125 units) is given for each 1 mL of FMH greater than 4 mL.

Sensitizing episodes during pregnancy Anti-D IgG should be given to Rh D-negative women who have potentially sensitizing episodes during pregnancy: 250 units is given if the event occurs up to week 20 of gestation and 500 units thereafter, followed by a Kleihauer test. Potentially sensitizing events include therapeutic termination of pregnancy, spontaneous

miscarriage after 12 weeks' gestation, ectopic pregnancy and invasive antenatal diagnostic procedures.

Treatment of established anti-D sensitization

If anti-D antibodies are detected during pregnancy they should be quantified at regular intervals. The strength of anti-D present in maternal serum is related to the clinical severity of HDN but this is also affected by such factors as the IgG subclass, rate of rise of antibody and past history. The development of haemolytic disease in the fetus can be assessed by velocimetry of the fetal middle cerebral artery by Doppler ultrasonography as increased velocities correlate with fetal anaemia (Fig. 28.6). If anaemia is detected, fetal blood sampling and intrauterine transfusion of irradiated Rh D-negative packed red cells may be indicated.

Clinical features of HDN

1 Severe disease Intrauterine death from hydrops fetalis (Fig. 28.7a).

2 Moderate disease The baby is born with anaemia and jaundice and may show pallor, tachycardia, oedema and hepatosplenomegaly. If the unconjugated bilirubin is not controlled and reaches levels exceeding 250 $\mu\text{mol/L}$, bile pigment deposition in the basal ganglia may lead to *kernicterus*—central nervous system damage with generalized spasticity and possible subsequent mental deficiency, deafness and epilepsy. This problem becomes acute after birth as maternal clearance of fetal bilirubin ceases




Fig. 28.5 Kleihauer test for fetal red cells; a deeply eosin-staining cell containing fetal haemoglobin is seen at the centre. Haemoglobin has been eluted from the other red cells by an incubation at acid pH and these appear as colourless ghosts.




Fig. 28.6 Doppler ultrasonography of the circle of Willis in a fetus. The cursor is placed over the middle cerebral artery and an increased blood velocity correlates with anaemia. (From Kumar and Regan, 2005 with permission)

and conjugation of bilirubin by the neonatal liver has not yet reached full activity.

3 Mild disease Mild anaemia with or without jaundice.

Investigations will reveal variable anaemia with a high reticulocyte count; the baby is Rh D-positive, the direct antiglobulin test is positive and the serum bilirubin raised. In moderate and severe cases, many erythroblasts are seen in the blood film (Fig. 28.7b) and this is known as *erythroblastosis fetalis*.

Treatment

Exchange transfusion may be necessary; the indications for this include severe anaemia (Hb <10 g/dL at birth) and severe or rapidly rising hyperbilirubinaemia. More than one exchange transfusion may be required and 500 mL is usually sufficient for each exchange. The donor blood should be less than 5 days old, CMV negative and irradiated. Rh D-negative and ABO compatible with the baby's and mother's serum. Phototherapy (exposure of the infant to bright light of appropriate wavelength) degrades bilirubin and reduces the likelihood of kernicterus.

ABO haemolytic disease of the newborn

In 20% of births, a mother is ABO incompatible with the fetus. Group A and group B mothers usually have only IgM ABO antibodies (p. 338). The major-

ity of cases of ABO HDN are caused by 'immune' IgG antibodies in group O mothers. Although 15% of pregnancies in white people involve a group O mother with a group A or group B fetus, most mothers do not produce IgG anti-A or anti-B and very few babies have severe enough haemolytic disease to require treatment. Exchange transfusions are needed in only 1 in 3000 infants. The mild course of ABO HDN is partly explained by the A and B antigens not being fully developed at birth and by partial neutralization of maternal IgG antibodies by A and B antigens on other cells, in the plasma and tissue fluids.


In contrast to Rh HDN, ABO disease may be found in the first pregnancy and may or may not affect subsequent pregnancies. The direct antiglobulin test on the infant's cells may be negative or only weakly positive. Examination of the blood film shows autoagglutination and spherocytosis, polychromasia and erythroblastosis.

Bibliography

- Finning K.M., Martin P.G., Soothill P.W., Avent N.D. (2002) Prediction of fetal D status from maternal plasma: introduction of a new non-invasive fetal RHD genotyping service. *Transfusion* 42,1079–85.
- Greer I.A. and Nelson-Piercy C. (2005) Low-molecular-weight heparins for thromboprophylaxis and treatment



(a)



(b)

Fig. 28.7 (a) Ultrasound features of hydrops fetalis showing skin oedema, hepatomegaly and ascites. (From Kumar and Regan, 2005 with permission.) (b) Rh haemolytic disease of the newborn (erythroblastosis fetalis): peripheral blood film showing large numbers of erythroblasts, polychromasia and crenated cells.

of venous thromboembolism in pregnancy: a systematic review of safety and efficacy. *Blood* 106,401–7.
 Kumar S. and Regan F. (2005) Management of pregnancies with Rh D alloimmunisation. *BMJ* 330,1255–8.
 Mari G., Deter R.L., Delle Chiaie L. *et al.* (2000) Noninvasive diagnosis by Doppler ultrasonography of fetal anemia due to maternal red-cell alloimmunization. Collabora-

tive Group for Doppler Assessment of the Blood Velocity in Anemic Fetuses. *N Engl J Med* 342,9–14.
 Moise K.J. (2005) Red blood cell alloimmunization in pregnancy. *Semin Hematol* 42,169–78.
 Nathan D.G., Orkin S.H., Look A.T., Ginsberg D. (eds) (2003) *Hematology of Infancy and Childhood*. W.B. Saunders, Philadelphia.

Cluster	Main cellular distribution	Name/comments/function/diagnostic value
CD158	NK cells and T-cell subsets	KIR
CD161	NK cells and some T cells	NKR-P1 family
CD162	Lymphocytes, neutrophils and monocytes	P-selectin glycoprotein ligand
CD163	Monocyte macrophage lineage	
CD164	Haemopoietic cells	MGC-24
CD165	Thymocytes, thymic epithelial cells	AD2/gp37
CD166	Thymic epithelial cells, activated T cells	CD6L
CD167	Epithelial cells	DDR1/receptor tyrosine kinase
CD168	Haemopoietic cells	RHAMM/hyaluronan-binding receptor
CD169	Macrophages	Sialoadhesin, siglec-1/cell–cell and cell–matrix interactions
CD171	Neurons, Epi (sub), lymphocyte (sub)	Homotypic and heterotypic cell adhesion
CD178	Activated T cells, neutrophils	FasL/cytokine
CD179a	Pro B and early pre B cells	VpreB/surrogate light chain
CD179b	Pro B and early pre B cells	λ 5/surrogate light chain
CD184	Wide expression on haemopoietic cells	CXCR4/chemokine receptor/binds to SDF-1
CD233	Erythrocytes	Band 3/anion exchanger and attachment site for cytoskeleton
CD246	Few neural cells, subset of B-cell lymphoma	ALK/tyrosine kinase

ADAM, A disintegrin and metalloprotease; ALK, anaplastic lymphoma kinase; B-CLL, B-chronic lymphocytic leukaemia; c-ALL, common (CD10⁺) acute lymphoblastic leukaemia; CD, cluster differentiation; C3dR, complement 3d receptor; EBV, Epstein–Barr virus; FcR, immunoglobulin Fc fragment receptor; FDC, follicular dendritic cell; G-CSFR, granulocyte colony-stimulating factor receptor; GM-CSFR, granulocyte–macrophage colony-stimulating factor; GP, glycoprotein; ICAM, intercellular adhesion molecule; IFN, interferon; Ig, immunoglobulin; IL, interleukin; Ki, antibody to CD30; KIR, killer inhibitory receptors; LFA, leucocyte function-associated antigen; LPS, lipopolysaccharide; Mac-I, integrin molecule; MHC, major histocompatibility complex; NK, natural killer; PDGF, platelet-derived growth factor; SRCR, sheep red cell receptor; TGF, transforming growth factor; TNFR, tumour necrosis factor receptor; VLA, very late antigens; w, workshop.

Appendix 2

Normal values

	Males	Females	Males and females
Haemoglobin	13.5–17.5 g/dL	11.5–15.5 g/dL	
Red cells (erythrocytes)	$4.5\text{--}6.5 \times 10^{12}/\text{L}$	$3.9\text{--}5.6 \times 10^{12}/\text{L}$	
PCV (haematocrit)	40–52%	36–48%	
MCV			80–95 fL
MCH			27–34 pg
MCHC			20–35 g/dL
White cells (leucocytes)			
total			$4.0\text{--}11.0 \times 10^9/\text{L}$
neutrophils			$2.5\text{--}7.5 \times 10^9/\text{L}$
lymphocytes			$1.5\text{--}3.5 \times 10^9/\text{L}$
monocytes			$0.2\text{--}0.8 \times 10^9/\text{L}$
eosinophils			$0.04\text{--}0.44 \times 10^9/\text{L}$
basophils			$0.01\text{--}0.1 \times 10^9/\text{L}$
Platelets			$150\text{--}400 \times 10^9/\text{L}$
Red cell mass	$30 \pm 5 \text{ mL/kg}$	$27 \pm 5 \text{ mL/kg}$	
Plasma volume	$45 \pm 5 \text{ mL/kg}$	$45 \pm 5 \text{ mL/kg}$	
Serum iron			10–30 $\mu\text{mol/L}$
Total iron-binding capacity			40–75 $\mu\text{mol/L}$ (2.0–4.0 g/L as transferrin)
Serum ferritin*	40–340 $\mu\text{g/L}$	14–150 $\mu\text{g/L}$	
Serum vitamin B ₁₂ *			160–925 ng/L
Serum folate*			3.0–15.0 $\mu\text{g/L}$
Red cell folate*			160–640 $\mu\text{g/L}$

* Normal ranges differ with different commercial kits.

MCH, mean corpuscular haemoglobin; MCHC, mean corpuscular haemoglobin concentration; MCV, mean corpuscular volume; PCV, packed cell volume.

Appendix 3

Table 1 Proposed World Health Organization (WHO) classification of myeloid neoplasms.*

Myeloproliferative disease (MPD)

Chronic myelogenous leukaemia, Philadelphia chromosome (Ph1) [t(9;22)(q34;q11), BCR/ABL] +
 Chronic neutrophilic leukaemia
 Chronic eosinophilic leukaemia/hypereosinophilic syndrome
 Chronic idiopathic myelofibrosis
 Polycythaemia vera
 Essential thrombocythaemia
 Myeloproliferative disease, unclassifiable

Myelodysplastic/myeloproliferative diseases

Chronic myelomonocytic leukaemia (CMML)
 Atypical chronic myelogenous leukaemia (aCML)
 Juvenile myelomonocytic leukaemia (JMML)

Myelodysplastic syndromes (MDS)

Refractory anaemia (RA)
 with ringed sideroblasts (RARS)
 without ringed sideroblasts
 Refractory cytopenia (myelodysplastic syndrome) with multilineage dysplasia (RCMD)
 Refractory anaemia (myelodysplastic syndrome) with excess blasts (RAEB)
 5q-syndrome
 Myelodysplastic syndrome, unclassifiable

Acute myeloid leukaemias (AML)*

Acute myeloid leukaemias with recurrent cytogenetic translocations

AML with t(8;21)(q22;q22), AML1(CBF α)/ETO
 Acute promyelocytic leukaemia (AML with t(15;17)(q22;q11–12) and variants, PML/RAR α)
 AML with abnormal bone marrow eosinophils (inv(16)(p13q22) or t(16;16)(p13;q11), CBF β /MYH11X)
 AML with 11q23 (MLL) abnormalities

Acute myeloid leukaemia with multilineage dysplasia

with prior myelodysplastic syndrome
 without prior myelodysplastic syndrome

Acute myeloid leukaemia and myelodysplastic syndrome, therapy related

Alkylating agent related
 Etoposide/anthracycline related (some may be lymphoid)
 Other types

(Continued)

Table 1 Proposed World Health Organization (WHO) classification of myeloid neoplasms (*Continued*).**Acute myeloid leukaemia not otherwise categorized*

- AML minimally differentiated
- AML without maturation
- AML with maturation
- Acute myelomonocytic leukaemia
- Acute monocytic leukaemia
- Acute erythroid leukaemia
- Acute megakaryocytic leukaemia
- Acute basophilic leukaemia
- Acute panmyelosis with myelofibrosis

Acute biphenotypic leukaemias

* Only major disease categories are listed.

Table 2 Proposed World Health Organization (WHO) classification of lymphoid neoplasms.*

B-CELL NEOPLASMS**Precursor B-cell neoplasm**

Precursor B-lymphoblastic leukaemia / lymphoma (precursor B-cell acute lymphoblastic leukaemia)

Mature (peripheral) B-cell neoplasms**

- B-cell chronic lymphocytic leukaemia/small lymphocytic lymphoma*
- B-cell prolymphocytic leukaemia
- Lymphoplasmacytic lymphoma
- Splenic marginal zone B-cell lymphoma (+/- villous lymphocytes)
- Hairy cell leukaemia
- Plasma cell myeloma / plasmacytoma*
- Extranodal marginal zone B-cell lymphoma of MALT type*
- Nodal marginal zone B-cell lymphoma (+/- monocytoid B cells)
- Follicular lymphoma*
- Mantle cell lymphoma*
- Diffuse large B-cell lymphoma*
- Mediastinal large B-cell lymphoma
- Primary effusion lymphoma
- Burkitt's lymphoma / Burkitt's cell leukaemia*

T AND NK-CELL NEOPLASMS**Precursor T-cell neoplasm**

Precursor T-lymphoblastic lymphoma / leukaemia (precursor T-cell acute lymphoblastic leukaemia)

Mature (peripheral) T-cell neoplasms**

- T-cell prolymphocytic leukaemia
- T-cell granular lymphocytic leukaemia
- Aggressive NK-cell leukaemia
- Adult T-cell lymphoma / leukaemia (HTLV1+)
- Extranodal NK/T-cell lymphoma, nasal type
- Enteropathy-type T-cell lymphoma
- Hepatosplenic $\gamma\delta$ T-cell lymphoma
- Subcutaneous panniculitis-like T-cell lymphoma

(Continued)

Table 2 Proposed World Health Organization (WHO) classification of lymphoid neoplasms (*Continued*).*

<i>Mycosis fungoides</i> /Sézary syndrome
Anaplastic large-cell lymphoma, T/null cell, primary cutaneous type
<i>Peripheral T-cell lymphoma, not otherwise characterized</i>
<i>Angioimmunoblastic T-cell lymphoma</i>
<i>Anaplastic large cell lymphoma, T/null cell, primary systemic type</i>
HODGKIN'S LYMPHOMA (HODGKIN'S DISEASE)
Nodular lymphocyte predominance Hodgkin's lymphoma
<i>Classic Hodgkin's lymphoma</i>
Nodular sclerosis Hodgkin's lymphoma (Grades 1 and 2)
Lymphocyte-rich classical Hodgkin's lymphoma
Mixed cellularity Hodgkin's lymphoma
Lymphocyte depletion Hodgkin's lymphoma

* More common entities are in italics.

** B- and T/NK-cell neoplasms are grouped according to major clinical presentations (predominantly disseminated/leukaemic, primary extranodal, predominantly nodal).

Table 3 Mast cell diseases.

Cutaneous mastocytosis
Systematic mast cell disease (+/- skin involvement)
Systematic mast cell disease with associated haematological disorder (+/- skin involvement)
Mast cell leukaemia/sarcoma

Table 4 Histiocytic and dendritic-cell neoplasms.

Macrophage/histiocytic neoplasm
Histiocytic sarcoma
Dendritic-cell neoplasms
Langerhans cell histiocytosis
Langerhans cell sarcoma
Interdigitating dendritic cell sarcoma/tumour
Follicular dendritic cell sarcoma/tumour
Dendritic cell sarcoma, not otherwise specified (NOS)

Table 5 Plasma cell disorders: subtypes and variants.

Monoclonal gammopathy of undetermined significance (MGUS)
<i>Plasma cell myeloma variants:</i>
Indolent myeloma
Smoldering myeloma
Osteosclerotic myeloma (POEMS syndrome)
Plasma cell leukaemia
Non-secretory myeloma
<i>Plasmacytoma variants:</i>
Solitary plasmacytoma of bone
Extramedullary plasmacytoma

Table 6 Immunosecretory disorders (clinical manifestations of diverse lymphoid neoplasms).

Clinical syndrome	Underlying neoplasm
Waldenström's macroglobulinemia	Lymphoplasmacytic lymphoma
Heavy chain diseases (HCD):	
γ HCD	Lymphoplasmacytic lymphoma
α HCD	Extranodal marginal zone lymphoma (immunoproliferative small intestinal disorder)
μ HCD	B-cell chronic lymphocytic leukaemia
Immunoglobulin deposition diseases:	
Systemic light chain disease	Plasma cell myeloma, monoclonal gammopathy
Primary amyloidosis	Plasma cell myeloma, monoclonal gammopathy

Index

Note: Page numbers in *italic* refer to figures and tables

- abciximab 287, 318, 319
- ABO blood group system 338–9, 340, 358
- abortion, recurrent 309
- acanthocytes 20, 24
- N*-acetylglucosaminyltransferase 247
- aciclovir, prophylactic 151
- acid phosphatase 95
- acrocyanosis 68
- acrolein 261
- actin 19, 20
- activated oxygen species 100
- activated partial thromboplastin time 276, 300, 310, 356
- activated protein C resistance 306, 307
- acute biphenotypic leukaemias 367
- acute lymphoblastic leukaemia (ALL) 157, 158–67, 214
 - aetiology 130–1
 - and aplastic anaemia 244
 - diagnosis 142, 158
 - differentiation from AML 157–8
 - genetics 133, 135, 137, 138, 161–2, 164, 174
 - minimal residual disease 144–6, 165
 - treatment 98, 156, 164–6, 166
- acute myeloid leukaemia (AML) 157, 167–72
 - and aplastic anaemia 244, 245
 - classification 158, 160, 167, 168, 169, 366–7
 - diagnosis 142, 158
 - differentiation from ALL 157–8
 - establishment of treatment protocol 143–4
 - familial tendency 129
 - genetics 133, 135, 136, 138, 167
 - immunophenotype 159, 167
 - minimal residual disease 144–6
 - progression from myelodysplasia 182
 - transformation 180, 230, 239, 244
 - treatment 98, 156, 169–72
 - types 155–6, 158, 168, 169, 170, 299
- acute phase proteins 333
- acute phase response 333, 335–6
- acute sickle chest syndrome 87
- acylated plasminogen streptokinase activator complex 317
- ADAMTS13 269, 284, 285
- ADCC 109
- S*-adenosyl homocysteine 51
- adhesion molecules 11, 99
- adipocytes 3
- ADP 267, 269
- adriamycin 153, 154
- adult T-cell leukaemia/lymphoma 131, 195, 214
- adverse drug reactions
 - aplastic anaemia 242, 244
 - haematological malignancies 129–30
 - haemolytic anaemia 66, 68
 - neutropenia 104, 105
 - thrombocytopenia 280, 281, 284, 313, 314
- African macroglobulinaemia 125–6, 330
- AGM region 1
- agnogenic myeloid metaplasia 238
- AIDS *see* HIV infection
- AIHAs 66–8, 189, 282, 321
- ALA synthase 16, 40
- albumin 219, 349
- alcohol 49, 56, 198
- Alder's anomaly 102
- alemtuzumab 213
- ALG 186, 244, 245
- alkylating agents 129, 153, 155, 244
- ALL *see* acute lymphoblastic leukaemia
- all-*trans* retinoic acid 138, 154, 155–6, 170
- allopurinol 149, 179, 239
- α_2 -antiplasmin 275
- α_2 -macroglobulin 275
- α -heavy chain disease 223
- α -mannosidase 247
- α -spectrin 19, 20
- Alzheimer's disease 51
- aminocaproic acid 280
- δ -aminolaevulinic acid synthase 16, 40
- AML *see* acute myeloid leukaemia
- AML1 protein 10, 162
- amniotless 49
- amniotic fluid embolism 299
- amoxicillin in infectious mononucleosis 118
- ampicillin 68, 118, 119
- amyloid P 225
- amyloidosis 217, 219, 221, 225, 226, 227, 228
- anaemia 20–1
 - in ALL 159
 - in AML 169
 - classification 21
 - clinical features 21–3
 - in CLL 189
 - in CML 175
 - effects of age on ability to tolerate 22
 - in hereditary spherocytosis 62
 - in HIV infection 328–9
 - in Hodgkin's lymphoma 198
 - in infection 327
 - laboratory findings 23–6
 - in lymphoplasmacytoid lymphomas 211
 - macrocytic 21, 44–57
 - in malignant disease 320–1, 322
 - microcytic hypochromic 21, 28, 28, 39, 41
 - in multiple myeloma 217, 218, 220, 223
 - in myelodysplasia 182, 183
 - neonatal 355
 - in non-Hodgkin's lymphoma 204, 206
 - normocytic normochromic 21
 - physiological 352
 - in renal failure 324
 - severity 21–2
 - speed of onset 21
 - in Waldenström's macroglobulinaemia 224
- anaemia of chronic disorders 28, 39–40, 41, 320, 323
- anaemia of prematurity 355
- anaesthesia and sickle cell anaemia 89
- anagrelide 238
- anaphylaxis 114, 346
- anaplastic large cell lymphoma 133, 208, 214–15
- androgen therapy in Fanconi's anaemia 244, 245
- anencephaly 51
- angiocentric lymphomas 214
- angular stomatitis/cheilosis 34, 50
- ankle ulcers 59, 88
- ankyrin 19
- anorexia 175, 198
- anthracyclines 153, 154, 242
- anti-A 338
- anti-B 338
- antibiotics
 - cytotoxic 153, 154
 - in fever associated with haematological malignancy 150–1
 - in glucose-6-phosphate dehydrogenase deficiency 64
 - immune haemolytic anaemia due to 68
 - neutropenia due to 104
 - prophylactic 106, 127, 150, 261
- antibody-dependent cell-mediated cytotoxicity 109
- anti-C 338, 338, 356
- anticardiolipin antibodies 309
- anti-CD20 *see* rituximab
- anti-CD52 67, 68, 154, 156, 192, 246
- anticoagulants 96, 311–16
 - and chemotherapy 148
 - in DIC 300
 - following splenectomy 127
 - in paroxysmal nocturnal haemoglobinuria 70
 - in pregnancy 313, 314, 354

- thrombocytopenia due to 284, 313, 314
 as vitamin K antagonist 296–7
 anticonvulsants 49, 104, 242
 anti-D 338, 339, 356–8
 antidepressants 104, 242
 anti-DNA antibodies 324
 anti-E 339, 356, 339
 antiemetics in chemotherapy 148–9
 anti-erythropoietin antibodies 246
 antifolate drugs 48, 56
 antifungal agents 106, 258
 antigen-presenting cells 99, 100, 115
 antigen-receptor gene arrangements 112–13, 114
 antiglobulin (Coombs') test 66, 114, 343
 antihelminthic agents 64
 anti-inflammatory drugs 242
 anti-K 356
 antilymphocyte globulin 186, 244, 245
 antimetabolites 153, 244
 antinuclear factor 324
 antiphospholipid syndrome 309
 antiplatelet drugs 287, 317–19
 antithrombin 273, 275, 356
 deficiency 306
 antithymocyte globulin 245
 antithyroid drugs 242
 aorta-gonads-mesonephros region 1
 aortic embolism 304
 Apaf-1 9
 APCs 99, 100, 115
 aplastic anaemia 70, 241–6, 326, 328
 aplastic crises 59, 62, 88
 apoferritin 29
 apoptosis 9–10, 132
 APSAC 317
 APTT 276, 300, 310, 356
 arachidonate 269
 argatroban 313
 arsenic 70, 154, 156
 arsine 70
 arterial embolism 303, 304
 asparaginase 154, 156
 aspergillosis 151, 152, 258, 259
 aspirin 317, 318
 contraindications 287
 discontinuation before chemotherapy 148
 effects on platelet function 287
 in essential thrombocythaemia 238
 following splenectomy 127
 in glucose-6-phosphate dehydrogenase deficiency 64
 in polycythaemia rubra vera 235
 ataxia telangiectasia 129
 ATG 245
 atherosclerosis 303–4
 ATLL 131, 195, 214
 ATRA 138, 154, 155–6, 170
 ATRA syndrome 156
 atrophic gastritis 35, 49
 autoimmune thrombocytopenic purpura 66, 118, 281–3, 287–8, 353–4
 autospontaneous in sickle cell anaemia 88
 avascular necrosis 86
 azacytidine 136, 156
 azathioprine 153
 AZT 56
 azurophilic granules 195
 B-cell neoplasms 367
 B-cell receptor 108, 115
 B-lymphocytes 108
 circulation 109
 in CLL 189, 190
 differentiation 2
 functional aspects 111
 in immune response 115–17
 naïve/virgin 115
 babesiosis 342
 bacteria, phagocytosis and destruction 99–101
 bacterial infection 326–7, 328
 adjuvant growth factor therapy 99
 in β -thalassaemia major 80
 in CLL 189
 following stem cell transplantation 258, 260, 261
 in haematological malignancies 149–51
 in multiple myeloma 217, 223
 in neutropenia 105–6
 prevention in hyposplenism 127
 purpura in 279
 transmission by blood transfusion 342, 343
 band 3 protein 20, 63
 barbiturate-induced folate deficiency 49
 bartonellosis 327
 basket cells 24
 basophilia 104, 175
 basophils 2, 94, 95, 96, 365
 BAX 10
 BCL-1 gene 212
 BCL-2 gene/BCL-2 protein 10, 132, 135, 138, 209, 212
 BCNU 129, 153
 BCR-ABL gene/BCR-ABL fusion protein 135, 174, 175–6
 Behçet's disease 309
 Bence-Jones protein 112, 218, 219, 221
 benzene 129
 Bernard-Soulier syndrome 266, 287
 β_2 -microglobulin in multiple myeloma 219, 223
 β spectrin 19, 20
 β -storage pool disease 287
 β -thromboglobulin 267
 BFUs 2, 12, 234
 big spleen disease 125–6, 330
 biliary obstruction 297
 bilirubin 26, 52, 58, 59
 Birbeck granules 106
 bisphosphonates in multiple myeloma 223
 bite cells 65
 bivalirudin 316
 blackwater fever 69, 330
 blast cells 157, 206
 bleeding disorders
 due to defective coagulation 278, 290–302
 due to defective platelet function 278, 286–8
 with thrombocytopenia 280–6
 vascular 278–80
 and venous thrombosis 305–8
 bleeding time 277, 288
 bleomycin 153, 154
 blister cells 65
 blood
 fetal sampling 92
 peripheral stem cell collection 249, 251
 blood coagulation 268, 270–4, 276–7
 disorders 278, 290–301
 in malignant disease 322
 blood film
 in ALL 159
 in anaemia 23, 24, 25
 in autoimmune haemolytic anaemia 67
 in β -thalassaemia major 80, 81
 in bleeding disorders 276
 in CLL 189, 190
 in CML 175, 177
 dimorphic 35–6
 in hairy cell leukaemia 193
 in iron deficiency 35–6
 in myelodysplasia 182–3, 184
 in platelet disorders 287–8
 in polymphocytic leukaemia 193
 in thrombophilia 310
 blood group systems 337–41
 blood loss 21, 350
 blood products 147–8, 346–50
 blood transfusion 337
 ABO incompatible 68
 in aplastic anaemia 245
 autologous 347, 349
 in β -thalassaemia major 78, 82
 blood components/products 147–8, 347–50
 blood group serology 343, 344
 complications 78, 283–4, 344–6
 cross-matching and pre-transfusion tests 343–4
 donors 337, 338
 following stem cell transplantation 260
 in haematological malignancy 147
 haemolytic transfusion reactions 344–6
 infection due to 80, 341–3
 massive 286, 301
 in myelodysplastic syndromes 186
 in myelofibrosis 239
 neonatal 358
 red cell antigens and blood group antibodies 337–41
 reduction of need for 346–7
 in sickle cell anaemia 89
 in sideroblastic anaemia 41
 in warm autoimmune haemolytic anaemia 67
 Bloom's syndrome 129
 bone lesions in multiple myeloma 218, 219, 220, 223
 bone marrow
 in ALL 159
 in anaemia 24, 25, 26
 in anaemia of chronic disorders 39
 in aplastic anaemia 242, 244
 aspiration 24, 25, 26
 in CLL 189
 in CML 177
 erythropoiesis 12
 failure 281
 fetal 1
 granulopoiesis 95, 96–7
 in haemolytic anaemias 59
 haemopoiesis 1
 hyperplasia 78
 hypoplasia 70
 infancy and childhood 1
 iron assessment 36
 irradiation 130
 in megaloblastic anaemia 52
 metastatic carcinoma in 320, 321
 in multiple myeloma 217, 218
 in myelodysplasia 182, 183, 184
 in myelofibrosis 235, 238–9
 in non-Hodgkin's lymphoma 206, 208, 209
 in polycythaemia rubra vera 233, 235
 as primary lymphoid organ 108, 110
 stem cells 1–3, 249
 stroma 3–4
 structure 2
 transplantation see stem cell transplantation
 trephine biopsy 24, 25, 26
 bortezomib 213, 223
 BRCA2 gene 241

- bronchiolitis obliterans 261
brucellosis 342
bruising, easy 279
Burkitt's lymphoma 131, 213–14
 genetics 133, 135, 137, 138, 207, 208
 in HIV infection 329
 immunophenotype 207
burns 70
burr cells 324
burst-forming units 2, 12, 234
busulfan 153, 235, 242, 244
- c-Kit 4
c-Mpl receptors 265
C-reactive protein 198, 218, 320, 335–6
cachexia in Hodgkin's lymphoma 198
caeruloplasmin 29
calcium 267
cAMP 267, 269
Campath-1H 67, 68, 154, 156, 192, 246
Candida infection 151, 258
cannabinoids in chemotherapy 149
carbamazepine 104
carbimazole 104
carbon dioxide exchange 17–18
cardiolipin 301
cardiovascular disease 56
carotid artery thrombosis 303
caspases 9, 10
catalase 30
cathepsin G 100
CBF 137, 138, 167
CCNU 129, 153
CD molecules 70, 108, 159, 160, 191, 207,
 245, 360–4
 see also transferrin receptors
CDAs 246–7
Cdk 9
cell cycle 7, 9
central nervous system in ALL 159, 164–5
central venous catheter insertion 147, 148
centroblasts 117, 206
centrocytes 117, 206
cerebrospinal fluid in ALL 159, 164–5
CFU-GEMM 2, 3, 12
CFUs 2, 12, 234
Chagas' disease 342–3
Chédiak-Higashi syndrome 101, 102
chemokine receptors 100
chemokines 100
chemotaxis 99, 101
chemotherapy 153, 153–4
 ALL 164, 165, 166
 AML 170, 171
 before stem cell transplantation 250, 252
 Burkitt's lymphoma 213–14
 CLL 191–2
 CML 179
 diffuse large B-cell lymphoma 213
 follicular lymphoma 209–10
 growth factor therapy following 97
 Hodgkin's lymphoma 201–2
 immunodeficiency following 121
 late effects 188, 202
 leukaemia following 188
 mantle cell lymphoma 212
 multiple myeloma 220, 222
 in myelodysplastic syndromes 186
 polycythaemia rubra vera 234–5
 prolymphocytic leukaemia 193
 and stem cell population expansion 249
 support therapy 147–9
 Waldenström's macroglobulinaemia
 224
chickenpox 104
chlorambucil 68, 129, 153, 191–2
chloramphenicol 104, 242, 244
chlorate poisoning 70
2-chlorodeoxyadenosine 154, 194
chloroquine 64
chlorpromazine 104
chondroitin 4
Christmas disease 294, 295, 350
chromosomes
 duplications 135, 136
 nomenclature 134–5
 sex 134
 translocations 132, 134, 135, 136, 137,
 138
chronic eosinophilic leukaemia 104, 133
chronic lymphocytic leukaemia (CLL)
 188–92
 familial tendency 129
 genetics 133, 135, 136, 138, 189–90, 191,
 207
 and idiopathic thrombocytopenic
 purpura 282
 immunodeficiency in 121, 189
 immunophenotype 189, 190, 207
 and lymphocytic lymphoma 205, 206,
 207, 211
 and red cell aplasia 246
 Richter's transformation 192
chronic lymphoid leukaemias 188–96
chronic myeloid leukaemia (CML) 174–81,
 232
 basophilia in 104, 175
 donor leucocyte infusion 261–2
 genetics 133, 135, 138, 174, 175–6
 minimal residual disease 144–6
 Philadelphia-negative (atypical) 174,
 180, 186
 splenomegaly in 125, 175
chronic myelomonocytic leukaemia 181,
 186–7
chronic neutrophilic leukaemia 181
chronic obstructive airways disease 236
ciclosporin 186, 192, 244, 245
ciprofloxacin 150
cisplatin 153, 156
CLL see chronic lymphocytic leukaemia
clodronate 223
clonal progression 132
clopidogrel 148, 287, 318–19
Clostridium perfringens 69–70, 327, 328
clozapine 104
CLP 3
cluster differentiation molecules see CD
 molecules
CML see chronic myeloid leukaemia
CMV infection 147, 151, 258–60, 328, 342,
 346
coagulation cascade 270, 276
coagulation factors 266, 268, 269, 270–3,
 276, 277
 antibodies to 300–1
 hereditary disorders 290–6
 inhibitors 273–4
cobalamins see vitamin B₁₂
cobalt and erythropoiesis 15
cold agglutinin disease 112
cold antibodies 338
collagen vascular diseases 309, 324
collagenase 95
colon carcinoma 38
colonoscopy 37, 38
colony-forming units 2, 12, 234
common iliac artery embolism 304
common lymphoid precursor 3
common variable immunodeficiency 121
complement 94, 99, 113–14
compression stockings 316
- computed tomography (CT)
 combined with PET scanning 209, 211
 Hodgkin's lymphoma 200, 201
 non-Hodgkin's lymphoma 208–9, 211
 spleen 125
computed tomography pulmonary
 angiography 311
congenital dyserythropoietic anaemias
 246–7
conjunctiva 22, 233
connective tissue disorders 279, 323–4
contrast venography in deep vein
 thrombosis 310
Coombs' test 66, 114, 343
core binding factor 137, 138, 167
corticosteroids
 in CLL 192
 in haematological malignancy 154, 155
 in idiopathic thrombocytopenic purpura
 283
 in infectious mononucleosis 119
 in neutropenia 106
 purpura associated with therapy 280
 in warm autoimmune haemolytic
 anaemia 67
co-trimoxazole 48, 64, 104
Creutzfeldt-Jacob disease, new variant 343,
 346
Crohn's disease 49
CRP 198, 218, 320, 335–6
cryoprecipitate 148, 349
CT see computed tomography
CT pulmonary angiography 311
cubilin 45, 49
cyanosis in polycythaemia rubra vera 233
cyclic adenosine monophosphate 267, 269
cyclin-dependent protein kinases 9
cyclins 9, 207, 212, 216
cyclozine 149
cyclophosphamide 153, 192, 242
cystathione β -synthase deficiency 307
cytochrome c 9–10
cytokinesis 9
cytomegalovirus infection 147, 151,
 258–60, 328, 342, 346
cytosine arabinoside 56, 153
cytotoxic drugs see chemotherapy
- D-dimer plasma concentration following
 DVT 310
dactylitis in sickle cell anaemia 87
danaparoid 313
danazol 239, 283
dapsone 64, 70
DAT (Coombs') test 66, 114, 343
daunorubicin 154, 244
DDAVP 293
DEB test 242
deep vein thrombosis 305, 310–11
deferasirox 83
deferiprone 83, 104
deferroxamine 83
demarcation membrane 265
dendritic-cell neoplasms 368
dendritic cells 99, 106, 115, 117, 124
deoxycoformycin 154, 194
desmopressin 293
dexamethasone 148
DHF 47, 48
diabetes 309
dialysis 223, 324
Diamond-Backfan syndrome 246, 247
diaphorase deficiency 18
DIC see disseminated intravascular
 coagulation
diepoxybutane test 242

- diffuse large B-cell lymphoma 138, 205, 206, 207, 208, 213, 329
- dihydrofolate 47, 48
- 2,3-diphosphoglycerate 17–18, 22
- dipyridamole 287, 317–18
- direct antiglobulin (Coombs') test 66, 114, 343
- disseminated intravascular coagulation (DIC) 298–300
- in AML 169
- in bacterial infection 326
- in giant cavernous haemangioma 279
- in malaria 330
- in malignant disease 322, 323
- red cell fragmentation in 69
- thrombocytopenia in 285
- diuretics in blood transfusion 147
- divalent metal transporter-1 30, 32–3
- DKC1* gene 242
- DLCL 138, 205, 206, 207, 208, 213, 329
- DLI 261–2
- DMT-1 30, 32–3
- DNA
- complementary (cDNA) 140
- microarray analysis 140–1, 142, 208
- prenatal diagnostic techniques 90, 91–2
- Döhle bodies 101, 102, 103, 326
- domperidone in chemotherapy 149
- Donath–Landsteiner antibody 68
- donor leucocyte infusions 261–2
- Doppler ultrasound in deep vein thrombosis 305, 310
- Down's syndrome 129
- doxorubicin 153
- 2,3-DPG 17–18, 22
- drug resistance 132
- dsRNA 115
- DVT 305, 310–11
- dysfibrinogenaemia 298, 326
- dyskeratosis congenita 242
- dyskerin 242
- dyspnoea in polycythaemia rubra vera 233
- easy bruising 279
- EBV *see* Epstein–Barr virus
- ecchinocytes 24
- eculizumab 70
- Ehlers–Danlos syndrome 279
- elastase 100
- electrocardiography in pulmonary embolism 311
- elliptocytes 24, 59
- elliptocytosis 20
- Embsden–Meyerhof pathway 18, 19, 65–6
- encephalocoele 51
- endothelial cells 3, 275
- endothelins 275
- endotoxins 97, 98
- eosinophilia 103–4, 198
- eosinophilic granuloma 106
- eosinophilic leukaemia 104, 133, 181
- eosinophils 2, 94, 95, 96, 365
- epidodophyllotoxins 129–30, 154
- epirubicin 153
- epistaxis, recurrent 278
- Epstein–Barr virus (EBV)
- and Burkitt's lymphoma 131, 213
- and Hodgkin's lymphoma 197
- infection following stem cell transplantation 260
- infection in haematological malignancy 151
- and infectious mononucleosis 117–19, 120
- and post-transplant lymphoproliferative disease 131
- transmission by blood transfusion 342, 346
- eptifibatide 287, 318, 319
- ERK 1/2 7
- erythroblastosis fetalis 358, 359
- erythroblasts *see* normoblasts
- erythrocyte sedimentation rate (ESR) 333, 335
- in anaemia of chronic disorders 320
- in Hodgkin's lymphoma 198
- in multiple myeloma 218
- erythrocytes *see* red cells
- erythrocytosis *see* polycythaemia
- erythromelalgia 237
- erythromycin, prophylactic 127
- erythropoiesis 1, 12
- assessment 27
- effective 27
- extramedullary 12, 13
- ineffective 26, 29
- regulation 12–13, 15
- total 27
- erythropoietin 6, 12–13, 15, 17
- in polycythaemia rubra vera 233
- radioimmunoassay 13, 15
- recombinant 14–15, 40, 148, 186, 320, 324–5, 347
- Escherichia coli* 149, 285
- ESR *see* erythrocyte sedimentation rate
- essential thrombocythaemia 230, 234, 237–8, 309
- etoposide 129–30, 153, 154
- ETV6 10
- Evans' syndrome 66
- extrinsic Xase 271, 276
- FA 129, 241–4
- facial appearance in β -thalassaemia major 78
- FACS 145
- factor V Leiden gene mutation 306, 307
- factor VIII concentrate 293, 350
- factor IX
- concentrate 309, 350
- deficiency 294, 295, 350
- see also* coagulation factors
- Fanconi's anaemia 129, 241–4
- FAND1* gene 241
- Fansidar 64
- Fas 9
- fava beans 64
- FDCs 117
- Felty's syndrome 323
- ferric hydroxide-sucrose 38
- ferritin 28, 29, 30, 365
- in anaemia of chronic disorders 39
- in β -thalassaemia major 81–2
- depletion 33
- in iron deficiency 37
- in iron overload 42, 43
- regulation of synthesis 30–1
- ferroportin 30, 31, 32, 42, 43
- ferrous gluconate 38
- ferrous sulphate 38
- fertility and haematological malignancy 149
- fetomaternal alloimmune thrombocytopenia 356
- fetomaternal haemorrhage 357
- fetus
- blood sampling 92
- haemopoiesis 1
- fever
- in ALL 159
- in Hodgkin's lymphoma 198
- in infectious mononucleosis 118
- in neutropenia 150
- in non-Hodgkin's lymphoma 204
- FFP 147, 148, 300, 300, 348, 349
- fibrin 268, 270, 271, 272–3, 274, 276, 298–9
- fibrinogen 267, 268, 270, 272, 276
- assay 276, 277
- defects 308
- deficiency 299
- elevated levels 304
- structure 273
- fibrinolysis 274–5, 277
- fibrinolytic agents 275, 317
- fibroblasts 3
- fibronectin 4
- filariasis 331
- FIP1L1-PDGFR* fusion gene 181
- FISH 138–9, 140
- 5q-syndrome 183, 185, 186
- flow cytometry 141, 143
- Flt-3* gene 135, 167
- Flt ligand (Flt-L) 6
- fludarabine 66, 67, 68, 153, 154, 192
- fluorescence-activated cell sorting 145
- fluorescent *in situ* hybridization analysis 138–9, 140
- FMAIT 356
- FMH 357
- FMS gene 183
- foam cells 333
- FoG transcription factor 12–13, 73
- folate
- absorption 46, 47
- deficiency 40, 47–8, 49–50, 51–2, 52–6
- aplastic crisis in 59
- with iron deficiency 35–6
- in pregnancy 49, 353
- and erythropoiesis 15
- function 47
- metabolic abnormalities 56
- nutritional aspects 45
- red cell 52, 365
- reduction 48
- serum 52, 365
- structure 47
- supplements 54–6
- in β -thalassaemia major 82
- in hereditary spherocytosis 63
- in myelofibrosis 239
- in pregnancy 51, 55, 353
- in sickle cell anaemia 89
- in sideroblastic anaemia 40
- in warm autoimmune haemolytic anaemia 67
- transport 47
- folate polyglutamates 47, 48
- folinic acid 48, 153
- follicular dendritic cells 117
- follicular lymphoma 194, 205, 209–10, 212
- genetics 10, 133, 135, 138, 207, 208
- immunophenotype 190, 207
- fondaparinux 315–16
- fresh frozen plasma 147, 148, 300, 300, 348, 349
- frusemide 104, 147
- fungal infection
- adjuvant growth factor therapy 99
- in CLL 189
- following stem cell transplantation 258
- in haematological malignancy 151–2
- in multiple myeloma 217, 223
- in neutropenia 105–6
- fusion proteins 135, 136, 137, 138, 141, 156, 162, 174, 175–6

- G-CSF
clinical applications 97–9, 106
in haemopoiesis 5, 6, 97, 98
in large granular lymphocytic leukaemia 195
in multiple myeloma 220
in myelodysplastic syndromes 186
pegylated 97, 98
in stem cell mobilization 4
and stem cell population expansion 249, 250, 251
- G6PD 18, 20, 63–4, 63–5
gallstones 58–9, 60, 62, 66, 88
gastrectomy 35, 48, 49, 55
gastric mucosa B-cell lymphoma 131, 133, 207, 212
gastrointestinal tract
atrophic gastritis 35, 49
blood loss from 37, 38, 278–9, 291
malignant disease 321, 322
in non-Hodgkin's lymphoma 204
- GATA-1 5, 12–13, 73
GATA-2 10
Gaucher's disease 125, 331–3, 334
gene therapy 262–3, 294
genetic counselling in haemoglobin disorders 90–2
genetic markers in haematological malignancy 143–6
germ cells 134–5
giant cavernous haemangioma 279
Gilbert's disease 62
glandular fever 67, 68, 117–19, 120, 125
Glanzmann's disease 266, 287
Glivec (imatinib) 104, 154, 155, 177–9
globin 72–3, 74
glossitis 34, 50
glucose-6-phosphate 18, 20
glucose-6-phosphate dehydrogenase 18, 20, 63–4, 63–5
glucosylceramide deficiency 310
glutathione deficiency 65
gluten-induced enteropathy 35, 37, 49, 54
glycoproteins
platelet 266, 267, 268–9
stromal 4
glycosaminoglycans 4
glycosylphosphatidylinositol anchor 70
GM-CSF 4, 6, 97–9, 195
gold 104, 242, 244, 323
gout 233, 234
GPI anchor 70
GPIIb/IIIa 268
GPIIb/IIIa receptors 269
graduated compression stockings 316
graft-versus-host disease 252, 254, 255, 256–8, 346
graft-versus-leukaemia effect 254, 255, 261–2
granisetron 148
granulocyte colony-stimulating factor *see* G-CSF
granulocyte-macrophage colony-stimulating factor 4, 6, 97–9, 195
granulocytes 94–6, 99–101
concentrates 349
differentiation 2, 95, 96–7, 98
granulocytic sarcoma 169
granulopoiesis 2, 95, 96–7, 98
grey platelet syndrome 287
growth factor receptors 6–7, 8
growth factors 5–7, 8, 97–9
gum hypertrophy in AML 169, 170
GVHD 252, 254, 255, 256–8, 346
GVL 254, 255, 261–2
- haem 16–17, 17, 32, 58
haemarthrosis 291, 292, 293
haematocrit 16, 365
haematological malignancy 125, 129–56
haematuria 37, 90, 291
haemochromatosis 41–3
haemoglobin
in acute blood loss 21, 350
in anaemia of chronic disorders 39
in β -thalassaemias 80–1, 82
deoxygenation 17
electrophoresis 80–1, 82, 89
function 17–18
genetic disorders 74–93
Gower 1 72
Gower 2 72
in haemolytic anaemias 60
Hb A (adult) 16, 72
Hb A₂ 16, 41, 72, 80–1, 82
Hb C 74, 75, 82, 89, 90
Hb Constant Spring 77
Hb D 74, 75, 90
Hb E 74, 75, 85, 90
Hb F (fetal) 16, 18, 41, 72, 80, 85, 89, 355
Hb H 41, 76, 77, 82, 84
Hb Lepore 75, 84, 92
Hb M 18
Hb S (sickle) 17, 22, 74, 82, 85, 87, 90
high performance liquid chromatography 81, 82
iron in 29, 30
neonatal 354
normal red cell values 16, 20, 21, 365
in normoblasts 12, 13
oxygenation 17
Portland 72
in sickle cell trait/anaemia 82
structure 16, 72
synthesis 15–17, 72–4
defects 28
unstable 74
haemoglobin oxygen dissociation curve 17, 21, 22
haemoglobinaemia 61
haemoglobinuria 61, 64, 69
haemolysis 58, 59, 60–1, 74
in malaria 69, 330
in sickle cell anaemia 88
haemolytic anaemias
acquired 58–71
autoimmune 66–8, 189, 282, 321
chronic 74
in infectious mononucleosis 118
and macrocytic anaemia 56
microangiopathic 69, 299, 321, 322
in non-Hodgkin's lymphoma 206
haemolytic disease, compensated 58
haemolytic disease of the newborn 68, 356–8, 359
haemolytic shock 344
haemolytic transfusion reactions 344–6
haemolytic uraemic syndrome 285, 328
haemopexin 60
haemophagocytic lymphohistiocytosis/syndrome 106, 107, 328
haemophilias 290–4, 295, 350
haemophilic pseudotumours 291, 292
Haemophilus influenzae 127
haemopoiesis 1, 96–7
bone marrow stroma 3–4
extramedullary 1, 124
growth factors 5–7, 8
ineffective 182
regulation 5–7, 8, 97, 98
sites of 1
transcription factors 5, 10
- haemopoietic malignancy 125, 129–56
haemopoietic progenitors 2–3
haemopoietic stem cells 1–3, 4–5, 108
haemopoietin receptor superfamily 6–7
haemorrhage
in essential thrombocythaemia 237
fetomaternal 357
in heparin therapy 313
intracerebral 291
operative 291
in polycythaemia rubra vera 233
post-traumatic 291
in renal failure 325
retinal 22
haemorrhagic cystitis 261
haemorrhagic disease of the newborn 296, 297
haemorrhagic syndrome 169–70
haemosiderin 29, 30, 33, 60, 79
haemosiderinuria 37, 61, 70
haemostasis 264–5
blood coagulation 268, 270–4
components of response 265–9, 270
during pregnancy 354
and fibrinolysis 274–5
responses 275–6
role of endothelium 275
tests of function 276–7
hair cell leukaemia/variant 190, 193–4
hair cells 193
Ham's test 70
hand-foot syndrome 87
Hand-Schüller-Christian disease 106
haptocorrin 46
haptoglobins 58, 59, 60
Hashimoto's disease 326
Hb H disease 41, 76, 77, 82, 84
Hb S/ β -thalassaemia 90
HCL 190, 193–4
HCP-1 32
HDN 68, 356–9
heavy chain disease 223
Heinz bodies 25, 65, 70
Helicobacter pylori 35, 49, 131, 212
HELLP syndrome 353
hemojuvelin 31, 42, 43
HEMPAS 247
Henoch-Schönlein syndrome 279, 280
heparin 96, 127, 148, 284, 300, 311–13
heparin cofactor II 274
hepatitis
and aplastic anaemia 242, 244, 326, 328
chronic immune 326
and haemophilia 292
immunization 84
transmission in blood transfusion 80, 84, 341, 342
hepatomegaly 77–8, 118, 159, 189, 198, 204, 238
hepcidin 31, 33, 40, 42, 43, 320
hephaestin 33
hereditary elliptocytosis 62, 63
hereditary erythroblast multinuclearity with a positive serum lysis test 247
hereditary haemorrhagic telangiectasia 278–9
hereditary pyropoikilocytosis 63
hereditary spherocytosis 20, 60, 61–3
herpes simplex infection 151, 258
herpes zoster 151, 189
hexose monophosphate pathway 18, 20, 65
HFE gene/HFE protein 31, 41–2
HH-8 131, 342
high molecular weight kininogen 271
hirudin 313
histamine 96, 239

- histiocytic disorders 106–7
 histiocytic neoplasms 368
 histiocytosis X 106
 HIV infection 328–9
 and haemophilia 291
 and Hodgkin's disease 131, 329
 and idiopathic thrombocytopenic purpura 282
 lymphoma associated 131, 329
 lymphopenia in 119
 transmission in blood transfusion 80, 341, 342, 346
 virus entry into cells 100
 HLA system 108, 109, 115, 254–5, 256, 346
 HMWK 271
 Hodgkin cells 198
 Hodgkin's lymphoma 129, 131, 197–202, 282, 323, 329, 368
 homocysteine 46, 49, 51, 52, 56, 304, 307–8
 homocystinuria 307
 hookworm infection 37
 hormone replacement therapy 309
 Howell–Jolly bodies 25, 81, 88, 89, 123, 124, 126
 Hox genes 10
 HPA 268, 356
 HS 20, 60, 61–3
 HS40 73
 HTLV-1 131, 195, 214, 342
 human herpes virus-8 131, 342
 human leucocyte antigens 108, 109, 115, 254–5, 256, 346
 human platelet antigens 268, 356
 human T-cell leukaemia/lymphoma virus type I 131, 195, 214, 342
 Hurler's syndrome 101
 HUS 285, 328
 hyaluronic acid 4
 hybrid acute leukaemia 158
 hydrogen peroxide 100
 hydrops fetalis 74, 75, 76, 357, 359
 hydroxybutyrate 52
 hydroxycobalamin therapy 54, 55
 hydroxydaunorubicin 153, 154
 hydroxyurea 154, 155
 in essential thrombocythaemia 238
 megaklastic anaemia due to 56
 in myelofibrosis 239
 polycythaemia rubra vera 234–5
 in sickle cell anaemia 89–90
 hypercalcaemia 217, 218, 219, 223
 hypereosinophilic syndrome 103–4
 hyperferritinaemia 42
 hyperglobulinaemia 287
 hyperhomocysteinaemia 304, 307–8
 hypersplenism 126
 hyperuricaemia 149, 175, 177, 207, 219, 233, 234
 hyperviscosity syndrome 211, 218, 224, 225, 226–8
 hypogammaglobulinaemia 101
 hypoplastic anaemia 70, 241–6, 326, 328
 hyposplenism 126–7, 127
 hypothyroidism 326

 ibritumomab 154
 idarubicin 154
 idiopathic cold haemagglutinin syndrome 67
 idiopathic thrombocytopenic purpura 66, 118, 281–3, 287–8, 353–4
 idraparinux 316
 IF 45, 46, 49
 IGH gene 216
 IGVH gene 190, 191
 ileal resection 55
 imatinib 104, 154, 155, 177–9
 imipenem 104
 imipramine 104
 immobility and thrombosis 308
 immune paresis 218
 immune response 115–17, 124
 immunoblasts 206
 immunocytes 94
 immunodeficiency 119, 120, 121, 189
 immunoglobulin superfamily 11
 immunoglobulins 11, 94, 109–12, 218
 in CLL 189, 190
 genes 112–13, 190, 208
 in phagocytosis 99
 structure 111, 112
 therapy 67, 192, 283, 350
 immunohistochemistry 141
 immunohistology 141
 immunosecretory disorders 369
 immunosuppressive therapy 67, 283
 inborn errors of metabolism 331–3, 334
 incidental thrombocytopenia of pregnancy 353
 industrial solvents 129
 infancy and childhood
 CML 180–1
 haemoglobin levels 20, 21
 haemopoiesis 1
 iron deficiency 34
 iron requirements 33
 see also neonates
 infection 326–31
 adjuvant growth factor therapy 99
 in aplastic anaemia 244
 in β -thalassaemia major 80
 in CLL 189
 following stem cell transplantation 258–60, 261
 haematological malignancies associated 130–1
 haemolytic anaemia due to 69–70
 in multiple myeloma 217, 223
 in myelodysplasia 182, 183
 in neutropenia 105–6
 in non-Hodgkin's lymphoma 204
 prevention in hyposplenism 127
 prevention and treatment in haematological malignancies 149–52
 purpura in 279
 in sickle cell anaemia 89
 and thrombocytopenia 283
 transmission by blood transfusion 80, 84, 341–3, 346
 infectious mononucleosis 67, 68, 117–19, 120, 125
 inferior vena cava filter 316
 infertility 52, 149
 inflammatory bowel disease 309
 integrins 11
 interferon- α therapy 154, 156, 179, 194, 223, 235, 238
 interferon- γ 6, 116
 interleukins
 in granulopoiesis 97
 in haemopoiesis 6, 6, 97, 98
 IL-1 6, 97, 98, 115
 IL-6 31, 217
 synthesis by T cells 116
 interphase 9
 interstitial pneumonitis 260
 intracerebral haemorrhage 291
 intrinsic factor 45, 46, 49
 intrinsic Xase 271
 IREs 30
 iron
 absorption 32–3
 daily cycle 29
 deficiency 28, 30, 33–9, 41, 323, 352–3
 dietary 31–2
 distribution and transport within the body 28–30
 and erythropoiesis 15
 nutritional and metabolic aspects 28–33
 overload 30, 31, 41–3, 78–9, 80, 81–2, 186, 346
 requirements 33
 serum 36, 39, 365
 therapy 34–5, 36, 38–9, 70, 353
 iron chelation therapy 78, 79, 83, 186
 iron dextran 38
 iron regulatory protein 30
 iron response elements 30
 iron sorbitol 38
 IRP 30
 isochromosomes 134
 ITP 66, 118, 281–3, 287–8, 353–4

 Jagged proteins 4
 JAK2 gene/JAK2 protein 135, 230, 231, 232, 233, 237, 238
 JAK/STAT pathway 6–7, 8
 jaundice 22
 in cold autoimmune haemolytic anaemia 68
 in haemolytic anaemias 58
 in hereditary spherocytosis 62
 in infectious mononucleosis 118
 in megaloblastic anaemia 50
 in pyruvate kinase deficiency 66
 jejunal resection and folate deficiency 49
 juvenile myelomonocytic leukaemia 187

 kala-azar 331
 kallikrein 271
 Kaposi's sarcoma 131
 karyotype 134, 138, 139
 keratinocyte growth factor, recombinant human 150
 kernicterus 357
 kidney
 erythropoietin production 12, 15
 in Fanconi's anaemia 241, 243
 in multiple myeloma 217, 219, 221
 polycystic 324
 see also renal failure
 Klebsiella 149
 Kleihauer test 357
 Klinefelter's syndrome 129
 koilonychia 22, 34
 Kostmann's syndrome 104–5

 lactate dehydrogenase 26, 52, 198, 206–7, 233, 284–5
 lactoferrin 95, 100, 101
 Langerhans' cell histiocytosis 106
 large B-cell lymphoma 194
 large granular lymphocytic leukaemia 195
 lazy leucocyte syndrome 101
 LCH 106
 LDH 26, 52, 198, 206–7, 233, 284–5
 lead poisoning 41, 70
 leg ulcers 22
 leishmaniasis 125, 331, 342
 lepirudin 313
 Letterer–Siwe disease 106
 leucocytes see white blood cells
 leucocytosis
 basophil 104, 175
 in CML 175
 eosinophilic 103–4, 198
 leukaemoid reaction 102–3
 neutrophil 101–2, 198, 233

- leucodepletion of blood products 347
 leucopenia 126, 244
 leucopheresis in hyperviscosity syndrome 228
 leukaemia
 acute 157–8
 chronic 174
 classification 157
 see also specific disorders
 leukaemia/lymphoma syndromes 188, 194–5
 leukaemoid reaction 102–3, 323, 326, 327
 LGL-L 195
 ligands 11
 lipopolysaccharide and dendritic cell maturation 115
 livedo reticularis 309
 liver
 in β -thalassaemia major 77–8, 79, 80
 disease 297–8, 325–6
 erythropoietin production 12
 fetal 1
 haemopoiesis 1, 124
 in iron overload 41, 42–3, 79, 80
 veno-occlusive disease following stem cell transplantation 261
 LM02 10
 loiasis 331, 332
 lorazepam in chemotherapy 149
 LPS and dendritic cell maturation 115
 Luebering–Rapoport shunt 18, 19
 lumbar puncture in ALL 159
 lupus anticoagulant 300–1, 309, 324
 Lyme disease 342
 lymph nodes 116–17
 in non-Hodgkin's lymphoma 204, 205, 206
 as secondary lymphoid organ 110
 lymphadenopathy 197
 in ALL 159, 162
 angioidimmunoblastic 214
 in Burkitt's lymphoma 213
 in CLL 189
 differential diagnosis 121–2
 in diffuse large B-cell lymphoma 213
 in follicular lymphoma 209
 in Hodgkin's lymphoma 197–8
 in infectious mononucleosis 118
 in mantle cell lymphoma 212
 in non-Hodgkin's lymphoma 203–4
 lymphoblastic lymphomas 214
 lymphoblasts 157
 lymphocytes 94, 95, 108–9, 110, 365
 see also specific cells
 lymphocytic lymphomas 205, 206, 207, 210–11
 lymphocytosis 117–19, 126, 188, 189
 lymphoma 197
 distinction from chronic lymphoid leukaemias 188
 leukaemic phase 197
 and red cell aplasia 246, 321
 splenomegaly in 125
 therapy 99
 lymphoma/leukaemia syndromes 188, 194–5
 lymphopenia 119, 198
 lymphoplasmacytoid lymphomas 205, 211, 223–4
 lysosomes 267
 lysozyme 95, 100
 M-CSF 6, 7, 97, 98
 M proteins 216, 217, 218, 224–5, 329
 macrocytes 23, 24, 52
 macrophage colony-stimulating factor 6, 7, 97, 98
 macrophages 3, 99, 100, 124
 magnetic resonance imaging *see* MRI
 magnetic resonance pulmonary angiography 311
 major histocompatibility complex 254
 malaria 25
 and Burkitt's lymphoma 131
 and DIC 299
 haemolysis in 69, 330
 in hyposplenism 127
 resistance to 64, 74
 splenomegaly in 125
 transmission by blood transfusion 342, 346
 and tropical splenomegaly syndrome 125–6, 330
 Maloprim in glucose-6-phosphate dehydrogenase deficiency 64
 MALT lymphoma 131, 133, 207, 212
 mantle cell lymphoma 9, 194, 205, 211–12
 genetics 133, 207, 208
 histology 206
 immunophenotype 190, 207
 mantle zone cells 206
 MAPK pathway 6, 7, 8
 march haemoglobinuria 69
 marginal zone lymphomas 205, 212
 massive transfusion syndrome 286, 301
 mast cell diseases 368
 mast cells 96, 239
 mastocytosis, systemic 239–40
 May–Hegglin anomaly 101, 102
 MCH *see* mean cell haemoglobin
 MCV *see* mean corpuscular volume
 MDR 132
 mean cell haemoglobin (MCH) 16, 365
 in anaemia 21, 28
 in anaemia of chronic disorders 39
 in pregnancy 352–3
 mean cell haemoglobin concentration 16, 28, 365
 mean corpuscular volume (MCV) 16, 365
 in anaemia 21, 23, 28
 in anaemia of chronic disorders 39
 in megaloblastic anaemia 52
 neonatal 354
 in pregnancy 352
 mediastinal compression in ALL 159, 161, 163
 megakaryoblasts 265
 megakaryocytes 2, 265, 266, 281
 megaloblastic anaemias 44–56, 101, 102, 353
 MEK 1/2 7
 melagatran 316
 melanin pigmentation 50, 52, 79
 melphalan 129, 153
 membrane phospholipid 266
 memory B cells 108, 115, 117, 205
 meningeal syndrome in ALL 159
 menorrhagia 35, 37, 282, 326
 menstruation 33, 35, 37, 326
 mepacrine 104
 mercaptopurine 56, 153
 mesenchymal stem cells 4
 metamyelocytes 52, 53, 55, 95, 96, 101, 326
 methaemalbumin 60, 61
 methaemoglobinemia 18, 74
 methionine synthase 46
 methotrexate 48, 56, 153, 244
 methylolopa 66, 67, 68
 methylene tetrahydrofolate reductase 307
 methylmalonic acid 46, 49, 52
 methylmalonyl CoA 46
 methyltetrahydrofolate (methyl THF) 46, 47–8
 metoclopramide 149
 MGUS 216, 224–5
 MHC 254
 mianserin 104
 microcytes 23, 24
 microRNAs 137
 microspherocytes 24, 59, 62, 67
 minimal residual disease 144–6, 165
 minor histocompatibility antigens 254–5
 MIP-1 α 100
 miRNAs 137
 mitogen activated protein kinase pathway 6, 7, 8
 mitosis 9
 mitoxantrone 153, 154
 MLL gene/MLL protein 136, 162
 monoblasts 96
 monoclonal band proteins 216, 217, 218, 224–5, 329
 monoclonal gammopathy of undetermined significance 216, 224–5
 monocytes 94, 95–6, 99
 blood count 94, 365
 differentiation 2, 96
 functional defects 101
 functions 99–101
 monocytopenia in hairy cell leukaemia 193
 monocytosis 104
 monospot test 118, 120
 Mpl in polycythaemia rubra vera 233
 MRD 144–6, 165
 MRI
 deep vein thrombosis 311
 Hodgkin's lymphoma 200
 multiple myeloma 219
 non-Hodgkin's lymphoma 208, 210
 sickle cell anaemia 86
 spleen 125
 mucin-secreting adenocarcinoma 321, 322, 323
 mucopolysaccharidoses 101
 mucosa-associated lymphoid tissue lymphoma 131, 133, 207, 212
 mucositis 149, 150
 mucous membrane pallor 22, 58
 multidrug resistance 132
 multiple myeloma 121, 149, 205, 216–23, 227
 muscle
 haematoma 291
 weakness in Hodgkin's lymphoma 198
 mustine 129
 MYC gene 135, 137, 138
 Mycoplasma 67, 327
 mycosis fungoides 204, 214
 myeloblasts 95, 96, 102, 157, 183
 myelocytes 95, 96, 101, 102
 myelodysplastic syndromes/myelodysplasia 40, 182–6, 321
 and aplastic anaemia 244, 245
 classification 183, 185, 366
 genetics 133, 183, 185
 in HIV infection 328
 hypoplastic 245
 treatment 98–9, 185–6
 myelodysplastic/myeloproliferative diseases 186–7, 287, 366
 myelofibrosis 125, 230, 234, 235, 238–9
 myeloid:erythroid ratio 24, 27, 96
 myeloid growth factors 97
 myeloid metaplasia 238
 myelomatosis 121, 149, 205, 216–23, 227
 myeloperoxidase 95
 deficiency 101
 myelopoiesis 1
 myeloproliferative disorders 133, 135, 230–40, 366

- myelosclerosis 125, 230, 234, 235, 238–9
 myoglobin 29, 30
 myxoedema 104, 326
- nabilone 149
 NADH 18
 NADPH 18, 63, 100
 nail bed in anaemia 22
 nails in iron deficiency 34
 NAP score 102, 103, 175, 233, 326
 natural killer cells 2, 108–9
 nausea in chemotherapy 148–9
Neisseria meningitidis 127
 neonates
 anaemia 355
 coagulation tests 356
 G6PD deficiency 65
 haemolytic disease 68, 356–8, 359
 haemorrhagic disease 296, 297
 normal blood count 354
 polycythaemia 355–6
 nephrocalcinosis 221
 nephrotic syndrome 325
 neural tube defects 51
 neutropenia 104
 in ALL 159, 161
 in anaemia 23
 autoimmune 105
 causes 104
 clinical features 105
 congenital 104–5
 cyclical 105
 diagnosis 105
 drug-induced 104, 105
 in haematological malignancy 149
 in HIV infection 328
 idiopathic benign 105
 in large granular lymphocytic leukaemia 195
 management 99, 105–6
 in multiple myeloma 217, 218
 in non-Hodgkin's lymphoma 204, 206
 prevention and treatment of infection 150–1
 neutrophil alkaline phosphatase score 102, 103, 175, 233, 326
 neutrophil count 23, 94
 neutrophils 94–5, 365
 band/stab/juvenile 95, 96
 differentiation 2, 95, 96
 functional defects 101
 functions 99–101
 leucocytosis (neutrophilia) 101–2, 198, 233
 morphological abnormalities 101, 102
 in the neonate 354
 segmented 96
 new variant Creutzfeldt–Jacob disease 343, 346
 NF-E2 73
 NHL 129, 131, 133, 203–15, 323, 329
 nicotinamide adenine dinucleotide 18
 nicotinamide adenine dinucleotide phosphate 18, 63, 100
 Niemann–Pick disease 333
 night sweats 175, 204, 233
 nitric oxide 101, 270, 275
 nitrosoureas 129, 153, 242
 NK cells 2, 108–9
 NO 101, 270, 275
 nocturnal enuresis 88
 non-Hodgkin's lymphoma 129, 131, 133, 203–15, 323, 329
 normoblasts 12, 13, 14, 25
 in anaemia of chronic disorders 39
 in iron deficiency 36
 in megaloblastic anaemia 44, 52, 53
- Notch 4, 10
 nuclear weapons 130
nucleophosmin gene 135, 167
 nutritional support in chemotherapy 149
- oestrogen therapy and thrombosis 309
 oncogenes 132, 133
 ondansetron 148
 operative haemorrhage 291
 opsonization 99–100, 113
 oral cavity disorders 105, 169, 170, 204
 oral contraceptives 309
 orotic aciduria 56
 Oroya fever 327
 osteomyelitis 88
 osteoporosis 80, 313
 oxygen exchange 17–18
- p53 9–10, 132, 135
 packed cell volume 16, 365
 PAI 275
 pain
 alcohol-induced 198
 in haematological malignancy 149, 217
 Palifermin® 150
 pamaquine 64
 pamidronate 223
 pancytopenia 23, 104, 241
 following stem cell transplantation 252
 in hairy cell leukaemia 193
 in malignant disease 322
 in myelodysplasia 182
 papilloedema, in ALL 159
 Pappenheimer bodies 24, 25, 124, 126
 paraproteinaemia 112, 216, 217
 paraproteins 216, 217, 218, 224–5, 329
 parietal cell antibody 49
 paroxysmal cold haemoglobinuria 68, 328
 paroxysmal nocturnal haemoglobinuria 58, 60, 70, 245, 309
 parvovirus infection 59, 246, 247, 328, 342
 Paterson–Kelly syndrome 34
 Paul–Bunnell test 118, 120
 PBSC transplantation *see* stem cell transplantation
 PCR 90, 91–2, 139, 141, 145–6
 PCV 16, 365
 PDGF 267, 270, 303
 PECAM-1 270
 PEL 131
 Pelger–Huët anomaly 101, 102, 183
 pencil cells 24, 35
 penicillamine 104
 penicillin 68
 pentose phosphate pathway 18, 20, 65
 peptic ulceration in polycythaemia rubra vera 233
 peripheral blood stem cell transplantation *see* stem cell transplantation
 pernicious anaemia 48–9, 53, 55, 321, 326
 Persantine (dipyridamole) 287, 317–18
 PET scanning
 combined with CT 209, 211
 Hodgkin's lymphoma 200
 non-Hodgkin's lymphoma 208–9, 210, 211
 spleen 125
 PF4 267
 PFA-100 277, 288, 302
 Ph chromosome 138, 155, 162, 174, 175–6
 phagocytes 2, 94–7, 98, 99–101
 phagocytosis 99–101
 phagosome 100
 phenoxymethylpenicillin 127
 phenylbutazone 104
 phenytoin 104
 Philadelphia chromosome 138, 155, 162, 174, 175–6
- phosphatidylinositol 3 kinase pathway 6, 7, 8, 231
 phosphatidylinositol glycan protein-A 70
 6-phosphogluconate 18, 20
 phospholipase A₂ 269
 phosphorus-32 therapy in polycythaemia rubra vera 235
 PI3 kinase pathway 6, 7, 8, 231
 pica in iron deficiency 34
 PIG-A 70
 pipobroman 235
 PIVKA factors 296
 placenta
 infarction 309
 premature separation 299
 plasma cell leukaemia 194, 223
 plasma cells 108, 109, 115, 117, 194, 205
 disorders 368
 in HIV infection 329
 in multiple myeloma 216, 217, 218
 plasma exchange 285
 plasma viscosity measurement 335
 plasma volume 365
 plasmacytoma, solitary 223
 plasmapheresis 224, 225, 228
 plasmin 274–5
 plasminogen 274–5
 plasminogen activator inhibitor 275
 platelet count
 in anaemia 23
 in bleeding disorders 276, 300
 in CML 175
 following splenectomy 127
 following stem cell transplantation 252, 253
 in haematological malignancy 147
 in Hodgkin's lymphoma 198
 in iron deficiency 36
 normal 265, 365
 in platelet disorders 287–8
 in polycythaemia rubra vera 233
 in pregnancy 353
 raised 237
 platelet-derived growth factor 267, 270, 303
 platelet factor-3 266, 269
 platelet function analysis-100 test 277, 288, 302
 platelets
 activation 268–9
 adhesion 266, 267, 268–9
 aggregation 269
 aggregometry 277, 288
 amplification 269
 antigens 268
 in atherosclerosis 303
 clot formation and retraction 269
 in CML 175
 differentiation 2
 disorders 278, 286–8
 functions 268–70
 granules 266–8, 269
 in haemostasis 265–9, 270, 275–6
 increased destruction 281–6
 lifespan 265, 282
 in malignant disease 322
 in multiple myeloma 217
 in myelodysplasia 183
 natural inhibitors of function 270
 procoagulant activity 269
 production 265
 failure 281
 release reaction 269
 in splenomegaly 285–6
 structure 265–8
 transfusion 147, 245, 260, 283, 288–9, 348, 349

- pleural effusion 198
 PLL 133, 190, 192–3
 Plummer–Vinson syndrome 34
 PML gene 135, 136, 138
 PML-RAR α fusion protein 135, 136, 138, 141, 156
Pneumocystis carinii pneumonia 151–2, 260
 PNH 58, 60, 70, 245, 309
 point mutations 135
 polyarteritis nodosa 69
 polycythaemia 13, 74, 226, 232, 236–7, 323, 355–6
 polycythaemia rubra vera 7, 13, 103, 230, 232–6, 309
 polyendocrine syndrome 48
 polymerase chain reaction 90, 91–2, 139, 141, 145–6
 polymorphs *see* neutrophils
 polymyalgia rheumatica 324
 portal hypertension 125, 126
 positron emission tomography *see* PET scanning
 post-capillary venules 109
 post-transplant lymphoproliferative disease 131, 151
 post-traumatic haemorrhage 291
 pregnancy 352
 anticoagulants in 313, 314, 354
 folate deficiency 49, 353
 folate supplementation 51, 55, 353
 haemostasis 354
 idiopathic thrombocytopenic purpura in 353–4
 iron deficiency 34–5, 37, 352–3
 iron requirements 33
 physiological anaemia 352
 and sickle cell anaemia 89
 thrombocytopenia 353–4
 thrombosis in 354
 vitamin B₁₂ deficiency 353
 pre-implantation diagnosis of haemoglobin disorders 92
 prenatal diagnosis
 haemoglobin disorders 90–2
 haemophilia 292
 priapism 88, 175
 prickle cells 66
 primaquine 64
 primary effusion lymphoma 131
 primary lymphoid organs 108, 110
 procabazine 129
 prochlorperazine 149
 progenitor cells 2–3
 progesterone therapy during chemotherapy 148
 prolymphocytes 193
 prolymphocytic leukaemia 133, 190, 192–3
 promonocytes 96
 promyelocytes 95, 96, 102
 promyelocytic leukaemia 143
 pronormoblasts 12, 13, 14
 prostacyclin
 intravenous therapy 319
 see also prostaglandins
 prostaglandins 239, 267, 269, 275
 protamine 313
 protein 4.1 19, 20
 protein 4.2 20
 protein C 272, 274, 275, 306, 350, 356
 deficiency 306
 recombinant human 300
 protein S 274, 307
Proteus 149
 prothrombin 266, 268, 269, 271
 allele G20210A 307
 prothrombin time 276, 300, 310, 356
 prothrombinase 269, 271
 proto-oncogenes 132
 protoporphyrin 41, 58
 pruritus 198, 233
 PRV 7, 13, 103, 230, 232–6, 309
 PRV-1 233
Pseudomonas 149, 342
 pseudo-Pelger cells 184
 pseudopolycythaemia 232, 236
 pseudotumours, haemophilic 291, 292
 pseudoxanthoma elasticum 279
 psychotropic drugs 242
 PT 276, 300, 310, 356
 PTLID 131, 151
 PU.1 5
 pulmonary embolism 305, 311
 pulmonary haemosiderosis 37
 pulmonary hypertension 88
 purine analogues 153, 154, 192
 purpura
 in CLL 189
 in infection 279
 in megaloblastic anaemia 50
 in myelodysplasia 182, 183
 in non-Hodgkin's lymphoma 204
 post-transfusion 283–4, 346
 senile 279
 steroid 280
 pyelonephritis in multiple myeloma 219, 221
 pyridoxal-6-phosphate 40
 pyridoxine 15, 16–17, 40
 pyrimethamine 48, 56
 pyrimidine analogues 153
 pyruvate kinase deficiency 65–6

 Q fever 342
 quinidine 68, 284
 quinine 284, 330

 RA 183, 185
 radiation exposure 130
 radiography
 ALL 159, 161, 163
 Hodgkin's lymphoma 200, 201
 multiple myeloma 218, 219, 220
 non-Hodgkin's lymphoma 208
 pulmonary embolism 311
 sickle cell anaemia 86
 radiotherapy
 ALL 164, 165, 166
 CLL 192
 follicular lymphoma 209
 growth factor therapy following 97
 Hodgkin's lymphoma 201
 immunodeficiency following 121
 late effects 188, 202
 leukaemia following 188
 multiple myeloma 223
 RANKL 218
 RANTES 100
 RAR α gene 135, 136, 138
 rare autosomal recessive syndrome 246
 RARS 40, 185
 RAS oncogenes 135, 183
 rasburicase 149
 Raynaud's phenomenon 224
 RCMD 185
 REAL classification 203
 receptors 11
 recombinases 113, 135
 recurrent abortion 309
 red cell count 16
 red cell fragmentation syndromes 69
 red cell indices 16, 21, 23, 35–6, 365
 red cells 12, 18
 alloimmunization 356
 in anaemia 23, 24, 25
 antigens and blood groups 337–41
 aplasia 246, 247, 321, 328
 basophilic stippling 25, 41
 development 12, 13, 14
 differentiation 2
 fragments 24, 59, 69, 299
 functions 15–16, 17–18
 haemoglobin content 16
 in haemolytic anaemias 59
 inclusions 23, 25
 lifespan 26, 27
 in malignant disease 322
 membrane 18–20
 metabolism 18, 19, 20
 defective 63–6
 normal destruction 58, 59
 in renal failure 324
 role of spleen 123–4
 rouleaux formation 218, 220, 324
 storage 347
 substitutes 347
 transfusion 147, 347, 348
 red man syndrome 195
 Reed–Sternberg cells 197, 198, 199
 refractory anaemia 183, 185
 refractory anaemia with excess blasts 185
 refractory anaemia with ring sideroblasts 40, 185
 refractory cytopenia with multilineage dysplasia 185
 refractory cytopenia with multilineage dysplasia and ring sideroblasts 185
 renal failure 217, 219, 221, 223, 324–5
 restrictive pneumonitis 261
 reticulocyte count 16, 23, 35, 354
 reticulocytes 12, 14, 25, 62, 67
 reticulocytosis 58, 59, 252
 reticuloendothelial system 99
 retina
 haemorrhage in anaemia 22
 in hyperviscosity syndrome 224, 225, 227
 in polycythaemia rubra vera 233
 in sickle cell anaemia 88
 Revised European–American Lymphoma classification 203
 rhesus blood group system 68, 338, 339, 340, 341, 356–8, 359
 rheumatoid arthritis 323–4
 rhombotin 10
 riboflavin 15
 rifampicin 68
 ring sideroblasts 40, 183, 184, 185
 rituximab 154, 156
 in CLL 192
 in cold autoimmune haemolytic anaemia 68
 in diffuse large B-cell lymphoma 213
 in follicular lymphoma 210
 in idiopathic thrombocytopenic purpura 283
 in mantle cell lymphoma 212
 in neutropenia 106
 potential mechanisms of action 212
 in prolymphocytic leukaemia 193
 in red cell aplasia 246
 in thrombotic thrombocytopenic purpura 285
 in Waldenström's macroglobulinaemia 224
 in warm autoimmune haemolytic anaemia 67
 RNA 115, 137
 Rocky Mountain spotted fever 342

- Rosai–Dorfman disease 107
 RS cells 197, 198, 199
 rubella 328
 Runx-1 (AML1 protein) 10, 162
- Salazopyrin 64, 70
Salmonella 342
 SAP scan 228
 SCF 4, 6, 7, 97
 schistosomiasis 125, 331
 Schwachman–Diamond syndrome 246
 SCL 10
 SCT *see* stem cell transplantation
 SCU-PA 317
 scurvy 279, 280
 SDF-1 4
 secondary lymphoid organs 108, 110, 116
 secondary myeloid leukaemia 133
 selectins 11
 septicaemia 69–70, 105
 serine proteases 270, 272, 273
 serotonin 267
Serratia 342
 serum amyloid P component scan 228
 serum free light chain assay 218
 severe combined immunodeficiency 121
 Sézary syndrome 195, 204, 214
Shigella 285
 shingles 151, 189
 sickle cell anaemia 74, 75, 82, 85–90, 91, 309
 sickle cell/C disease 90
 sickle cell trait 82, 90
 sickle cells 24, 88, 89
 sideroblastic anaemia 28, 39, 40–1
 siderotic granules 24, 25, 124, 126
 single-chain urokinase-type plasminogen activator 317
 sinus histiocytosis with massive lymphadenopathy 107
 skull bossing in β -thalassaemia major 78, 79
 smallpox 104
 smudge cells 189, 190
 snake bites 299
 solitary plasmacytoma 223
 South-East Asian ovalocytosis 63
 Southern blot analysis 139, 145
 spectrin 19, 20
 spherocytes 20, 61–3, 67
 spina bifida 51
 spinal cord, subacute combined degeneration 51
 spleen
 anatomy 123
 atrophy 124
 circulation 123, 124
 cords 123, 124
 fetal 1
 functions 123–4
 haemopoiesis 1, 124
 imaging 125
 in immune response 124
 irradiation in myelofibrosis 239
 marginal zone 123
 periarteriolar lymphatic sheath (PALS) 123
 perifollicular zone 123
 and red cell integrity 123–4
 red pulp 123, 124
 rupture 127
 as secondary lymphoid organ 110
 white pulp 123, 124
 splenectomy 127
 in β -thalassaemia major 83
 in CLL 192
 in hereditary spherocytosis 62–3
 in idiopathic thrombocytopenic purpura 283
 indications 127
 in myelofibrosis 239
 in neutropenia 106
 prevention of infection following 127
 in polymphocytic leukaemia 193
 in pyruvate kinase deficiency 66
 in tropical splenomegaly syndrome 126
 in warm autoimmune haemolytic anaemia 67
 splenic artery 123
 splenic lymphoma with villous lymphocytes 194, 212
 splenic marginal zone lymphoma 194, 212
 splenic sequestration in sickle cell anaemia 87
 splenomegaly 125
 in ALL 159
 in β -thalassaemia major 77–8
 causes 126
 in CLL 189
 in CML 125, 175
 in cold autoimmune haemolytic anaemia 68
 cryptogenic 125–6
 in Felty's syndrome 323
 in haemolytic anaemias 58
 in hereditary spherocytosis 62
 in Hodgkin's lymphoma 198
 in infectious mononucleosis 118
 massive 125, 238
 in myelofibrosis 125, 234, 238
 in non-Hodgkin's lymphoma 204
 platelet pooling 285–6
 in polycythaemia rubra vera 233, 234
 in polymphocytic leukaemia 193
 in sickle cell anaemia 88
 tropical splenomegaly syndrome 125–6
Staphylococcus 105, 149, 150–1, 342
 STAT transcription factors 6–7, 8, 231
 stem cell factor 4, 6, 7, 97
 stem cell transplantation (SCT)
 in ALL 165
 allogeneic 249, 250, 251, 254–61
 in AML 171
 in aplastic anaemia 245–6
 autologous 249, 250, 251, 253–4
 in β -thalassaemia major 84
 in CLL 192
 in CML 179–80
 complications 256–61
 engraftment and immunity following 252–3
 failure 261
 in Fanconi's anaemia 244
 in follicular lymphoma 210
 and gene therapy 262–3
 growth factor therapy following 97, 98
 haploidentical 255
 HLA typing 254–5, 256
 in idiopathic thrombocytopenic purpura 283
 immunodeficiency following 121
 indications 251
 in mantle cell lymphoma 212
 in multiple myeloma 220
 in myelodysplastic syndromes 186
 non-myeloablative conditioning 252, 262
 patient conditioning 250, 252, 262
 potential donors 251
 principles 249–53
 relapse 261
 in sickle cell anaemia 90
 stem cell collection and processing 249, 251, 252
 syngeneic 249, 251
 umbilical cord blood 261
 use of growth factors in 99
 viral infection following 151
 stem cells 134–5, 249, 251, 252
 haemopoietic 1–3, 4–5, 108
 mesenchymal 4
 stercobilin 58
 stercobilinogen 58, 59
 stomatocytes 24
Streptococcus pneumoniae 127
 streptokinase 275, 317
 stroke in sickle cell anaemia 87
 stromal-derived factor-1 4
 subacute combined degeneration of the cord 51
 succinic dehydrogenase 30
 succinyl CoA 46
 sulfanilamide 64
 sulfasalazine 49, 104
 sulfapyrazone 318
 sulphaemoglobinaemia 18
 sulphonamides 242
 superior vena cava obstruction 198
 superoxide 100
 syphilis 342, 346
 systemic lupus erythematosus 66, 246, 282, 300, 309, 323–4
- T-cell lymphomas 214–15
 T-cell neoplasms 367–8
 T-cell receptors 11, 108, 113, 114, 115–16, 135, 139, 208
 T lymphocytes 108, 115, 116
 antigen presentation 99, 100
 circulation 109
 differentiation 2
 functional aspects 111
 HLA-restricted 115
 in immune response 115–17
 in infectious mononucleosis 118, 119
 naïve/virgin 115
 TAFI 275
 Tal-1 10
 target cells 24, 35, 80, 81, 88, 89, 325, 326
 TBI 250, 252, 261
 TC 45–6, 233
 TCRs 11, 108, 113, 114, 115–16, 135, 139, 208
 TdT 113, 159, 160
 tear drop poikilocytes 24, 238, 239
 TEG 301–2
 Tel 10
 TEL-AML1 gene/TEL-AML1 fusion protein 135, 162
 telomerase 134–5
 telomeres 134–5
 temporal arteritis 324
 tenase 269
 TERC gene 242
 terminal deoxynucleotidyl transferase 113, 159, 160
 tetrahydrofolate 47, 48, 153
 TFPI 271, 273, 275
 TGF- β 6
 thalassaemias 28, 39, 40, 72, 74–85, 90, 92
 thalidomide 186, 220, 222–3, 239
 THF 47, 48, 153
 thiamine and erythropoiesis 15
 thioridazine 104
 thrombasthenia 266, 287
 thrombi 303
 thrombin 266, 268, 269, 270–3, 276
 inactivation 273
 intravascular 299
 thrombin-activated fibrinolysis inhibitor 275

- thrombin time 276, 277, 300, 356
 thrombocytopenia 280–6
 in ALL 159, 162
 in AML 169
 in anaemia 23
 in antiphospholipid syndrome 309
 in aplastic anaemia 244
 causes 281
 in CLL 189
 drug-induced 280, 281, 284, 313, 314
 in HIV infection 328
 of hypertensive disorders 353
 and infection 283
 in infection 327
 in infectious mononucleosis 118
 in liver disease 326
 in malaria 330
 in malignant disease 323
 in multiple myeloma 217, 218
 in non-Hodgkin's lymphoma 204, 206
 in pregnancy 353–4
 in renal failure 325
 in tropical splenomegaly syndrome 126
 thromboelastography 301–2
 thrombolytic agents 275, 317
 thrombomodulin 274, 275
 thrombophilia 305–10
 thrombopoiesis 1
 thrombopoietin 6, 265
 thrombosis 273
 arterial 303–4, 309
 cerebral 303
 in essential thrombocythaemia 237, 309
 neonatal 356
 in paroxysmal nocturnal
 haemoglobinuria 70, 309
 in polycythaemia rubra vera 233, 309
 in pregnancy 354
 venous 304–19, 323
 thrombospondin 4
 thrombotic thrombocytopenic purpura 69,
 284–5, 286, 328
 thromboxane A₂ 267, 269, 275, 287
 thymoma 246, 321
 thymus
 aplasia 121
 as primary lymphoid organ 108, 110
 thyroxine deficiency 326
 TIAs 303
 TIBC 36, 37, 39, 365
 ticlopidine 318
 tioguanine 153
 tirofiban 287, 318, 319
 tissue factor 268, 270, 271, 272, 275, 298
 tissue factor pathway inhibitor 271, 273,
 275
 tissue plasminogen activator 274–5, 275,
 317
 TNF 6, 97, 98, 115, 116
 tolbutamide 104
 tositumomab 154
 total-body irradiation 250, 252, 261
 total iron-binding capacity 36, 37, 39, 365
 total plasma volume in polycythaemia
 232
 total red cell volume in polycythaemia 232,
 233
 toxoplasmosis 330, 342, 346
 tPA 274–5, 275, 317
 trabecular arteries 123, 124
 trabecular vein 124
 TRALI 346
 tranexamic acid 148, 280
 transcobalamin 45–6, 233
 transcription factors 5, 8, 10, 73
 transferrin 28–9, 30, 31, 32, 33, 58
 transferrin receptors 13, 28–9, 30–1, 33,
 36–7, 39, 42, 43
 transforming growth factor- β 6
 transfusion transmitted virus 342
 transfusion-related acute lung injury 346
 transient ischaemic attacks 303
 trimethoprim 48, 56
 trisomy 12 136
 tropical splenomegaly syndrome 125–6,
 330
 tropical sprue 49
 trypanosomiasis 331
 TT 276, 277, 300, 356
 TTP 69, 284–5, 286, 328
 TTV 342
 tuberculosis 309, 327
 tumour lysis syndrome 149
 tumour necrosis factors 6, 97, 98, 115, 116
 tumour-suppressor genes 132, 133
 UIBC 37
 ulcerative colitis, basophilia in 104
 ultrasound
 deep vein thrombosis 305, 310
 spleen 125
 umbilical cord blood, stem cell
 transplantation 261
 unsaturated serum iron-binding capacity
 37
 uraemia 287
 urobilin 58
 urobilinogen 58, 59
 urokinase 275
 urticaria pigmentosa 239
 vaccinations in hyposplenism 127
 valaciclovir 151
 varicella zoster infection 151, 258, 260
 vascular endothelial growth factor 13
 vasculitis 69
 vasoconstriction 275
 vegan diet 48
 VEGF 13
 Velcade 213, 223
 venesection 43, 227, 234
 venous stasis and thrombosis 308
 ventilation-perfusion scintigraphy in
 pulmonary embolism 311
 vinblastine 154
 vincristine 153, 154
 vindesine 154
 viral infection 328–9
 in β -thalassaemia major 80
 in CLL 189
 following stem cell transplantation
 258–61
 haematological malignancies associated
 131
 in haematological malignancy 151
 in neutropenia 105–6
 purpura in 279
 and thrombocytopenia 283
 transmission by blood transfusion 80,
 341–2, 346
 visual disorders 175, 233
 vitamin B₆ 15, 16–17, 40
 vitamin B₁₂ 44
 absorption 45, 46
 tests 52–3, 54
 biochemical function 46
 deficiency 47, 48
 and cardiovascular disease 56
 causes 48–9
 clinical features 49–50
 diagnosis 52–4
 effects 50, 51–2
 with iron deficiency 35–6
 laboratory findings 52
 and neural tube defects 51
 neuropathy due to 51
 in pregnancy 353
 treatment 54–6
 and erythropoiesis 15
 forms 45
 malabsorption 49
 metabolic abnormalities 56
 nutritional aspects 45
 serum 52, 365
 structure 45
 supplements 54–5
 transport 45–6
 vitamin C 15, 83, 279, 280
 vitamin E 15
 vitamin K 148, 273, 296–7
 vomiting in chemotherapy 148–9
 von Willebrand disease 37, 287, 288, 294,
 295–6
 von Willebrand factor (VWF) 266, 267, 269,
 275, 285, 295
 assay 277
 ultra large 284
 Waldenström's macroglobulinaemia 211,
 223–4, 225, 226
 Waldeyer's ring 204
 warfarin 313–14
 discontinuation before chemotherapy 148
 drug interactions 314, 315
 during surgery 315
 overdose 315, 316
 in paroxysmal nocturnal
 haemoglobinuria 70
 as vitamin K antagonist 296–7
 Weibel–Palade bodies 269
 weight loss 149, 175, 198, 204
 white blood cells 94–107, 108–22, 322, 323,
 327, 365
 Wilson's disease 70, 325
 Wiskott–Aldrich syndrome 129
Wuchereria bancrofti 332
 X-linked agammaglobulinaemia 121
 ximelagatran 316
Yersinia enterocolitica 80, 342
 yolk sac 1
 ZAP-70 190, 191
 zidovudine 56
 Zieve's syndrome 325, 326
 zoledronic acid 223


Essential Best Seller

Postgraduate Haematology, 5e

Edited by: A. Victor Hoffbrand, Daniel Catovsky, Edward G.D. Tuddenham

A practical, readable text which will give trainees, residents and practising hematologists up-to-date knowledge of the pathogenesis, clinical and laboratory features and management of blood disorders.

ISBN: 1405108215



Postgraduate Haematology is ideal for:

- Trainees and residents in hematology
- Hematologists in practice

Why Buy This Book?

A well established and respected review of hematology

- Practical and readable text
- Essential information for everyday use as well as the scientific background
- Up-to-date knowledge of the pathogenesis, clinical and laboratory features and management of blood disorders
- Complete revision of all chapters and the addition of new chapters to reflect latest advances in the speciality

For further information, a free sample chapter or to place your order online please visit:

www.blackwellhematology.com

Alternatively, you can place your order by calling:

Toll Free Phone: 800-216-2522 (The Americas)

Direct Sales Tel No: +44 (0)1235 465500 (Rest of World)



**Blackwell
Publishing**

www.blackwellhematology.com

www.blackwellhematology.com

